


Table 27.20  
Cancer of the Urinary Bladder (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2009-2013

Females

State	Rate	SE	Rank(CI) <sup>d</sup>	PD
TOTAL U.S.	2.17	0.01		
High Five States				
D.C.	3.07	0.43	01(01,40)	41.45 <sup>c</sup>
Delaware	2.96	0.32	02(01,34)	36.53 <sup>c</sup>
Vermont	2.93	0.38	03(01,42)	35.06 <sup>c</sup>
Maine	2.78	0.24	04(01,32)	27.90 <sup>c</sup>
Massachusetts	2.73	0.11	05(01,16)	25.72 <sup>c</sup>
Low Five States				
New Mexico	1.78	0.17	47(17,50)	-18.08 <sup>c</sup>
Utah	1.77	0.18	48(17,50)	-18.36 <sup>c</sup>
Alabama	1.77	0.11	49(31,50)	-18.56 <sup>c</sup>
South Dakota	1.76	0.25	50(07,51)	-19.15 <sup>c</sup>
Hawaii	0.98	0.14	51(50,51)	-54.68 <sup>c</sup>

State	Rate	SE	Rank(CI) <sup>d</sup>	PD	State	Rate	SE	Rank(CI) <sup>d</sup>	PD
Alabama	1.77	0.11	49(31,50)	-18.56 <sup>c</sup>	Montana	2.00	0.25	36(04,50)	-7.81
Alaska	1.94	0.43	40(01,51)	-10.85	Nebraska	2.01	0.19	35(07,50)	-7.33
Arizona	1.81	0.10	44(31,50)	-16.59 <sup>c</sup>	Nevada	2.54	0.19	08(01,37)	16.84 <sup>c</sup>
Arkansas	2.21	0.15	23(05,45)	1.99	New Hampshire	2.18	0.23	28(02,50)	0.23
California	1.95	0.04	39(30,45)	-9.95	New Jersey	2.50	0.09	11(03,25)	15.03 <sup>c</sup>
Colorado	1.88	0.12	41(21,50)	-13.21	New Mexico	1.78	0.17	47(17,50)	-18.08 <sup>c</sup>
Connecticut	2.53	0.14	09(02,32)	16.76 <sup>c</sup>	New York	2.19	0.06	26(14,37)	1.07
Delaware	2.96	0.32	02(01,34)	36.53 <sup>c</sup>	North Carolina	2.20	0.09	25(11,40)	1.34
D.C.	3.07	0.43	01(01,40)	41.45 <sup>c</sup>	North Dakota	1.87	0.29	42(04,51)	-14.04
Florida	2.04	0.05	32(24,43)	-6.21	Ohio	2.52	0.08	10(03,22)	16.16 <sup>c</sup>
Georgia	2.20	0.09	24(10,40)	1.47	Oklahoma	2.27	0.14	18(05,43)	4.68
Hawaii	0.98	0.14	51(50,51)	-54.68 <sup>c</sup>	Oregon	2.14	0.13	30(09,47)	-1.47
Idaho	1.81	0.20	45(12,50)	-16.63 <sup>c</sup>	Pennsylvania	2.31	0.07	17(09,33)	6.33
Illinois	2.25	0.08	22(10,37)	3.46	Rhode Island	2.67	0.27	06(01,42)	22.92 <sup>c</sup>
Indiana	2.35	0.11	16(05,36)	8.45	South Carolina	2.25	0.13	21(06,42)	3.79
Iowa	2.03	0.14	33(12,49)	-6.71	South Dakota	1.76	0.25	50(07,51)	-19.15 <sup>c</sup>
Kansas	2.02	0.15	34(10,50)	-6.92	Tennessee	1.99	0.10	38(18,49)	-8.52
Kentucky	2.38	0.13	15(03,37)	9.57	Texas	1.81	0.05	46(37,50)	-16.82 <sup>c</sup>
Louisiana	2.27	0.13	19(05,42)	4.40	Utah	1.77	0.18	48(17,50)	-18.36 <sup>c</sup>
Maine	2.78	0.24	04(01,32)	27.90 <sup>c</sup>	Vermont	2.93	0.38	03(01,42)	35.06 <sup>c</sup>
Maryland	2.47	0.12	13(03,32)	13.79	Virginia	2.19	0.10	27(10,42)	0.93
Massachusetts	2.73	0.11	05(01,16)	25.72 <sup>c</sup>	Washington	2.38	0.11	14(04,34)	9.80
Michigan	2.48	0.09	12(04,25)	14.14	West Virginia	2.61	0.20	07(01,36)	20.21 <sup>c</sup>
Minnesota	1.99	0.11	37(17,49)	-8.18	Wisconsin	2.16	0.11	29(10,44)	-0.38
Mississippi	1.86	0.14	43(19,50)	-14.23	Wyoming	2.12	0.37	31(01,50)	-2.19
Missouri	2.26	0.11	20(07,39)	4.32					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p < .0002$ ).

<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.

<sup>d</sup> Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.