

Table 28.8

5-Year Relative Survival (Percent)  
By Selected Primary Cancer Site and Year of Diagnosis  
All Races, Males and Females

Site	Ages 0-14										
	1975-1977	1978-1980	1981-1983	1984-1986	1987-1989	1990-1992	1993-1995	1996-1998	1999-2001	2002-2005	2006-2012
All Sites											
All Races	58.0	62.4	67.0	68.1	71.6	75.8	77.4	79.2	80.8	82.6	84.5 <sup>c</sup>
Whites	57.9	63.0	67.7	69.8	72.4	76.8	78.3	80.6	81.8	84.5	85.3 <sup>c</sup>
Blacks	57.3	57.7	62.2	57.0	65.3	70.8	73.3	75.6	73.9	72.7	82.1 <sup>c</sup>
Bone & Joint	49.9 <sup>d</sup>	47.8	56.8 <sup>d</sup>	57.3 <sup>d</sup>	66.8 <sup>d</sup>	67.4	74.1	70.3	70.0	78.4	78.4 <sup>c</sup>
Brain & ONS	56.9	57.7	56.7	61.7	64.3	64.4	70.7	75.2	74.0	75.5	74.7 <sup>c</sup>
Hodgkin lymphoma	80.9	86.8	88.1	89.9	87.1	96.8	94.6	96.1	94.4	97.2	97.7 <sup>c</sup>
Leukemia	49.7	58.0	62.9	63.6	71.0	75.5	75.9	80.3	82.8	85.7	88.0 <sup>c</sup>
Acute lymphocytic	57.2	65.7	71.3	72.2	77.7	83.1	83.9	86.9	88.5	90.8	92.3 <sup>c</sup>
Acute myeloid	18.8	25.8	26.7 <sup>d</sup>	30.6 <sup>d</sup>	37.1 <sup>d</sup>	42.2	40.6 <sup>d</sup>	48.7	58.6	63.3	66.8 <sup>c</sup>
Neuroblastoma <sup>a</sup>	52.5	56.6	54.5	52.3	63.2	76.0	66.8	66.5	72.1	72.2	77.3 <sup>c</sup>
Non-Hodgkin lymphoma	43.2	52.7	66.9	69.8	70.7	76.9	80.7	83.2	89.8	83.9	90.8 <sup>c</sup>
Soft tissue	61.3	74.2	69.2	72.9	66.4	79.8	76.7	70.5	76.9	82.5	81.4 <sup>c</sup>
Wilms' tumor <sup>b</sup>	73.1	79.0	86.7	90.7	92.2	91.9	91.7	91.5	93.8	90.8	93.0 <sup>c</sup>

Site	Ages 0-19										
	1975-1977	1978-1980	1981-1983	1984-1986	1987-1989	1990-1992	1993-1995	1996-1998	1999-2001	2002-2005	2006-2012
All Sites											
All Races	61.5	65.2	68.3	70.5	73.5	76.3	77.8	79.8	80.3	82.7	84.7 <sup>c</sup>
Whites	61.3	65.8	69.2	72.4	74.9	77.6	78.5	81.1	81.4	84.3	85.8 <sup>c</sup>
Blacks	59.5	60.3	62.5	58.5	64.4	68.6	74.2	73.9	74.6	73.1	80.5 <sup>c</sup>
Bone & Joint	50.4	48.1	51.2	56.1	64.0	68.8	68.9	67.0	66.7	75.0	74.5 <sup>c</sup>
Brain & ONS	58.9	58.1	58.4	63.9	65.9	66.4	71.5	76.0	74.4	75.5	75.1 <sup>c</sup>
Hodgkin lymphoma	86.2	88.3	85.5	90.6	88.7	94.2	93.9	95.2	95.0	96.6	97.3 <sup>c</sup>
Leukemia	44.9	53.2	58.0	60.3	67.6	71.3	71.9	76.6	77.7	82.2	86.1 <sup>c</sup>
Acute lymphocytic	53.8	61.9	67.1	69.9	75.0	79.8	81.4	84.3	84.8	87.8	91.0 <sup>c</sup>
Acute myeloid	18.7	26.2	26.4	31.4	37.6	41.9	38.6	46.3	51.9	61.5	66.2 <sup>c</sup>
Neuroblastoma <sup>a</sup>	53.1	56.7	53.4	52.5	61.7	75.7	66.8	66.7	71.9	72.2	76.8 <sup>c</sup>
Non-Hodgkin lymphoma	44.6	53.4	63.6	67.6	70.4	72.7	78.0	81.0	84.3	83.9	88.0 <sup>c</sup>
Soft tissue	64.8	68.9	68.2	72.6	68.0	69.7	73.8	72.3	73.0	77.0	78.2 <sup>c</sup>
Wilms' tumor <sup>b</sup>	72.6	78.4	86.8	91.0	92.2	91.3	91.8	91.6	93.8	91.0	92.6 <sup>c</sup>

Source: SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Based on follow-up of patients into 2013.

Note: Neuroblastoma and Wilms' tumor are not mutually exclusive from the other tumors presented in table.

<sup>a</sup> Neuroblastoma is defined as histologies 9490-9509.

<sup>b</sup> Wilms' tumor is defined as histologies 8959-8960.

<sup>c</sup> The difference between 1975-1977 and 2006-2012 is statistically significant ( $p < .05$ ).

<sup>d</sup> The standard error is between 5 and 10 percentage points.

<sup>e</sup> The standard error is greater than 10 percentage points.

- Statistic could not be calculated due to fewer than 25 cases during the time period.