

Table 32.9

Age-Adjusted SEER Incidence^a, by Year and Sex
By Adapted Classification Scheme for Tumors of Adolescents and Young Adults^b
All Races, Ages 25-29

| Both Sexes | 1975-2013 | | | 2004-2013 | | |
|---|-----------|------|-------------------|-----------|------|-------------------|
| | No. | Rate | AAPC ^c | No. | Rate | AAPC ^c |
| All Sites Combined | 43,116 | 55.8 | 0.7* | 12,401 | 62.0 | 1.1* |
| 1. Leukemias | 1,839 | 2.4 | 0.7* | 535 | 2.7 | 0.7* |
| 2. Lymphomas | 6,413 | 8.3 | 0.6 | 1,694 | 8.5 | 0.4 |
| 3. CNS, intracranial, intraspinal | 2,241 | 2.9 | 1.0* | 648 | 3.2 | 1.0* |
| 4. Osseous & chondromatous neoplasms | 530 | 0.7 | 1.2* | 161 | 0.8 | 1.2* |
| 5. Soft tissue sarcomas | 3,146 | 4.1 | 1.3 | 581 | 2.9 | -0.4 |
| 6. Germ cell and trophoblastic neoplasms | 5,510 | 7.1 | 1.5* | 1,673 | 8.4 | 2.0* |
| 7. Melanoma and skin carcinomas | 5,906 | 7.6 | 0.7* | 1,773 | 8.9 | -1.2 |
| 8. Carcinomas (excluding skin) | 16,903 | 21.9 | 0.5* | 5,150 | 25.7 | 1.5* |
| 9. Miscellaneous specified neoplasms, NOS | 419 | 0.5 | 1.9* | 140 | 0.7 | 1.9* |
| 10. Unspecified malignant neoplasms | 209 | 0.3 | - | 46 | 0.2 | - |

| Males | 1975-2013 | | | 2004-2013 | | |
|---|-----------|------|-------------------|-----------|------|-------------------|
| | No. | Rate | AAPC ^c | No. | Rate | AAPC ^c |
| All Sites Combined | 19,439 | 50.1 | 0.8* | 5,416 | 53.7 | 1.1* |
| 1. Leukemias | 1,061 | 2.7 | 0.3 | 294 | 2.9 | 0.3 |
| 2. Lymphomas | 3,659 | 9.4 | 0.7 | 934 | 9.3 | 0.9 |
| 3. CNS, intracranial, intraspinal | 1,226 | 3.2 | 1.0* | 370 | 3.7 | 1.0* |
| 4. Osseous & chondromatous neoplasms | 306 | 0.8 | 0.5 | 88 | 0.9 | 0.5 |
| 5. Soft tissue sarcomas | 2,302 | 5.9 | 1.7 | 380 | 3.8 | -0.4 |
| 6. Germ cell and trophoblastic neoplasms | 5,091 | 13.1 | 1.6* | 1,578 | 15.6 | 2.0* |
| 7. Melanoma and skin carcinomas | 2,093 | 5.4 | 0.2 | 557 | 5.5 | 0.2 |
| 8. Carcinomas (excluding skin) | 3,455 | 8.9 | 0.4 | 1,132 | 11.2 | 1.9* |
| 9. Miscellaneous specified neoplasms, NOS | 156 | 0.4 | - | 58 | 0.6 | - |
| 10. Unspecified malignant neoplasms | 90 | 0.2 | - | 25 | 0.2 | - |

| Females | 1975-2013 | | | 2004-2013 | | |
|---|-----------|------|-------------------|-----------|------|-------------------|
| | No. | Rate | AAPC ^c | No. | Rate | AAPC ^c |
| All Sites Combined | 23,677 | 61.6 | 0.7* | 6,985 | 70.3 | 1.0* |
| 1. Leukemias | 778 | 2.0 | 1.1* | 241 | 2.4 | 1.1* |
| 2. Lymphomas | 2,754 | 7.2 | 0.7 | 760 | 7.7 | -0.5 |
| 3. CNS, intracranial, intraspinal | 1,015 | 2.6 | 0.9* | 278 | 2.8 | 0.9* |
| 4. Osseous & chondromatous neoplasms | 224 | 0.6 | 1.5* | 73 | 0.7 | 1.5* |
| 5. Soft tissue sarcomas | 844 | 2.2 | 0.5 | 201 | 2.0 | -1.5* |
| 6. Germ cell and trophoblastic neoplasms | 419 | 1.1 | -0.8 | 95 | 1.0 | -0.8 |
| 7. Melanoma and skin carcinomas | 3,813 | 9.9 | 1.1* | 1,216 | 12.2 | -1.1 |
| 8. Carcinomas (excluding skin) | 13,448 | 35.0 | 0.5* | 4,018 | 40.5 | 1.3* |
| 9. Miscellaneous specified neoplasms, NOS | 263 | 0.7 | 1.5* | 82 | 0.8 | 1.5* |
| 10. Unspecified malignant neoplasms | 119 | 0.3 | - | 21 | 0.2 | - |

Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^a Rates are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta).

^b Adapted from: RD Barr, EJ Holowaty, JM Birch. Classification Scheme for tumors diagnosed in adolescents and young adults. Cancer. 106:7; 1425-1430. (Restricted to malignant tumors and updated based on ICD-O-3 definitions).

^c The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint over the time range of 1975-2013.

- Statistic could not be calculated. Count and rate based on less than 16 cases for the time interval. AAPC could not be calculated due to insufficient data.

* The AAPC is significantly different from zero (p<.05).