

Table 33.1
Five-Year Cause-Specific Survival^a(%) for Malignant Cancers, 2007-2013
Both Sexes by Race/Ethnicity and Primary Site

Primary Site	White			Black			Asian/ Pacific Islander			American Indian/ Alaska Native ^b			Hispanic ^c		
	Cases	Percent	SE	Cases	Percent	SE	Cases	Percent	SE	Cases	Percent	SE	Cases	Percent	SE
All Sites	1,815,757	68.1	0.0	248,020	63.2	0.1	151,645	66.8	0.1	9,830	60.5	0.6	238,595	68.2	0.1
Oral Cavity & Pharynx	44,889	69.7	0.3	5,015	52.8	0.8	3,814	69.2	0.9	264	64.3	3.4	4,443	65.7	0.8
Esophagus	17,582	22.0	0.4	2,224	15.0	0.9	916	19.7	1.6	93	22.8	5.0	1,637	19.9	1.2
Stomach	25,101	31.1	0.3	4,965	33.4	0.8	5,197	39.7	0.8	291	23.9	2.8	6,671	31.0	0.7
Colon & Rectum	160,525	65.8	0.1	25,495	60.0	0.4	17,470	69.0	0.4	1,184	60.9	1.7	22,303	66.2	0.4
Liver & Intrahepatic Bile Duct	30,793	20.4	0.3	5,704	17.0	0.7	6,879	28.0	0.7	429	16.5	2.3	8,448	20.9	0.6
Pancreas	47,279	8.5	0.2	7,315	8.8	0.4	4,432	10.3	0.6	280	6.6	2.0	6,254	9.1	0.5
Larynx	13,036	67.4	0.5	2,471	58.0	1.2	540	69.5	2.3	63	58.8	7.1	1,334	67.5	1.5
Lung & Bronchus	214,490	20.7	0.1	30,221	17.9	0.3	16,357	22.6	0.4	987	17.5	1.5	15,449	19.5	0.4
Melanoma of the Skin	92,110	89.6	0.1	462	66.2	2.6	627	76.9	2.0	138	87.8	3.0	3,167	84.2	0.8
Breast	269,966	89.0	0.1	37,817	80.6	0.2	27,878	91.6	0.2	1,436	86.7	1.1	36,961	87.6	0.2
Cervix uteri	16,205	70.9	0.4	3,067	59.5	1.0	1,975	71.7	1.2	166	64.3	4.2	4,918	73.3	0.7
Corpus & Uterus, NOS	59,247	83.2	0.2	7,574	64.1	0.7	5,978	83.7	0.6	408	83.3	2.1	8,637	81.8	0.5
Ovary ^d	27,808	46.8	0.4	2,947	39.1	1.1	2,767	56.9	1.2	180	41.1	4.6	4,394	54.0	0.9
Prostate	274,076	94.2	0.1	54,465	93.0	0.1	17,099	94.4	0.2	940	87.9	1.3	33,015	93.2	0.2
Testis	14,285	95.8	0.2	499	91.5	1.4	618	94.3	1.0	120	93.6	2.3	3,684	93.8	0.5
Urinary Bladder	78,418	78.9	0.2	4,951	67.3	0.8	3,610	78.9	0.8	191	64.9	4.0	5,485	77.0	0.7
Kidney & Renal Pelvis	61,082	75.5	0.2	8,587	77.8	0.5	3,959	74.5	0.8	607	73.4	2.0	10,500	77.3	0.5
Brain & Nervous System	29,117	34.1	0.3	2,464	42.6	1.2	1,947	41.5	1.3	140	43.0	4.6	4,607	46.3	0.9
Thyroid	57,192	97.4	0.1	4,753	96.6	0.3	7,398	96.8	0.2	324	96.2	1.1	11,070	97.5	0.2
Hodgkin Lymphoma	11,893	88.4	0.3	1,877	86.5	0.9	773	88.5	1.3	45	95.1	3.3	2,519	86.7	0.8
Non-Hodgkin Lymphoma	78,456	72.5	0.2	7,827	68.2	0.6	6,509	70.2	0.6	365	74.5	2.4	11,975	71.0	0.5
Myeloma	22,780	53.7	0.4	6,393	56.6	0.8	1,822	55.7	1.5	130	49.1	5.8	3,682	55.3	1.1
Leukemia	52,851	64.2	0.2	5,138	60.0	0.8	3,648	54.8	1.0	281	63.2	3.1	8,968	63.8	0.6
Mesothelioma	3,954	9.4	0.6	227	17.5	2.9	136	7.8	3.2	-	-	-	450	11.3	1.9

Source: SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

^a Five-Year Cause-Specific survival was calculated by actuarial method using diagnosis years 2007-2013 with follow-up through 2014.

^b Data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^c Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

^d Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

- Statistic could not be calculated due to less than 25 cases in the time interval.

Table 33.2
 Five-Year Cause-Specific Survival^a(%) for Malignant Cancers, 2007-2013
 Males by Race/Ethnicity and Primary Site

Primary Site	White			Black			Asian/ Pacific Islander			American Indian/ Alaska Native ^b			Hispanic ^c		
	Cases	Percent	SE	Cases	Percent	SE	Cases	Percent	SE	Cases	Percent	SE	Cases	Percent	SE
All Sites	935,336	67.6	0.1	131,034	65.1	0.1	69,427	62.1	0.2	4,653	56.5	0.8	115,567	66.1	0.2
Oral Cavity & Pharynx	32,420	69.3	0.3	3,494	49.5	1.0	2,512	66.6	1.1	187	64.1	4.1	3,043	62.1	1.0
Esophagus	14,204	21.9	0.4	1,535	13.7	1.1	695	17.3	1.8	69	22.1	6.0	1,326	19.7	1.4
Stomach	15,687	28.7	0.4	2,786	28.7	1.0	2,891	39.4	1.1	170	21.5	3.5	3,774	29.3	0.9
Colon & Rectum	83,500	66.1	0.2	12,392	57.8	0.5	9,152	69.6	0.6	616	61.5	2.3	11,988	65.5	0.5
Liver & Intrahepatic Bile Duct	22,629	20.2	0.3	4,240	16.4	0.8	4,819	27.8	0.8	302	16.9	2.7	6,140	20.3	0.7
Pancreas	24,165	8.7	0.2	3,395	7.8	0.6	2,128	10.8	0.9	147	7.1	2.9	3,043	8.5	0.6
Larynx	10,565	67.5	0.5	1,944	58.4	1.4	464	68.3	2.5	55	62.8	7.2	1,157	66.8	1.7
Lung & Bronchus	111,396	17.9	0.1	16,734	14.7	0.3	9,247	19.4	0.5	522	14.0	2.0	8,222	15.4	0.5
Melanoma of the Skin	52,262	87.3	0.2	198	59.6	4.2	315	71.2	3.1	63	87.1	4.6	1,370	79.0	1.3
Prostate	274,076	94.2	0.1	54,465	93.0	0.1	17,099	94.4	0.2	940	87.9	1.3	33,015	93.2	0.2
Testis	14,285	95.8	0.2	499	91.5	1.4	618	94.3	1.0	120	93.6	2.3	3,684	93.8	0.5
Urinary Bladder	59,670	80.0	0.2	3,309	71.6	0.9	2,734	80.2	0.9	148	67.3	4.5	4,060	79.0	0.8
Kidney & Renal Pelvis	38,264	74.9	0.3	5,094	76.1	0.7	2,538	74.7	1.0	363	70.9	2.8	6,205	75.9	0.6
Brain & Nervous System	16,304	32.8	0.4	1,288	39.9	1.6	1,057	41.8	1.8	85	43.5	5.9	2,524	45.8	1.2
Thyroid	13,524	95.0	0.2	828	93.6	0.9	1,535	94.7	0.6	61	96.7	2.3	1,970	94.7	0.6
Hodgkin Lymphoma	6,515	87.9	0.4	1,017	83.3	1.3	421	87.7	1.9	-	-	-	1,388	84.6	1.1
Non-Hodgkin Lymphoma	42,831	71.7	0.2	4,157	65.1	0.8	3,528	69.9	0.9	195	73.8	3.3	6,440	68.9	0.7
Myeloma	12,985	54.2	0.6	3,086	57.4	1.2	978	53.2	2.1	62	56.8	7.3	2,013	55.7	1.5
Leukemia	30,489	65.0	0.3	2,790	61.0	1.1	2,047	56.6	1.3	153	63.8	4.2	5,038	64.2	0.8
Mesothelioma	3,013	7.4	0.6	155	11.5	3.0	91	5.5	3.2	-	-	-	321	9.0	2.1

Source: SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

^a Five-Year Cause-Specific survival was calculated by actuarial method using diagnosis years 2007-2013 with follow-up through 2014.

^b Data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^c Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

- Statistic could not be calculated due to less than 25 cases in the time interval.

Table 33.3
Five-Year Cause-Specific Survival^a(%) for Malignant Cancers, 2007-2013
Females by Race/Ethnicity and Primary Site

Primary Site	White			Black			Asian/ Pacific Islander			American Indian/ Alaska Native ^b			Hispanic ^c		
	Cases	Percent	SE	Cases	Percent	SE	Cases	Percent	SE	Cases	Percent	SE	Cases	Percent	SE
All Sites	880,421	68.6	0.1	116,986	61.0	0.2	82,218	70.7	0.2	5,177	64.1	0.8	123,028	70.1	0.2
Oral Cavity & Pharynx	12,469	70.5	0.5	1,521	60.4	1.4	1,302	74.1	1.4	77	64.5	6.2	1,400	73.5	1.4
Esophagus	3,378	22.7	0.9	689	17.9	1.8	221	27.9	3.6	-	-	-	311	20.7	3.0
Stomach	9,414	35.0	0.6	2,179	39.3	1.2	2,306	40.1	1.2	121	27.6	4.6	2,897	33.4	1.1
Colon & Rectum	77,025	65.6	0.2	13,103	62.0	0.5	8,318	68.4	0.6	568	60.3	2.4	10,315	67.1	0.6
Liver & Intrahepatic Bile Duct	8,164	20.7	0.6	1,464	18.6	1.5	2,060	28.5	1.2	127	15.3	4.5	2,308	22.4	1.2
Pancreas	23,114	8.3	0.2	3,920	9.6	0.6	2,304	9.9	0.8	133	5.9	2.6	3,211	9.6	0.7
Larynx	2,471	67.1	1.1	527	56.9	2.6	76	76.8	5.5	-	-	-	177	71.7	4.1
Lung & Bronchus	103,094	23.8	0.2	13,487	21.7	0.4	7,110	26.8	0.7	465	21.1	2.2	7,227	23.9	0.6
Melanoma of the Skin	39,848	92.6	0.2	264	70.9	3.2	312	82.4	2.6	75	88.3	3.9	1,797	88.0	0.9
Breast	268,001	89.0	0.1	37,457	80.7	0.2	27,765	91.6	0.2	1,429	86.6	1.1	36,804	87.6	0.2
Cervix uteri	16,205	70.9	0.4	3,067	59.5	1.0	1,975	71.7	1.2	166	64.3	4.2	4,918	73.3	0.7
Corpus & Uterus, NOS	59,247	83.2	0.2	7,574	64.1	0.7	5,978	83.7	0.6	408	83.3	2.1	8,637	81.8	0.5
Ovary ^d	27,808	46.8	0.4	2,947	39.1	1.1	2,767	56.9	1.2	180	41.1	4.6	4,394	54.0	0.9
Urinary Bladder	18,748	75.4	0.4	1,642	58.6	1.4	876	74.8	1.7	43	59.2	7.7	1,425	71.2	1.4
Kidney & Renal Pelvis	22,818	76.6	0.3	3,493	80.1	0.8	1,421	74.3	1.3	244	77.2	2.9	4,295	79.4	0.7
Brain & Nervous System	12,813	35.8	0.5	1,176	45.5	1.6	890	41.2	2.0	55	42.6	7.2	2,083	46.9	1.2
Thyroid	43,668	98.1	0.1	3,925	97.3	0.3	5,863	97.3	0.2	263	96.1	1.3	9,100	98.1	0.2
Hodgkin Lymphoma	5,378	89.1	0.5	860	90.1	1.1	352	89.4	1.8	-	-	-	1,131	89.2	1.0
Non-Hodgkin Lymphoma	35,625	73.5	0.3	3,670	71.7	0.8	2,981	70.6	1.0	170	75.1	3.6	5,535	73.4	0.7
Myeloma	9,795	52.9	0.6	3,307	56.0	1.1	844	58.6	2.2	68	39.2	9.0	1,669	54.8	1.6
Leukemia	22,362	63.1	0.4	2,348	58.8	1.2	1,601	52.6	1.4	128	62.3	4.9	3,930	63.3	0.9
Mesothelioma	941	15.5	1.4	72	31.6	5.9	45	12.5	6.6	-	-	-	129	16.6	4.3

Source: SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

^a Five-Year Cause-Specific survival was calculated by actuarial method using diagnosis years 2007-2013 with follow-up through 2014.

^b Data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^c Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

^d Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

- Statistic could not be calculated due to less than 25 cases in the time interval.

Table 33.4

Cancer of the Female Breast

5-Year Cause-Specific Survival^a and Case Distribution by Race/Ethnicity
SEER 18 Registries, 2007-2013

Race/Ethnicity	All Ages				Ages <50				Ages 50+			
	Case Distribution		Survival		Case Distribution		Survival		Case Distribution		Survival	
	Count	Percent ^b	Percent	SE	Count	Percent ^b	Percent	SE	Count	Percent ^b	Percent	SE
All Races	337,571	100.0	88.3	0.1	79,177	100.0	89.9	0.1	258,394	100.0	87.9	0.1
White	268,001	79.4	89.0	0.1	58,544	73.9	90.9	0.1	209,457	81.1	88.5	0.1
White Non-Hispanic	232,689	68.9	89.2	0.1	46,162	58.3	91.7	0.2	186,527	72.2	88.6	0.1
White Hispanic	35,312	10.5	87.5	0.2	12,382	15.6	87.9	0.4	22,930	8.9	87.3	0.3
Black	37,457	11.1	80.7	0.2	10,611	13.4	81.5	0.5	26,846	10.4	80.3	0.3
American Indian/Alaska Native ^c	1,429	0.4	86.6	1.1	394	0.5	85.8	2.0	1,035	0.4	86.9	1.3
Asian	25,588	7.6	92.0	0.2	8,210	10.4	93.4	0.3	17,378	6.7	91.2	0.3
Asian Indian, Pakistani	2,556	0.8	90.4	0.8	1,030	1.3	91.1	1.2	1,526	0.6	90.0	1.0
Chinese	4,987	1.5	92.7	0.5	1,665	2.1	93.5	0.7	3,322	1.3	92.3	0.6
Filipino	6,976	2.1	91.3	0.4	1,860	2.3	93.5	0.7	5,116	2.0	90.5	0.5
Japanese	3,382	1.0	92.8	0.5	768	1.0	96.1	0.9	2,614	1.0	91.8	0.6
Korean	1,794	0.5	89.9	0.9	666	0.8	92.2	1.3	1,128	0.4	88.6	1.2
Vietnamese	1,648	0.5	91.1	0.9	672	0.8	92.5	1.3	976	0.4	90.1	1.2
Other Asian ^d	4,245	1.3	93.6	0.5	1,549	2.0	94.3	0.8	2,696	1.0	93.2	0.6
Pacific Islander	2,177	0.6	88.0	0.9	613	0.8	85.4	1.8	1,564	0.6	89.1	1.0
Hawaiian	1,375	0.4	89.9	1.0	364	0.5	86.0	2.2	1,011	0.4	91.5	1.1
Other Pacific Islander ^e	802	0.2	84.6	1.6	249	0.3	84.6	2.9	553	0.2	84.6	1.9
Hispanic ^f	36,804	10.9	87.6	0.2	12,918	16.3	87.8	0.4	23,886	9.2	87.4	0.3

Source: SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

^a Five-Year Cause-Specific survival was calculated by actuarial method using diagnosis years 2007-2013 with follow-up through 2014.

^b Percent of cases do not sum to 100 because All Races includes unknown race and other race which are not shown separately.

^c Survival estimates for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^d Other Asians include Laotian, Hmong, Kampuchean, Thai, and Asians, Not Otherwise Specified (NOS).

^e Other Pacific Islanders include Micronesian, Chamorro, Guamanian, Polynesian, Tahitian, Samoan, Tongan, Melanesian, Fiji Islander, New Guinean, and Pacific Islander, NOS.

^f Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

- Statistic not shown due to fewer than 25 cases during the time period.

Table 33.5

Cancer of the Colon and Rectum

5-Year Cause-Specific Survival^a and Case Distribution by Race/Ethnicity
SEER 18 Registries, 2007-2013

Race/Ethnicity	Both Sexes				Males				Females			
	Case Distribution		Survival		Case Distribution		Survival		Case Distribution		Survival	
	Count	Percent ^b	Percent	SE	Count	Percent ^b	Percent	SE	Count	Percent ^b	Percent	SE
All Races	206,912	100.0	65.6	0.1	106,911	100.0	65.6	0.2	100,001	100.0	65.5	0.2
White	160,525	77.6	65.8	0.1	83,500	78.1	66.1	0.2	77,025	77.0	65.6	0.2
White Non-Hispanic	139,129	67.2	65.8	0.1	71,954	67.3	66.2	0.2	67,175	67.2	65.4	0.2
White Hispanic	21,396	10.3	66.1	0.4	11,546	10.8	65.3	0.5	9,850	9.8	67.0	0.6
Black	25,495	12.3	60.0	0.4	12,392	11.6	57.8	0.5	13,103	13.1	62.0	0.5
American Indian/Alaska Native ^c	1,184	0.6	60.9	1.7	616	0.6	61.5	2.3	568	0.6	60.3	2.4
Asian	16,346	7.9	69.5	0.4	8,514	8.0	70.2	0.6	7,832	7.8	68.8	0.6
Asian Indian, Pakistani	990	0.5	71.8	1.8	585	0.5	72.6	2.2	405	0.4	70.4	2.8
Chinese	3,651	1.8	69.3	0.9	1,908	1.8	70.0	1.3	1,743	1.7	68.5	1.3
Filipino	3,554	1.7	70.4	0.9	1,818	1.7	70.2	1.3	1,736	1.7	70.6	1.3
Japanese	2,638	1.3	67.7	1.1	1,353	1.3	69.1	1.4	1,285	1.3	66.3	1.5
Korean	1,542	0.7	68.2	1.4	750	0.7	69.1	2.0	792	0.8	67.3	2.0
Vietnamese	1,538	0.7	66.0	1.5	845	0.8	65.6	2.0	693	0.7	66.5	2.2
Other Asian ^d	2,433	1.2	73.2	1.1	1,255	1.2	74.9	1.5	1,178	1.2	71.6	1.6
Pacific Islander	1,124	0.5	60.7	1.8	638	0.6	60.9	2.4	486	0.5	60.5	2.8
Hawaiian	725	0.4	61.0	2.2	432	0.4	60.2	2.9	293	0.3	62.2	3.5
Other Pacific Islander ^e	399	0.2	60.4	3.0	206	0.2	62.4	4.0	193	0.2	58.0	4.6
Hispanic ^f	22,303	10.8	66.2	0.4	11,988	11.2	65.5	0.5	10,315	10.3	67.1	0.6

Source: SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

^a Five-Year Cause-Specific survival was calculated by actuarial method using diagnosis years 2007-2013 with follow-up through 2014.

^b Percent of cases do not sum to 100 because All Races includes unknown race and other race which are not shown separately.

^c Survival estimates for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^d Other Asians include Laotian, Hmong, Kampuchean, Thai, and Asians, Not Otherwise Specified (NOS).

^e Other Pacific Islanders include Micronesian, Chamorroan, Guamanian, Polynesian, Tahitian, Samoan, Tongan, Melanesian, Fiji Islander, New Guinean, and Pacific Islander, NOS.

^f Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

- Statistic not shown due to fewer than 25 cases during the time period.

Table 33.6

Cancer of the Liver and Intrahepatic Bile Duct

5-Year Cause-Specific Survival^a and Case Distribution by Race/Ethnicity
SEER 18 Registries, 2007-2013

Race/Ethnicity	Both Sexes				Males				Females			
	Case Distribution		Survival		Case Distribution		Survival		Case Distribution		Survival	
	Count	Percent ^b	Percent	SE	Count	Percent ^b	Percent	SE	Count	Percent ^b	Percent	SE
All Races	44,125	100.0	21.2	0.3	32,234	100.0	21.0	0.3	11,891	100.0	21.9	0.5
White	30,793	69.8	20.4	0.3	22,629	70.2	20.2	0.3	8,164	68.7	20.7	0.6
White Non-Hispanic	22,621	51.3	20.2	0.3	16,693	51.8	20.2	0.4	5,928	49.9	20.1	0.7
White Hispanic	8,172	18.5	20.9	0.6	5,936	18.4	20.4	0.7	2,236	18.8	22.1	1.2
Black	5,704	12.9	17.0	0.7	4,240	13.2	16.4	0.8	1,464	12.3	18.6	1.5
American Indian/Alaska Native ^c	429	1.0	16.5	2.3	302	0.9	16.9	2.7	127	1.1	15.3	4.5
Asian	6,488	14.7	28.5	0.7	4,515	14.0	28.5	0.8	1,973	16.6	28.6	1.2
Asian Indian, Pakistani	282	0.6	28.4	3.5	208	0.6	32.8	3.8	74	0.6	17.4	6.5
Chinese	1,520	3.4	34.8	1.5	1,090	3.4	35.4	1.8	430	3.6	33.3	2.6
Filipino	1,038	2.4	21.4	1.6	700	2.2	19.5	1.9	338	2.8	25.2	2.9
Japanese	555	1.3	23.9	2.2	297	0.9	26.7	3.1	258	2.2	20.5	3.2
Korean	695	1.6	28.7	2.1	458	1.4	29.6	2.6	237	2.0	26.9	3.4
Vietnamese	1,385	3.1	30.5	1.5	1,059	3.3	28.7	1.8	326	2.7	36.4	3.2
Other Asian ^d	1,013	2.3	25.3	1.8	703	2.2	24.4	2.2	310	2.6	27.6	3.3
Pacific Islander	391	0.9	18.2	2.8	304	0.9	16.7	2.9	87	0.7	25.3	7.7
Hawaiian	213	0.5	18.6	4.0	166	0.5	18.4	4.3	47	0.4	21.2	10.4
Other Pacific Islander ^e	178	0.4	17.1	3.9	138	0.4	14.0	3.9	40	0.3	33.5	9.5
Hispanic ^f	8,448	19.1	20.9	0.6	6,140	19.0	20.3	0.7	2,308	19.4	22.4	1.2

Source: SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

^a Five-Year Cause-Specific survival was calculated by actuarial method using diagnosis years 2007-2013 with follow-up through 2014.

^b Percent of cases do not sum to 100 because All Races includes unknown race and other race which are not shown separately.

^c Survival estimates for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^d Other Asians include Laotian, Hmong, Kampuchean, Thai, and Asians, Not Otherwise Specified (NOS).

^e Other Pacific Islanders include Micronesian, Chamorroan, Guamanian, Polynesian, Tahitian, Samoan, Tongan, Melanesian, Fiji Islander, New Guinean, and Pacific Islander, NOS.

^f Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

- Statistic not shown due to fewer than 25 cases during the time period.

Table 33.7

Cancer of the Lung and Bronchus

5-Year Cause-Specific Survival^a and Case Distribution by Race/Ethnicity
SEER 18 Registries, 2007-2013

Race/Ethnicity	Both Sexes				Males				Females			
	Case Distribution		Survival		Case Distribution		Survival		Case Distribution		Survival	
	Count	Percent ^b	Percent	SE	Count	Percent ^b	Percent	SE	Count	Percent ^b	Percent	SE
All Races	263,026	100.0	20.6	0.1	138,433	100.0	17.6	0.1	124,593	100.0	23.7	0.1
White	214,490	81.5	20.7	0.1	111,396	80.5	17.9	0.1	103,094	82.7	23.8	0.2
White Non-Hispanic	199,661	75.9	20.8	0.1	103,473	74.7	18.1	0.1	96,188	77.2	23.7	0.2
White Hispanic	14,829	5.6	19.4	0.4	7,923	5.7	15.3	0.5	6,906	5.5	24.0	0.6
Black	30,221	11.5	17.9	0.3	16,734	12.1	14.7	0.3	13,487	10.8	21.7	0.4
American Indian/Alaska Native ^c	987	0.4	17.5	1.5	522	0.4	14.0	2.0	465	0.4	21.1	2.2
Asian	14,966	5.7	23.0	0.4	8,470	6.1	19.8	0.5	6,496	5.2	27.1	0.7
Asian Indian, Pakistani	732	0.3	28.7	2.2	479	0.3	24.7	2.5	253	0.2	35.9	4.0
Chinese	3,760	1.4	23.2	0.9	2,064	1.5	20.4	1.1	1,696	1.4	26.5	1.4
Filipino	3,730	1.4	24.1	0.9	2,196	1.6	20.5	1.1	1,534	1.2	29.1	1.5
Japanese	2,024	0.8	19.0	1.1	995	0.7	16.1	1.5	1,029	0.8	21.7	1.6
Korean	1,207	0.5	20.5	1.4	668	0.5	16.9	1.8	539	0.4	24.8	2.3
Vietnamese	1,768	0.7	24.0	1.2	1,153	0.8	20.6	1.4	615	0.5	30.2	2.3
Other Asian ^d	1,745	0.7	24.2	1.3	915	0.7	19.8	1.8	830	0.7	29.0	2.0
Pacific Islander	1,391	0.5	18.3	1.3	777	0.6	14.3	1.6	614	0.5	23.3	2.2
Hawaiian	901	0.3	18.0	1.6	486	0.4	14.8	2.0	415	0.3	21.8	2.6
Other Pacific Islander ^e	490	0.2	18.9	2.4	291	0.2	13.1	2.8	199	0.2	26.7	3.9
Hispanic ^f	15,449	5.9	19.5	0.4	8,222	5.9	15.4	0.5	7,227	5.8	23.9	0.6

Source: SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

^a Five-Year Cause-Specific survival was calculated by actuarial method using diagnosis years 2007-2013 with follow-up through 2014.

^b Percent of cases do not sum to 100 because All Races includes unknown race and other race which are not shown separately.

^c Survival estimates for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^d Other Asians include Laotian, Hmong, Kampuchean, Thai, and Asians, Not Otherwise Specified (NOS).

^e Other Pacific Islanders include Micronesian, Chamorro, Guamanian, Polynesian, Tahitian, Samoan, Tongan, Melanesian, Fiji Islander, New Guinean, and Pacific Islander, NOS.

^f Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

- Statistic not shown due to fewer than 25 cases during the time period.

Table 33.8

Cancer of the Prostate

5-Year Cause-Specific Survival^a and Case Distribution by Race/Ethnicity
SEER 18 Registries, 2007-2013

Race/Ethnicity	All Ages				Ages <65				Ages 65+			
	Case Distribution		Survival		Case Distribution		Survival		Case Distribution		Survival	
	Count	Percent ^b	Percent	SE	Count	Percent ^b	Percent	SE	Count	Percent ^b	Percent	SE
All Races	358,816	100.0	94.1	0.0	160,692	100.0	96.8	0.1	198,124	100.0	91.9	0.1
White	274,076	76.4	94.2	0.1	119,170	74.2	97.0	0.1	154,906	78.2	91.9	0.1
White Non-Hispanic	243,480	67.9	94.3	0.1	105,704	65.8	97.2	0.1	137,776	69.5	92.0	0.1
White Hispanic	30,596	8.5	93.1	0.2	13,466	8.4	95.4	0.2	17,130	8.6	91.2	0.3
Black	54,465	15.2	93.0	0.1	30,219	18.8	95.6	0.1	24,246	12.2	89.7	0.2
American Indian/Alaska Native ^c	940	0.3	87.9	1.3	379	0.2	92.4	1.6	561	0.3	84.7	1.9
Asian	15,795	4.4	94.7	0.2	5,515	3.4	97.4	0.3	10,280	5.2	93.2	0.3
Asian Indian, Pakistani	1,782	0.5	94.3	0.7	734	0.5	97.1	0.7	1,048	0.5	92.2	1.0
Chinese	3,197	0.9	95.5	0.4	985	0.6	98.9	0.4	2,212	1.1	93.8	0.6
Filipino	4,366	1.2	93.6	0.4	1,594	1.0	96.5	0.5	2,772	1.4	91.9	0.6
Japanese	2,170	0.6	95.4	0.5	620	0.4	97.5	0.7	1,550	0.8	94.5	0.7
Korean	777	0.2	93.0	1.1	193	0.1	94.7	1.9	584	0.3	92.3	1.4
Vietnamese	875	0.2	94.1	0.9	297	0.2	96.4	1.2	578	0.3	93.0	1.2
Other Asian ^d	2,628	0.7	96.0	0.5	1,092	0.7	98.0	0.5	1,536	0.8	94.4	0.7
Pacific Islander	1,304	0.4	90.9	1.0	538	0.3	93.0	1.3	766	0.4	89.4	1.4
Hawaiian	766	0.2	90.0	1.3	319	0.2	92.1	1.7	447	0.2	88.4	1.9
Other Pacific Islander ^e	538	0.1	92.3	1.4	219	0.1	94.4	1.9	319	0.2	90.8	2.0
Hispanic ^f	33,015	9.2	93.2	0.2	14,386	9.0	95.5	0.2	18,629	9.4	91.4	0.2

Source: SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

^a Five-Year Cause-Specific survival was calculated by actuarial method using diagnosis years 2007-2013 with follow-up through 2014.

^b Percent of cases do not sum to 100 because All Races includes unknown race and other race which are not shown separately.

^c Survival estimates for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^d Other Asians include Laotian, Hmong, Kampuchean, Thai, and Asians, Not Otherwise Specified (NOS).

^e Other Pacific Islanders include Micronesian, Chamorroan, Guamanian, Polynesian, Tahitian, Samoan, Tongan, Melanesian, Fiji Islander, New Guinean, and Pacific Islander, NOS.

^f Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

- Statistic not shown due to fewer than 25 cases during the time period.