Table 1.24 Age-Adjusted SEER Incidence Rates and Trends for the Top 15 Cancer Sites by Race/Ethnicity Both Sexes

All Races			White			Black		
	Rateb	APC ^c		Rateb	APC ^c		Rateb	APC ^c
	2010-2014	2005-2014		2010-2014	2005-2014		2010-2014	2005-2014
All Sites	442.7	-1.1*	All Sites	451.8	-1.1*	All Sites	459.3	-1.5*
Breast	67.0	0.0	Breast	67.8	-0.1	Prostate ^f	80.7	-4.5*
Lung and Bronchus	55.8	-2.3*	Lung and Bronchus	57.3	-2.2*	Breast	71.1	0.5
Prostate ^f	54.7	-4.8*	Prostate ^f	52.1	-5.2*	Lung and Bronchus	63.0	-2.5*
Colon and Rectum	40.1	-2.8*	Colon and Rectum	39.3	-2.8*	Colon and Rectum	48.7	-3.3*
Melanoma of the Skin	22.3	1.1*	Melanoma of the Skin	26.6	1.1*	Kidney and Renal Pelvis	18.0	1.0
Urinary Bladder	19.8	-1.2*	Urinary Bladder	21.7	-1.2*	Pancreas	15.5	-0.5
Non-Hodgkin Lymphoma	19.5	-0.5*	Non-Hodgkin Lymphoma	20.4	-0.6*	Non-Hodgkin Lymphoma	14.6	-0.3
Kidney and Renal Pelvis	15.6	0.7	Kidney and Renal Pelvis	16.1	0.7	Corpus and Uterus, NOSf	14.0	2.3*
Thyroid	14.2	4.0*	Thyroid	14.9	3.9*	Myeloma	13.2	0.8
Leukemia	13.7	0.5*	Leukemia	14.5	0.4	Urinary Bladder	12.5	-0.9*
Corpus and Uterus, NOSf	13.6	1.2*	Corpus and Uterus, NOSf	13.8	0.9*	Leukemia	10.9	1.2*
Pancreas	12.5	0.5*	Pancreas	12.5	0.6*	Stomach	10.2	-3.0*
Oral Cavity and Pharynx	11.2	0.7*	Oral Cavity and Pharynx	11.7	1.0*	Liver & IBD ^g	10.1	2.5*
Liver & IBD ^g	8.6	2.6*	Liver & IBD ^g	7.8	3.4*	Oral Cavity and Pharynx	9.1	-1.5*
Stomach	7.3	-0.9*	Brain and ONS ⁹	7.0	-0.5*	Thyroid	8.7	4.4*
Asian/Pacific Islander			American Indian/Alaska Native ^d			Hispanic ^e		
	Rateb	APC ^c		Rateb	APC ^c		Rateb	APC ^c
	2010-2014	2005-2014		2010-2014	2005-2014		2010-2014	2005-2014
All Sites	298.8	-1.2*	All Sites	316.0	-1.4*	All Sites	340.6	-1.3*
Breast	54.1	0.9*	Breast	44.1	0.0	Breast	49.8	0.1
Lung and Bronchus	35.7	-1.8*	Colon and Rectum	41.1	-0.6	Prostate ^f	43.7	-5.6*
Colon and Rectum	33.8	-2.8*	Lung and Bronchus	37.4	-3.8*	Colon and Rectum	33.9	-2.5*
Prostate ^f	27.5	-6.4*	Prostate ^f	26.5	-6.3*	Lung and Bronchus	28.8	-2.5*
Thyroid	13.7	3.3*	Kidney and Renal Pelvis	17.2	-3.2*	Non-Hodgkin Lymphoma	17.8	-0.2
Non-Hodgkin Lymphoma	13.5	0.2	Liver & IBDg	13.3	2.2	Kidney and Renal Pelvis	15.4	0.6
Liver & IBD ^g	13.2	-2.2*	Non-Hodgkin Lymphoma	12.1	-2.0	Liver & IBDg	13.2	1.9*
Corpus and Uterus, NOSf	11.2	2.1*	Stomach	10.6	0.5	Thyroid	12.3	4.2*
Stomach	10.6	-3.2*	Corpus and Uterus, NOSf	10.5	0.8	Corpus and Uterus, NOSf	11.6	2.5*
Pancreas	10.0	0.5	Pancreas	9.6	-2.9	Pancreas	11.1	-0.2
Urinary Bladder	8.7	-1.9*	Thyroid	9.3	5.0*	Urinary Bladder	11.0	-1.7*
Kidney and Renal Pelvis	8.2	0.5	Oral Cavity and Pharynx	8.9	1.4	Leukemia	10.6	0.2
Leukemia	7.8	0.5	Urinary Bladder	8.5	-0.4	Stomach	10.5	-1.7*
Oral Cavity and Pharynx	7.7	-0.5	Leukemia	8.1	0.7	Oral Cavity and Pharynx	6.6	0.1

Source: SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

- Top 15 cancer sites selected based on 2010-2014 age-adjusted rates for the race/ethnic group.
- Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups Census P25-1130).
- The APC is the Annual Percent Change over the time interval.
- Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups Census P25-1130).
- d Rates for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.
- Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives.
- Incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry. f
- The rates for sex-specific cancer sites are calculated using the population for both sexes combined.
- IBD = Intrahepatic Bile Duct. ONS = Other Nervous System.
- Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.
- The APC is significantly different from zero (p<.05).
- Statistic not shown. Rate based on less than 16 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.