Table 1.25

Age-Adjusted SEER Incidence Rates and Trends for the Top 15 Cancer Sites by Race/Ethnicity

Males

Source: SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

- a Top 15 cancer sites selected based on 2010-2014 age-adjusted rates for the race/ethnic group.
- Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups Census P25-1130).
- The APC is the Annual Percent Change over the time interval.
- Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups Census P25-1130).
- d Rates for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.
- Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.
- IBD = Intrahepatic Bile Duct. ONS = Other Nervous System.
- * The APC is significantly different from zero (p<.05).
- Statistic not shown. Rate based on less than 16 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.