

Table 2.17
All Cancer Sites (Invasive)

Age-Adjusted SEER Death Rates^a
By Registry, Race and Sex

| | All Races | | | Whites | | | Blacks | | |
|--|-----------|--------|---------|--------|--------|---------|--------|--------|---------|
| | Total | Males | Females | Total | Males | Females | Total | Males | Females |
| <u>SEER Death Rates^a, 2010-2014</u> | | | | | | | | | |
| California | 149.72 | 176.86 | 130.43 | 155.63 | 183.09 | 135.93 | 188.27 | 225.39 | 165.08 |
| Greater Bay Area | 139.85 | 164.24 | 122.81 | 148.16 | 172.12 | 131.46 | 192.42 | 231.26 | 168.01 |
| San Francisco-Oakland | 142.56 | 167.11 | 125.57 | 149.37 | 172.96 | 133.01 | 198.15 | 240.88 | 171.48 |
| San Jose-Monterey | 134.53 | 158.65 | 117.35 | 145.91 | 170.51 | 128.62 | 154.36 | 173.16 | 142.43 |
| Los Angeles | 145.14 | 171.43 | 127.11 | 148.56 | 174.41 | 130.51 | 193.38 | 232.88 | 170.55 |
| Greater California | 155.24 | 183.66 | 134.61 | 160.36 | 189.41 | 139.18 | 181.07 | 214.44 | 157.92 |
| Connecticut | 153.38 | 182.70 | 133.58 | 154.40 | 183.29 | 134.75 | 158.89 | 200.32 | 134.25 |
| Detroit | 179.47 | 214.96 | 155.75 | 173.85 | 207.31 | 151.23 | 209.89 | 260.48 | 178.46 |
| Georgia | 169.67 | 212.42 | 140.57 | 166.92 | 206.25 | 138.73 | 186.69 | 250.62 | 151.14 |
| Atlanta | 155.50 | 191.20 | 133.30 | 146.99 | 177.54 | 127.04 | 181.39 | 240.51 | 151.08 |
| Rural Georgia | 183.22 | 235.68 | 146.98 | 168.87 | 205.48 | 142.10 | 213.23 | 319.30 | 153.94 |
| Greater Georgia | 176.16 | 221.52 | 144.29 | 174.14 | 216.54 | 143.08 | 190.47 | 255.81 | 152.28 |
| Hawaii | 137.61 | 167.34 | 114.99 | 159.58 | 184.02 | 137.01 | 158.00 | 194.84 | 117.29 |
| Iowa | 168.93 | 206.72 | 142.23 | 168.64 | 206.24 | 142.08 | 224.83 | 268.69 | 195.60 |
| Kentucky | 201.52 | 249.61 | 167.15 | 201.89 | 250.20 | 167.10 | 214.69 | 268.09 | 181.40 |
| Louisiana | 190.96 | 237.81 | 157.16 | 180.78 | 222.54 | 149.99 | 223.51 | 291.92 | 178.67 |
| New Jersey | 161.27 | 191.42 | 141.67 | 164.65 | 195.16 | 144.48 | 182.00 | 226.02 | 157.70 |
| New Mexico | 146.55 | 176.20 | 123.87 | 149.13 | 178.68 | 126.30 | 163.07 | 203.11 | 131.35 |
| Seattle-Puget Sound | 158.75 | 187.82 | 137.78 | 161.86 | 190.21 | 141.31 | 180.96 | 224.69 | 146.40 |
| Utah | 128.56 | 151.04 | 111.23 | 128.93 | 151.67 | 111.36 | 144.27 | 151.88 | 135.09 |
| SEER 9 Areas ^b | 154.97 | 185.13 | 133.82 | 156.54 | 185.98 | 135.64 | 189.78 | 238.21 | 160.41 |
| SEER 13 Areas ^b | 151.98 | 181.06 | 131.62 | 154.54 | 183.01 | 134.34 | 190.27 | 236.75 | 162.26 |
| SEER 18 Areas ^b | 160.32 | 192.38 | 137.68 | 162.83 | 194.47 | 140.20 | 193.70 | 244.35 | 162.62 |
| Total U.S. | 166.12 | 200.49 | 141.53 | 166.24 | 199.79 | 141.86 | 194.16 | 247.33 | 161.75 |

^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b The SEER 9 areas are San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah and Atlanta. The SEER 13 areas comprise the SEER 9 areas plus San Jose-Monterey, Los Angeles, the Alaska Native Registry and Rural Georgia. The SEER 18 areas comprise the SEER 13 areas plus California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG.

- Statistic not shown. Rate based on less than 16 cases for the time interval.