

Table 12.13
Cancer of the Larynx (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years,
Lifetime Risk of Being Diagnosed with Cancer Given Alive and Cancer-Free at Current Age, and
Lifetime Risk of Dying from Cancer Given Alive at Current Age
Males, 2012-2014 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer
		+10 yrs	+20 yrs	+30 yrs	Ever				+10 yrs	+20 yrs	+30 yrs	Ever	
All Races	0	0.00	0.00	0.00	0.55	0.20	Asian/ Pacific Islander	0	0.00	0.00	0.00	0.25	0.12
	10	0.00	0.00	0.00	0.55	0.20		10	0.00	0.00	0.00	0.25	0.12
	20	0.00	0.00	0.03	0.55	0.20		20	0.00	0.00	0.01	0.25	0.12
	30	0.00	0.03	0.11	0.56	0.20		30	0.00	0.01	0.03	0.25	0.12
	40	0.02	0.11	0.27	0.57	0.21		40	0.01	0.03	0.08	0.25	0.12
	50	0.09	0.26	0.43	0.56	0.21		50	0.03	0.08	0.17	0.25	0.12
	60	0.18	0.37	0.48	0.50	0.20		60	0.05	0.15	0.21	0.23	0.12
	70	0.22	0.35	-	0.38	0.16		70	0.10	0.17	-	0.19	0.11
80	0.19	-	-	0.22	0.12	80	0.09	-	-	0.12	0.10		
White	0	0.00	0.00	0.00	0.54	0.19	American Indian/ Alaska Native ^a	0	0.00	0.00	0.00	0.33	0.15
	10	0.00	0.00	0.00	0.55	0.19		10	0.00	0.00	0.00	0.33	0.16
	20	0.00	0.00	0.03	0.55	0.19		20	0.00	0.00	0.02	0.33	0.16
	30	0.00	0.03	0.11	0.56	0.19		30	0.00	0.02	0.09	0.34	0.16
	40	0.02	0.11	0.27	0.56	0.20		40	0.02	0.09	0.14	0.35	0.17
	50	0.09	0.26	0.43	0.56	0.20		50	0.07	0.13	0.26	0.35	0.17
	60	0.18	0.37	0.48	0.50	0.19		60	0.06	0.20	0.30	0.30	0.17
	70	0.22	0.35	-	0.37	0.15		70	0.17	0.28	-	0.28	0.16
80	0.19	-	-	0.22	0.11	80	0.16	-	-	0.16	0.15		
Black	0	0.00	0.00	0.00	0.75	0.30	Hispanic ^b	0	0.00	0.00	0.00	0.46	0.20
	10	0.00	0.00	0.00	0.76	0.31		10	0.00	0.00	0.00	0.47	0.20
	20	0.00	0.00	0.03	0.77	0.31		20	0.00	0.00	0.01	0.47	0.20
	30	0.00	0.03	0.17	0.78	0.32		30	0.00	0.01	0.06	0.47	0.21
	40	0.03	0.17	0.43	0.80	0.33		40	0.01	0.06	0.18	0.47	0.21
	50	0.15	0.42	0.65	0.80	0.33		50	0.05	0.17	0.32	0.47	0.21
	60	0.30	0.56	0.70	0.73	0.32		60	0.13	0.28	0.40	0.45	0.21
	70	0.33	0.51	-	0.55	0.27		70	0.18	0.31	-	0.36	0.19
80	0.29	-	-	0.35	0.19	80	0.17	-	-	0.24	0.14		

Devcan 6.7.5, April 2017, National Cancer Institute (<https://surveillance.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, and Georgia excluding ATL/RG). Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

- Statistic could not be calculated.

A percent of 0.00 represents a value that is below 0.005.