

Table 12.17  
Cancer of the Larynx (Invasive)

Age-Adjusted SEER Death Rates<sup>a</sup>  
By Registry, Race and Sex

	All Races			Whites			Blacks		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
<u>SEER Death Rates<sup>a</sup>, 2010-2014</u>									
California	0.74	1.38	0.24	0.78	1.43	0.26	1.29	2.43	0.48
Greater Bay Area	0.58	1.18	0.13	0.61	1.22	0.14	1.24	2.43	-
San Francisco-Oakland	0.55	1.09	0.14	0.53	1.05	-	1.25	2.41	-
San Jose-Monterey	0.65	1.35	-	0.75	1.53	-	-	-	-
Los Angeles	0.70	1.35	0.20	0.70	1.33	0.20	1.47	2.96	-
Greater California	0.81	1.46	0.30	0.86	1.54	0.32	1.14	1.88	-
Connecticut	0.80	1.45	0.29	0.75	1.39	0.25	1.45	2.40	-
Detroit	1.28	2.25	0.56	1.05	1.85	0.44	2.27	4.12	1.03
Georgia	1.06	1.92	0.39	0.98	1.70	0.40	1.40	2.83	0.43
Atlanta	0.87	1.60	0.35	0.82	1.42	0.38	1.05	2.10	-
Rural Georgia	-	-	-	-	-	-	-	-	-
Greater Georgia	1.14	2.05	0.42	1.05	1.81	0.42	1.59	3.21	0.47
Hawaii	0.66	1.36	-	1.08	2.02	-	-	-	-
Iowa	1.01	1.77	0.41	1.01	1.77	0.41	-	-	-
Kentucky	1.49	2.52	0.67	1.44	2.44	0.63	2.53	4.23	-
Louisiana	1.32	2.44	0.45	1.13	1.92	0.50	1.92	4.25	-
New Jersey	1.09	1.93	0.45	1.07	1.91	0.42	1.65	2.91	0.84
New Mexico	0.92	1.63	0.31	0.97	1.70	0.35	-	-	-
Seattle-Puget Sound	0.89	1.53	0.38	0.91	1.51	0.42	-	-	-
Utah	0.34	0.54	-	0.33	0.54	-	-	-	-
SEER 9 Areas <sup>b</sup>	0.84	1.51	0.32	0.82	1.45	0.32	1.58	2.95	0.61
SEER 13 Areas <sup>b</sup>	0.80	1.46	0.28	0.79	1.42	0.29	1.56	2.98	0.59
SEER 18 Areas <sup>b</sup>	0.93	1.66	0.34	0.91	1.62	0.35	1.62	3.12	0.59
Total U.S.	1.03	1.85	0.38	0.99	1.76	0.37	1.71	3.38	0.57

<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

<sup>b</sup> The SEER 9 areas are San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah and Atlanta. The SEER 13 areas comprise the SEER 9 areas plus San Jose-Monterey, Los Angeles, the Alaska Native Registry and Rural Georgia. The SEER 18 areas comprise the SEER 13 areas plus California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG.

- Statistic not shown. Rate based on less than 16 cases for the time interval.