


Table 12.18  
Cancer of the Larynx (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2010-2014

Males and Females

State	Rate	SE	Rank(CI) <sup>d</sup>	PD
TOTAL U.S.	1.03	0.01		
High Five States				
D.C.	1.63	0.24	01(01,31)	58.23 <sup>c</sup>
West Virginia	1.55	0.11	02(01,11)	50.37 <sup>c</sup>
Kentucky	1.49	0.08	03(01,09)	44.11 <sup>c</sup>
Tennessee	1.32	0.06	04(02,18)	27.71 <sup>c</sup>
Louisiana	1.32	0.07	05(01,20)	27.64 <sup>c</sup>
Low Five States				
Oregon	0.70	0.06	47(36,50)	-32.27 <sup>c</sup>
Colorado	0.68	0.05	48(37,50)	-34.20 <sup>c</sup>
South Dakota	0.67	0.12	49(24,51)	-34.93 <sup>c</sup>
Hawaii	0.66	0.09	50(32,50)	-35.80 <sup>c</sup>
Utah	0.34	0.05	51(50,51)	-67.33 <sup>c</sup>

State	Rate	SE	Rank(CI) <sup>d</sup>	PD	State	Rate	SE	Rank(CI) <sup>d</sup>	PD
Alabama	1.23	0.07	10(03,27)	19.45 <sup>c</sup>	Montana	1.03	0.13	27(03,48)	-0.37
Alaska	0.82	0.19	40(03,51)	-20.85 <sup>c</sup>	Nebraska	1.02	0.10	28(05,44)	-0.83
Arizona	0.76	0.04	43(34,50)	-26.69 <sup>c</sup>	Nevada	0.92	0.08	35(13,48)	-11.36
Arkansas	1.01	0.08	30(10,41)	-2.17	New Hampshire	0.86	0.11	38(11,50)	-16.86 <sup>c</sup>
California	0.74	0.02	44(38,49)	-28.33 <sup>c</sup>	New Jersey	1.09	0.05	22(09,33)	5.36
Colorado	0.68	0.05	48(37,50)	-34.20 <sup>c</sup>	New Mexico	0.92	0.09	34(12,48)	-11.28
Connecticut	0.80	0.06	41(29,50)	-22.49 <sup>c</sup>	New York	1.10	0.03	20(11,30)	6.91
Delaware	1.16	0.14	14(01,43)	12.36	North Carolina	1.13	0.05	16(07,31)	9.06
D.C.	1.63	0.24	01(01,31)	58.23 <sup>c</sup>	North Dakota	1.05	0.16	26(02,50)	1.42
Florida	1.17	0.03	13(07,23)	13.45	Ohio	1.29	0.04	07(03,17)	24.70 <sup>c</sup>
Georgia	1.06	0.05	24(11,35)	2.87	Oklahoma	0.96	0.07	32(13,42)	-6.97
Hawaii	0.66	0.09	50(32,50)	-35.80 <sup>c</sup>	Oregon	0.70	0.06	47(36,50)	-32.27 <sup>c</sup>
Idaho	0.70	0.09	46(28,50)	-32.16 <sup>c</sup>	Pennsylvania	1.10	0.04	21(10,31)	6.87
Illinois	1.14	0.04	15(08,29)	10.74	Rhode Island	1.13	0.13	17(02,43)	8.98
Indiana	1.13	0.06	18(06,33)	8.96	South Carolina	1.31	0.07	06(01,19)	26.75 <sup>c</sup>
Iowa	1.01	0.07	29(09,41)	-2.12	South Dakota	0.67	0.12	49(24,51)	-34.93 <sup>c</sup>
Kansas	0.78	0.07	42(29,50)	-24.86 <sup>c</sup>	Tennessee	1.32	0.06	04(02,18)	27.71 <sup>c</sup>
Kentucky	1.49	0.08	03(01,09)	44.11 <sup>c</sup>	Texas	1.06	0.03	25(14,33)	2.81
Louisiana	1.32	0.07	05(01,20)	27.64 <sup>c</sup>	Utah	0.34	0.05	51(50,51)	-67.33 <sup>c</sup>
Maine	1.25	0.12	09(01,35)	21.45 <sup>c</sup>	Vermont	1.18	0.18	12(01,47)	13.91
Maryland	1.12	0.06	19(07,34)	8.29	Virginia	1.01	0.05	31(14,38)	-2.44
Massachusetts	0.94	0.05	33(21,41)	-9.03	Washington	0.87	0.05	37(26,46)	-15.45 <sup>c</sup>
Michigan	1.20	0.05	11(04,25)	16.61 <sup>c</sup>	West Virginia	1.55	0.11	02(01,11)	50.37 <sup>c</sup>
Minnesota	0.72	0.05	45(36,50)	-30.01 <sup>c</sup>	Wisconsin	0.89	0.05	36(25,45)	-14.05
Mississippi	1.26	0.09	08(02,29)	21.81 <sup>c</sup>	Wyoming	0.84	0.17	39(04,50)	-18.70 <sup>c</sup>
Missouri	1.07	0.05	23(09,36)	3.42					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p < .0002$ ).

<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.

<sup>d</sup> Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.