

Table 13.5
LeukemiaAge-adjusted SEER Incidence^a Rates by Year, Race and Sex

	All Races			Whites			Blacks		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
Year of Diagnosis:									
1975-2014	13.46	17.66	10.35	14.13	18.60	10.84	11.11	14.39	8.85
1975	12.80	16.93	9.96	13.10	17.49	10.10	11.58	16.33	8.47
1976	13.54	18.49	10.19	13.83	19.12	10.30	11.37	14.32	9.39
1977	12.74	16.34	10.22	13.20	17.04	10.52	10.15	13.61	7.72
1978	12.87	17.25	9.79	13.54	18.30	10.26	9.16	11.42	7.47
1979	12.56	16.79	9.70	13.03	17.56	10.02	10.60	13.52	8.64
1980	12.93	17.36	9.85	13.28	17.91	10.12	12.18	16.11	9.47
1981	12.38	16.47	9.65	12.65	17.03	9.82	12.27	14.47	10.37
1982	13.15	18.23	9.71	13.50	18.77	9.95	10.75	15.16	7.91
1983	13.15	17.81	9.91	13.48	18.43	10.07	12.48	15.95	10.06
1984	13.01	17.11	10.12	13.47	17.98	10.35	10.68	11.87	9.61
1985	13.38	17.78	10.38	13.75	18.30	10.64	12.32	17.03	9.30
1986	12.84	17.54	9.51	13.39	18.38	9.92	10.23	13.20	8.09
1987	13.36	17.76	10.33	13.81	18.47	10.62	12.57	17.60	9.26
1988	12.98	17.52	9.71	13.42	18.29	10.00	11.55	14.03	9.42
1989	13.46	17.69	10.48	14.00	18.33	11.01	11.26	15.80	8.11
1990	12.93	17.35	9.81	13.55	18.28	10.28	11.40	14.91	8.89
1991	13.08	16.82	10.25	13.71	17.78	10.60	10.81	12.09	9.98
1992	13.06	17.55	9.81	13.65	18.71	10.02	10.68	13.91	8.42
1993	13.09	17.10	10.08	13.66	18.02	10.37	11.68	15.29	9.14
1994	13.10	17.01	10.22	13.79	17.94	10.75	9.86	12.71	7.78
1995	13.64	17.80	10.54	14.52	18.91	11.27	9.97	13.69	7.42
1996	13.20	17.05	10.44	13.79	17.85	10.89	10.77	14.00	8.73
1997	13.25	17.66	10.04	14.08	18.76	10.68	10.67	14.23	8.29
1998	13.24	17.22	10.31	14.04	18.13	11.01	10.02	14.44	7.27
1999	12.88	17.20	9.68	13.52	18.02	10.16	10.62	14.18	8.32
2000	13.80	17.92	10.71	14.50	18.87	11.22	11.84	13.86	10.34
2001	14.47	18.83	11.17	15.42	19.93	12.00	11.93	14.60	10.07
2002	13.66	18.06	10.34	14.74	19.43	11.17	9.00	11.89	7.00
2003	13.72	17.98	10.49	14.50	19.13	10.98	11.48	14.95	9.00
2004	13.99	18.21	10.79	14.71	19.07	11.31	13.06	17.84	10.04
2005	13.93	18.25	10.60	14.86	19.72	11.06	11.42	12.87	10.36
2006	13.97	17.49	11.35	15.00	18.58	12.32	11.34	15.40	8.53
2007	13.55	17.77	10.35	14.61	19.08	11.15	9.92	13.63	7.64
2008	14.09	18.36	10.74	15.00	19.50	11.40	10.92	14.87	8.05
2009	13.69	17.65	10.61	14.55	18.80	11.16	11.23	15.29	8.55
2010	14.38	18.82	11.01	15.39	20.07	11.73	10.90	14.53	8.73
2011	14.55	18.63	11.27	15.41	19.67	11.92	12.46	14.84	10.56
2012	14.30	18.00	11.39	15.21	18.99	12.17	11.95	15.21	9.75
2013	14.14	18.14	10.96	15.19	19.30	11.87	11.37	15.06	8.65
2014	13.92	18.11	10.58	14.85	19.15	11.34	10.55	13.45	8.54

^a SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).
 Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
 - Statistic not shown. Rate based on less than 16 cases for the time interval.