


Table 15.25
Cancer of the Lung and Bronchus (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2010-2014

Males

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	55.89	0.09		
High Five States				
Kentucky	89.63	0.93	01(01,01)	60.38 ^c
Mississippi	82.73	1.11	02(02,03)	48.04 ^c
Arkansas	81.05	1.04	03(02,05)	45.03 ^c
Tennessee	78.43	0.72	04(03,06)	40.34 ^c
Alabama	77.16	0.82	05(04,06)	38.07 ^c
Low Five States				
Hawaii	40.83	1.06	47(44,49)	-26.94 ^c
California	39.84	0.22	48(46,49)	-28.72 ^c
New Mexico	38.14	0.87	49(47,50)	-31.76 ^c
Colorado	36.42	0.59	50(49,50)	-34.84 ^c
Utah	24.58	0.71	51(51,51)	-56.02 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	77.16	0.82	05(04,06)	38.07 ^c	Montana	46.23	1.29	43(37,46)	-17.27 ^c
Alaska	54.77	2.33	25(17,40)	-2.00	Nebraska	54.30	1.10	27(22,35)	-2.84
Arizona	42.96	0.50	45(43,47)	-23.14 ^c	Nevada	52.78	0.92	31(24,37)	-5.56 ^c
Arkansas	81.05	1.04	03(02,05)	45.03 ^c	New Hampshire	53.83	1.27	29(22,37)	-3.69
California	39.84	0.22	48(46,49)	-28.72 ^c	New Jersey	48.41	0.48	40(37,42)	-13.38 ^c
Colorado	36.42	0.59	50(49,50)	-34.84 ^c	New Mexico	38.14	0.87	49(47,50)	-6.91 ^c
Connecticut	46.31	0.72	42(40,44)	-17.14 ^c	New York	49.03	0.33	39(36,41)	-12.27 ^c
Delaware	62.35	1.63	16(12,22)	11.56 ^c	North Carolina	67.88	0.55	11(10,13)	21.46 ^c
D.C.	49.61	2.05	38(25,44)	-11.23 ^c	North Dakota	52.03	1.71	34(23,41)	-6.91 ^c
Florida	54.09	0.30	28(24,31)	-3.22	Ohio	66.53	0.48	13(11,14)	19.04 ^c
Georgia	64.17	0.59	15(13,16)	14.83 ^c	Oklahoma	71.55	0.88	08(07,10)	28.03 ^c
Hawaii	40.83	1.06	47(44,49)	-26.94 ^c	Oregon	50.57	0.71	36(31,40)	-9.51 ^c
Idaho	43.73	1.08	44(42,47)	-21.76 ^c	Pennsylvania	59.06	0.41	20(17,23)	5.68 ^c
Illinois	59.00	0.45	21(17,23)	5.56 ^c	Rhode Island	59.12	1.50	19(15,26)	5.78 ^c
Indiana	71.19	0.68	09(07,10)	27.39 ^c	South Carolina	67.06	0.77	12(10,14)	19.99 ^c
Iowa	59.15	0.85	18(16,24)	5.83 ^c	South Dakota	55.51	1.60	24(18,35)	-0.68
Kansas	57.35	0.91	23(18,26)	2.62	Tennessee	78.43	0.72	04(03,06)	40.34 ^c
Kentucky	89.63	0.93	01(01,01)	60.38 ^c	Texas	52.24	0.33	32(28,36)	-6.52 ^c
Louisiana	72.90	0.85	07(06,09)	30.45 ^c	Utah	24.58	0.71	51(51,51)	-56.02 ^c
Maine	64.45	1.29	14(11,17)	15.32 ^c	Vermont	54.33	1.79	26(19,38)	-2.80
Maryland	52.20	0.64	33(27,37)	-6.61 ^c	Virginia	57.41	0.57	22(19,26)	2.72
Massachusetts	51.86	0.57	35(28,37)	-7.21 ^c	Washington	49.88	0.57	37(34,40)	-10.74 ^c
Michigan	60.77	0.49	17(16,19)	8.74 ^c	West Virginia	77.15	1.21	06(03,07)	38.04 ^c
Minnesota	47.57	0.61	41(38,43)	-14.87 ^c	Wisconsin	53.43	0.61	30(24,35)	-4.40
Mississippi	82.73	1.11	02(02,03)	48.04 ^c	Wyoming	41.93	1.76	46(42,49)	-24.97 ^c
Missouri	69.36	0.68	10(08,12)	24.10 ^c					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 5% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.