

Table 17.6
MesotheliomaSEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Males

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013
Survival Time																							
1-year	32.9	36.0	33.8	32.8	34.9	45.0	32.7	32.8	28.8	41.3	36.5	30.6	44.9	37.5	40.5	39.9	40.7	41.8	46.2	30.9	45.8	39.9	41.0
2-year	11.1	15.9	14.2	11.7	11.1	24.5	14.3	17.6	13.7	13.5	20.3	14.5	18.5	14.7	17.3	22.0	15.2	19.3	25.6	16.0	22.4	23.5	
3-year	6.6	9.5	8.4	7.4	8.9	16.2	8.7	11.7	6.8	9.0	10.8	6.0	13.9	8.9	8.9	12.8	10.5	13.6	13.9	6.2	13.8		
4-year	5.2	6.6	5.8	4.7	5.9	11.3	6.2	10.1	4.4	6.7	8.8	3.1	12.6	6.6	6.3	9.4	5.9	11.2	11.1	3.5			
5-year	4.0	5.1	4.6	3.6	4.2	10.3	3.6	10.1	3.8	4.0	8.2	3.1	7.1	3.9	6.3	7.6	4.3	8.5	8.9				
6-year	3.3	4.3	3.2	3.1	3.5	10.3	2.2	8.2	3.8	1.7	7.5	2.4	6.5	3.9	6.3	6.3	4.3	6.9					
7-year	2.6	3.4	2.5	2.6	2.2	9.5	1.5	7.8	3.8	1.7	5.1	1.6	5.7	2.1	6.3	6.3	3.3						
8-year	2.3	3.3	2.5	2.1	2.2	6.7	1.5	4.8	3.8	0.9	5.1	1.6	5.7	2.1	6.2	5.1							
9-year	2.1	3.1	2.2	1.8	0.0	5.2	0.8	4.8	2.8	0.9	5.1	1.6	5.7	1.1	6.2								
10-year	2.1	3.1	1.9	1.7	0.0	4.5	0.8	4.1	2.8	0.9	5.1	1.6	4.5	1.1									
11-year	2.1	3.1	1.9	1.6	0.0	3.9	0.0	4.1	2.2	0.9	2.8	1.6	4.5										
12-year	1.9	2.9	1.9	1.4	0.0	3.2	0.0	3.6	2.2	0.9	2.8	1.0											
13-year	1.7	2.8	1.8	1.4	0.0	2.3	0.0	3.6	2.2	0.9	2.8												
14-year	1.4	2.8	1.7	1.4	0.0	2.3	0.0	3.6	1.6	0.9													
15-year	1.1	2.5	1.6	1.4	0.0	2.3	0.0	3.6	1.6														
16-year	1.1	2.5	1.6	1.4	0.0	2.3	0.0	3.6															
17-year	1.1	2.4	1.6	1.2	0.0	1.8	0.0																
18-year	1.1	2.4	1.6	1.2	0.0	1.8																	
19-year	1.1	2.4	1.6	0.7	0.0																		
20-year	1.1	2.4	1.6	0.7																			

a

Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).