

Table 17.8
Mesothelioma (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years,
Lifetime Risk of Being Diagnosed with Cancer Given Alive and Cancer-Free at Current Age, and
Lifetime Risk of Dying from Cancer Given Alive at Current Age
Both Sexes, 2012-2014 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer
		+10 yrs	+20 yrs	+30 yrs	Ever				+10 yrs	+20 yrs	+30 yrs	Ever	
All Races	0	0.00	0.00	0.00	0.04	0.09	Asian/ Pacific Islander	0	0.00	0.00	0.00	0.02	0.04
	10	0.00	0.00	0.01	0.04	0.10		10	0.00	0.00	0.00	0.02	0.04
	20	0.00	0.01	0.02	0.04	0.10		20	0.00	0.00	0.01	0.02	0.04
	30	0.01	0.02	0.02	0.04	0.10		30	0.00	0.01	0.01	0.02	0.04
	40	0.01	0.01	0.02	0.03	0.10		40	0.00	0.01	0.01	0.02	0.04
	50	0.01	0.01	0.02	0.03	0.10		50	0.00	0.01	0.01	0.01	0.04
	60	0.00	0.01	0.02	0.02	0.10		60	0.00	0.01	0.01	0.01	0.04
	70	0.01	0.02	-	0.02	0.10		70	0.00	0.01	-	0.01	0.03
80	0.01	-	-	0.01	0.07	80	0.01	-	-	0.01	0.03		
White	0	0.00	0.00	0.00	0.04	0.10	American Indian/ Alaska Native ^a	0	0.00	0.00	0.00	0.03	0.08
	10	0.00	0.00	0.01	0.04	0.11		10	0.00	0.00	0.00	0.03	0.08
	20	0.00	0.01	0.01	0.04	0.11		20	0.00	0.00	0.00	0.03	0.08
	30	0.01	0.01	0.02	0.03	0.11		30	0.00	0.00	0.01	0.03	0.08
	40	0.01	0.01	0.01	0.03	0.11		40	0.00	0.01	0.01	0.03	0.08
	50	0.00	0.01	0.01	0.02	0.11		50	0.00	0.00	0.02	0.03	0.09
	60	0.00	0.01	0.02	0.02	0.11		60	0.00	0.02	0.02	0.02	0.09
	70	0.01	0.01	-	0.02	0.10		70	0.02	0.03	-	0.03	0.09
80	0.01	-	-	0.01	0.08	80	0.00	-	-	0.00	0.09		
Black	0	0.00	0.00	0.02	0.08	0.04	Hispanic ^b	0	0.00	0.00	0.00	0.07	0.06
	10	0.00	0.02	0.04	0.08	0.04		10	0.00	0.00	0.01	0.07	0.07
	20	0.02	0.04	0.06	0.08	0.04		20	0.00	0.01	0.02	0.07	0.07
	30	0.03	0.04	0.05	0.06	0.04		30	0.01	0.02	0.02	0.07	0.07
	40	0.02	0.03	0.03	0.03	0.04		40	0.01	0.01	0.02	0.06	0.07
	50	0.01	0.01	0.02	0.02	0.04		50	0.01	0.01	0.03	0.06	0.07
	60	0.00	0.01	0.01	0.01	0.04		60	0.01	0.02	0.04	0.05	0.06
	70	0.00	0.01	-	0.01	0.04		70	0.02	0.04	-	0.05	0.06
80	0.00	-	-	0.00	0.03	80	0.02	-	-	0.04	0.04		

Devcan 6.7.5, April 2017, National Cancer Institute (<https://surveillance.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, and Georgia excluding ATL/RG). Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

- Statistic could not be calculated.

A percent of 0.00 represents a value that is below 0.005.