

Table 17.12
MesotheliomaAge-Adjusted SEER Incidence Rates^a
By Registry, Race and Sex

	All Races			Whites			Blacks		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
<u>SEER Incidence Rates^a, 2010-2014</u>									
California	0.89	1.55	0.40	1.02	1.77	0.46	0.63	1.23	-
Greater Bay Area	0.93	1.59	0.47	1.15	1.98	0.54	-	-	-
San Francisco-Oakland	0.97	1.61	0.51	1.22	2.02	0.64	-	-	-
San Jose-Monterey	0.87	1.55	0.38	1.04	1.92	0.37	-	-	-
Los Angeles	0.75	1.26	0.38	0.89	1.48	0.45	0.56	1.03	-
Greater California	0.94	1.67	0.39	1.04	1.83	0.44	0.73	1.52	-
Connecticut	1.01	1.83	0.44	1.09	1.95	0.48	-	-	-
Detroit	0.94	1.77	0.36	1.10	2.07	0.39	0.38	-	-
Georgia	0.61	1.14	0.22	0.71	1.31	0.25	0.29	0.52	-
Atlanta	0.52	1.02	-	0.69	1.24	-	-	-	-
Rural Georgia	-	-	-	-	-	-	-	-	-
Greater Georgia	0.64	1.17	0.25	0.71	1.31	0.25	0.34	-	-
Hawaii	0.49	0.94	-	0.78	-	-	-	-	-
Iowa	0.89	1.55	0.41	0.89	1.58	0.39	-	-	-
Kentucky	0.78	1.38	0.35	0.81	1.43	0.36	-	-	-
Louisiana	1.32	2.31	0.63	1.46	2.52	0.68	0.93	1.71	0.48
New Jersey	1.20	2.23	0.48	1.34	2.47	0.54	0.53	1.01	-
New Mexico	0.76	1.35	0.29	0.80	1.38	0.33	-	-	-
Seattle-Puget Sound	1.35	2.38	0.57	1.43	2.50	0.61	-	-	-
Utah	0.79	1.35	0.32	0.79	1.36	0.32	-	-	-
SEER 9 Areas ^b	0.93	1.66	0.40	1.05	1.85	0.46	0.41	0.72	0.22
SEER 13 Areas ^b	0.89	1.56	0.39	1.02	1.78	0.45	0.44	0.78	0.23
SEER 18 Areas ^b	0.93	1.67	0.40	1.05	1.85	0.45	0.53	0.98	0.26

^a Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130)

^b The SEER 9 areas are San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah and Atlanta.
The SEER 13 areas comprise the SEER 9 areas plus San Jose-Monterey, Los Angeles, the Alaska Native Registry and Rural Georgia.
The SEER 18 areas comprise the SEER 13 areas plus California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG.

- Statistic not shown. Rate based on less than 16 cases for the time interval.