

Table 18.13
Myeloma

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years,
Lifetime Risk of Being Diagnosed with Cancer Given Alive and Cancer-Free at Current Age, and
Lifetime Risk of Dying from Cancer Given Alive at Current Age
Males, 2012-2014 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer
		+10 yrs	+20 yrs	+30 yrs	Ever				+10 yrs	+20 yrs	+30 yrs	Ever	
All Races	0	0.00	0.00	0.00	0.89	0.48	Asian/ Pacific Islander	0	0.00	0.00	0.00	0.63	0.32
	10	0.00	0.00	0.01	0.90	0.48		10	0.00	0.00	0.01	0.63	0.33
	20	0.00	0.01	0.04	0.90	0.49		20	0.00	0.01	0.02	0.64	0.33
	30	0.01	0.04	0.14	0.91	0.49		30	0.01	0.02	0.08	0.64	0.33
	40	0.03	0.13	0.34	0.92	0.50		40	0.02	0.07	0.19	0.64	0.33
	50	0.10	0.32	0.64	0.91	0.51		50	0.06	0.18	0.39	0.63	0.33
	60	0.23	0.58	0.82	0.87	0.51		60	0.13	0.34	0.53	0.59	0.32
	70	0.40	0.69	-	0.74	0.49		70	0.23	0.44	-	0.51	0.29
80	0.41	-	-	0.48	0.40	80	0.25	-	-	0.34	0.22		
White	0	0.00	0.00	0.00	0.84	0.46	American Indian/ Alaska Native ^a	0	0.00	0.00	0.00	0.51	0.35
	10	0.00	0.00	0.01	0.84	0.46		10	0.00	0.00	0.01	0.51	0.35
	20	0.00	0.01	0.04	0.85	0.47		20	0.00	0.01	0.04	0.52	0.36
	30	0.01	0.04	0.12	0.86	0.47		30	0.01	0.04	0.08	0.52	0.37
	40	0.03	0.12	0.31	0.86	0.48		40	0.03	0.07	0.22	0.53	0.38
	50	0.09	0.29	0.59	0.86	0.49		50	0.04	0.20	0.34	0.53	0.39
	60	0.21	0.54	0.78	0.82	0.49		60	0.17	0.32	0.52	0.53	0.41
	70	0.38	0.66	-	0.71	0.47		70	0.18	0.41	-	0.42	0.39
80	0.40	-	-	0.47	0.38	80	0.33	-	-	0.35	0.34		
Black	0	0.00	0.00	0.00	1.39	0.70	Hispanic ^b	0	0.00	0.00	0.00	0.94	0.45
	10	0.00	0.00	0.02	1.42	0.72		10	0.00	0.00	0.01	0.95	0.46
	20	0.00	0.02	0.08	1.43	0.72		20	0.00	0.01	0.04	0.95	0.46
	30	0.02	0.08	0.27	1.46	0.73		30	0.01	0.04	0.13	0.96	0.46
	40	0.07	0.26	0.66	1.48	0.75		40	0.03	0.13	0.32	0.96	0.47
	50	0.20	0.62	1.14	1.47	0.77		50	0.10	0.30	0.63	0.95	0.47
	60	0.47	1.05	1.38	1.43	0.78		60	0.21	0.57	0.86	0.91	0.47
	70	0.74	1.14	-	1.21	0.76		70	0.40	0.73	-	0.78	0.44
80	0.65	-	-	0.76	0.63	80	0.43	-	-	0.50	0.33		

Devcan 6.7.5, April 2017, National Cancer Institute (<https://surveillance.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, and Georgia excluding ATL/RG). Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

- Statistic could not be calculated.

A percent of 0.00 represents a value that is below 0.005.