

Table 19.17
Non-Hodgkin Lymphoma

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years,
Lifetime Risk of Being Diagnosed with Cancer Given Alive and Cancer-Free at Current Age, and
Lifetime Risk of Dying from Cancer Given Alive at Current Age
Males, 2012-2014 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer
		+10 yrs	+20 yrs	+30 yrs	Ever				+10 yrs	+20 yrs	+30 yrs	Ever	
All Races	0	0.01	0.03	0.07	2.38	0.84	Asian/ Pacific Islander	0	0.01	0.04	0.07	2.13	0.85
	10	0.02	0.06	0.12	2.39	0.85		10	0.03	0.06	0.11	2.14	0.85
	20	0.03	0.10	0.23	2.38	0.85		20	0.03	0.08	0.18	2.12	0.85
	30	0.07	0.20	0.47	2.37	0.86		30	0.05	0.15	0.34	2.10	0.85
	40	0.14	0.41	0.93	2.35	0.87		40	0.10	0.29	0.68	2.06	0.86
	50	0.29	0.82	1.57	2.28	0.88		50	0.19	0.58	1.19	1.99	0.86
	60	0.57	1.38	2.01	2.15	0.89		60	0.41	1.04	1.65	1.87	0.85
	70	0.94	1.67	-	1.84	0.87		70	0.69	1.36	-	1.60	0.81
White	80	1.04	-	-	1.29	0.75	American Indian/ Alaska Native ^a	80	0.82	-	-	1.13	0.68
	0	0.01	0.03	0.07	2.48	0.88		0	0.01	0.04	0.06	1.33	0.62
	10	0.02	0.05	0.12	2.49	0.89		10	0.02	0.05	0.09	1.34	0.63
	20	0.03	0.09	0.23	2.48	0.89		20	0.02	0.07	0.13	1.32	0.64
	30	0.06	0.20	0.48	2.48	0.90		30	0.05	0.11	0.23	1.33	0.65
	40	0.14	0.42	0.97	2.46	0.91		40	0.07	0.19	0.45	1.32	0.67
	50	0.30	0.86	1.65	2.39	0.92		50	0.13	0.40	0.69	1.32	0.68
	60	0.60	1.46	2.12	2.26	0.93		60	0.30	0.61	0.97	1.30	0.72
Black	70	1.00	1.76	-	1.93	0.91	Hispanic ^b	70	0.37	0.79	-	1.19	0.77
	80	1.10	-	-	1.33	0.78		80	0.60	-	-	1.17	0.83
	0	0.01	0.03	0.07	1.42	0.48		0	0.01	0.03	0.05	2.34	0.87
	10	0.02	0.06	0.15	1.44	0.49		10	0.02	0.04	0.10	2.35	0.88
	20	0.04	0.12	0.26	1.43	0.49		20	0.02	0.08	0.19	2.34	0.89
	30	0.08	0.23	0.47	1.41	0.49		30	0.06	0.17	0.42	2.34	0.89
	40	0.15	0.40	0.75	1.36	0.49		40	0.12	0.37	0.83	2.31	0.89
	50	0.26	0.63	0.99	1.27	0.49		50	0.26	0.74	1.46	2.25	0.89
	60	0.41	0.82	1.07	1.13	0.48		60	0.50	1.27	1.91	2.11	0.89
	70	0.52	0.84	-	0.92	0.46		70	0.86	1.58	-	1.81	0.86
	80	0.50	-	-	0.64	0.39		80	0.95	-	-	1.25	0.72

Devcan 6.7.5, April 2017, National Cancer Institute (<https://surveillance.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, and Georgia excluding ATL/RG). Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives.

- Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

- Statistic could not be calculated.

A percent of 0.00 represents a value that is below 0.005.