

Table 19.25
Non-Hodgkin LymphomaEstimated United States Cancer Prevalence Counts^a on January 1, 2014
By Race/Ethnicity, Sex and Years Since Diagnosis

WARNING: The Prevalence Estimates presented on this page utilize a different tumor selection criteria from previous CSR releases and are NOT directly comparable to estimates from previous CSR releases.
Please see the Prevalence Comparison Chapter of this CSR for more information.

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <22 ^e	0 to <39 ^e	>=39 ^g	Complete ^h	
Race	Sex										
All Races ^b	Both Sexes	247,549	171,171	107,339	61,534	32,564	19,328	615,303	653,661	8,335	661,996
	Males	134,164	89,969	55,681	31,794	16,734	10,167	325,886	346,187	4,460	350,647
	Females	113,385	81,202	51,658	29,740	15,830	9,161	289,417	307,474	3,875	311,349
White ^b	Both Sexes	213,854	150,511	94,695	54,077	29,075	17,504	537,797	572,570	8,781	581,351
	Males	116,394	79,369	49,345	27,844	14,948	9,226	285,660	304,105	5,049	309,154
	Females	97,460	71,142	45,350	26,233	14,127	8,278	252,137	268,465	3,732	272,197
Black ^b	Both Sexes	20,241	13,137	8,292	4,757	2,078	1,220	48,274	50,604	724	51,328
	Males	10,423	6,541	4,121	2,515	1,036	641	24,517	25,727	320	26,047
	Females	9,818	6,596	4,171	2,242	1,042	579	23,757	24,877	404	25,281
Asian/ Pacific	Both Sexes	7,657	4,695	2,905	1,605	+	+	17,244	+	+	+
	Males	4,140	2,454	1,466	835	+	+	9,081	+	+	+
Islander ^c	Females	3,517	2,241	1,439	770	+	+	8,163	+	+	+
Hispanic ^d	Both Sexes	19,630	11,818	7,221	3,986	+	+	43,564	+	+	+
	Males	10,215	6,134	3,741	2,154	+	+	22,702	+	+	+
	Females	9,415	5,684	3,480	1,832	+	+	20,862	+	+	+

Estimated prevalence percent^a on January 1, 2014, of the SEER^c population diagnosed in the previous 22 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)								Age-Adjusted ^f		
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
Race	Sex										
All Races ^c	Both Sexes	0.1698%	0.0032%	0.0126%	0.0269%	0.0493%	0.1043%	0.2167%	0.4262%	0.7404%	0.9295%
	Males	0.1825%	0.0039%	0.0175%	0.0329%	0.0570%	0.1217%	0.2481%	0.4839%	0.8492%	1.1279%
	Females	0.1574%	0.0025%	0.0074%	0.0206%	0.0415%	0.0871%	0.1865%	0.3741%	0.6513%	0.8100%
White ^c	Both Sexes	0.1923%	0.0031%	0.0130%	0.0286%	0.0505%	0.1109%	0.2320%	0.4639%	0.8169%	1.0158%
	Males	0.2057%	0.0040%	0.0184%	0.0350%	0.0582%	0.1300%	0.2640%	0.5217%	0.9297%	1.2247%
	Females	0.1790%	0.0023%	0.0073%	0.0217%	0.0423%	0.0912%	0.2002%	0.4098%	0.7220%	0.8882%
Black ^c	Both Sexes	0.1058%	0.0034%	0.0111%	0.0255%	0.0577%	0.1013%	0.1899%	0.3178%	0.4713%	0.4945%
	Males	0.1132%	0.0032%	0.0149%	0.0327%	0.0698%	0.1152%	0.2197%	0.3730%	0.5095%	0.5934%
	Females	0.0991%	0.0035%	0.0071%	0.0184%	0.0470%	0.0892%	0.1639%	0.2746%	0.4446%	0.4455%
Asian/ Pacific	Both Sexes	0.1024%	0.0032%	0.0118%	0.0200%	0.0339%	0.0655%	0.1367%	0.2558%	0.4282%	0.5914%
	Males	0.1117%	0.0040%	0.0151%	0.0235%	0.0375%	0.0734%	0.1587%	0.2924%	0.5153%	0.7550%
Islander ^c	Females	0.0938%	0.0025%	0.0083%	0.0164%	0.0305%	0.0584%	0.1177%	0.2260%	0.3602%	0.4909%
Hispanic ^d	Both Sexes	0.0820%	0.0023%	0.0107%	0.0209%	0.0354%	0.0800%	0.1682%	0.3344%	0.5519%	0.7044%
	Males	0.0843%	0.0023%	0.0145%	0.0254%	0.0407%	0.0913%	0.1829%	0.3652%	0.5875%	0.7970%
	Females	0.0797%	0.0023%	0.0067%	0.0160%	0.0297%	0.0682%	0.1540%	0.3077%	0.5254%	0.6484%

^a US 2014 cancer prevalence counts are based on 2014 cancer prevalence proportions from the SEER registries and 1/1/2014 US population estimates based on the average of 2013 and 2014 population estimates from the US Bureau of the Census.

^b ^c Prevalence was calculated using the first invasive tumor for each cancer site diagnosed during the previous 39/22 years.

Statistics based on (b) SEER 9 Areas (c) SEER 13 Areas excluding the Alaska Native Registry

(d) NHIA for Hispanic for SEER 13 Areas excluding the Alaska Native Registry.

^e Maximum limited-duration prevalence: 39 years for 1975-2014 SEER 9 data; 22 years for 1992-2014 SEER 13 data

(excluding the Alaska Native Registry) used to calculate prevalence for Hispanics and Asian Pacific Islanders.

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^g ^h ⁱ (g) Cases diagnosed more than 39 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <39 and >=39. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.