

Table 19.28  
All Lymphoid Neoplasms With Detailed Non-Hodgkin Lymphoma Subtypes<sup>a</sup>  
SEER<sup>b</sup> Incidence Rates and Annual Percent Change by Age at Diagnosis

All Races, Females, 2005-2014

Site	All Ages			Ages 0-19			Ages 20-64			Ages 65+		
	Rate	Count	APC	Rate	Count	APC	Rate	Count	APC	Rate	Count	APC
Lymphoid Neoplasm	29.1	137,637	-0.5*	5.2	5,948	0.6	19.3	54,845	-0.4	129.2	76,844	-0.7*
1 Hodgkin Lymphoma	2.4	10,419	-1.9*	1.1	1,322	-1.5	2.8	7,179	-1.7*	3.3	1,918	-2.8*
2 Non-Hodgkin lymphoma	25.9	123,290	-0.2	4.0	4,572	1.1	16.0	46,516	0.0	121.6	72,202	-0.5*
2(a) Non-Hodgkin lymphoma, B-cell	24.0	114,627	0.0	3.3	3,774	2.3*	14.5	42,365	0.2	115.3	68,488	-0.2
2(a) 1. Precursor Non-Hodgkin lymphoma, B-cell	1.3	5,581	3.3*	2.8	3,245	2.7*	0.6	1,673	4.3*	1.1	663	3.7
2(a) 2. Mature Non-Hodgkin lymphoma, B-cell	21.4	102,875	-0.4*	0.4	500	0.2	13.3	38,881	-0.2	107.0	63,494	-0.6*
2(a) 2.1. Chronic/Small/Prolymphocytic/ Mantle B-cell Non-Hodgkin lymphoma	4.6	22,363	-1.5*	-	-	-	2.2	6,826	-1.2	25.9	15,526	-1.7*
2(a) 2.1.1. Chronic/Small lymphocytic leukemia/lymphoma	4.2	20,231	-1.8*	-	-	-	2.0	6,136	-1.4*	23.5	14,087	-2.0*
2(a) 2.1.2. Prolymphocytic leukemia, B-cell	0.0	64	-	-	-	-	-	-	-	0.1	46	-
2(a) 2.1.3. Mantle-cell lymphoma	0.4	2,068	1.4	-	-	-	0.2	675	0.9	2.4	1,393	1.6
2(a) 2.2. Lymphoplasmacytic lymphoma/ Waldenstrom macroglobulinemia	0.5	2,292	-1.0	-	-	-	0.2	703	-2.5	2.7	1,587	-0.4
2(a) 2.2.1. Lymphoplasmacytic lymphoma	0.2	1,109	-0.2	-	-	-	0.1	348	0.0	1.3	760	-0.3
2(a) 2.2.2. Waldenstrom macroglobulinemia	0.2	1,183	-1.6*	-	-	-	0.1	355	-4.3*	1.4	827	-0.5
2(a) 2.3. Diffuse large B-cell lymphoma (DLBCL)	5.7	27,273	0.2	0.2	273	-2.1	3.6	10,303	0.4	28.1	16,697	0.1
2(a) 2.3.1. DLBCL, NOS <sup>c</sup>	5.7	26,896	0.0	0.2	250	-1.9	3.5	9,998	0.1	28.0	16,648	0.0
2(a) 2.3.2. Intravascular large B-cell lymphoma	0.0	51	-	-	-	-	0.0	23	-	0.0	28	-
2(a) 2.3.3. Primary effusion lymphoma	-	-	-	-	-	-	-	-	-	-	-	-
2(a) 2.3.4. Mediastinal large B-cell lymphoma	0.1	316	8.4*	0.0	23	-	0.1	279	9.3*	-	-	-
2(a) 2.4. Burkitt lymphoma/leukemia	0.2	943	-0.9	0.1	160	-	0.2	469	-0.8	0.5	314	-4.5
2(a) 2.5. Marginal-zone lymphoma (MZL)	1.9	9,173	0.4	0.0	32	-	1.3	3,717	1.1*	9.2	5,424	-0.1
2(a) 2.5.1. Splenic MZL	0.2	806	-1.2	-	-	-	0.1	300	-2.6	0.9	506	-0.4
2(a) 2.5.2. Extranodal MZL, MALT <sup>c</sup> type	1.2	5,597	1.6*	0.0	27	-	0.8	2,444	1.9*	5.3	3,126	1.3
2(a) 2.5.3. Nodal MZL	0.6	2,770	-1.6*	-	-	-	0.3	973	0.4	3.0	1,792	-2.5*
2(a) 2.6. Follicular lymphoma	3.3	15,663	-2.2*	-	-	-	2.6	7,621	-2.2*	13.7	8,028	-2.2*
2(a) 2.7. Hairy-cell leukemia	0.1	520	0.7	-	-	-	0.1	306	1.8	0.4	214	-1.6
2(a) 2.8. Plasma cell neoplasms	5.1	24,635	0.7	-	-	-	3.0	8,929	1.1	26.5	15,698	0.5
2(a) 2.8.1. Plasmacytoma	0.3	1,359	2.5*	-	-	-	0.2	618	3.9*	1.2	734	1.3
2(a) 2.8.2. Multiple myeloma/ plasma-cell leukemia	4.8	23,276	0.6	-	-	-	2.8	8,311	0.9	25.3	14,964	0.5
2(a) 2.9. Heavy chain disease	-	-	-	-	-	-	-	-	-	-	-	-
2(a) 3. Non-Hodgkin lymphoma, B-cell, NOS <sup>c</sup>	1.3	6,171	4.8*	0.0	29	-	0.6	1,811	4.1*	7.2	4,331	5.1*

The APC is the Annual Percent Change over the time interval.

<sup>a</sup> Morton LM, Turner JJ, Cerhan JR, Linet MS, Treseler PA, Clarke CA, Jack A, Cozen W, Maynadié M, Spinelli JJ, Costantini AS, Rüdiger T, Scarpa A, Zheng T, Weisenburger DD. Proposed classification of lymphoid neoplasms for epidemiologic research from the Pathology Working Group of the International Lymphoma Epidemiology Consortium (InterLymph). *Blood*. 2007;110:695-708.

<sup>b</sup> SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

<sup>c</sup> NOS = Not otherwise specified. MALT = Mucosa-associated lymphoid tissue. NK = Natural killer.

\* The APC is significantly different from zero (p<.05).

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Trend based on less than 10 cases for at least one year within the time interval.

Table 19.28 - continued  
 All Lymphoid Neoplasms With Detailed Non-Hodgkin Lymphoma Subtypes  
 SEER<sup>b</sup> Incidence Rates and Annual Percent Change by Age at Diagnosis

All Races, Females, 2005-2014

Site	All Ages			Ages 0-19			Ages 20-64			Ages 65+		
	Rate	Count	APC	Rate	Count	APC	Rate	Count	APC	Rate	Count	APC
2(b) Non-Hodgkin lymphoma, T-cell	1.6	7,546	0.5	0.5	577	6.6*	1.4	3,855	0.0	5.3	3,114	-0.1
2(b) 1. Precursor Non-Hodgkin lymphoma, T-cell	0.0	213	-	0.1	90	-	0.0	98	-	0.0	25	-
2(b) 2. Mature Non-Hodgkin lymphoma, T-cell	1.6	7,297	1.5*	0.4	486	13.3*	1.4	3,735	1.0	5.2	3,076	0.2
2(b) 2.1. Mycosis fungoides/Sezary syndrome	0.4	1,958	1.6	0.1	63	-	0.4	1,166	1.8	1.2	729	1.2
2(b) 2.1.1. Mycosis fungoides	0.4	1,899	1.5	0.1	63	-	0.4	1,147	1.6	1.2	689	1.3
2(b) 2.1.2. Sezary syndrome	0.0	59	-	-	-	-	0.0	19	-	0.1	40	-
2(b) 2.2. Peripheral T-cell lymphoma	0.9	4,174	0.1	0.1	162	-	0.7	2,017	-0.5	3.4	1,995	0.0
2(b) 2.2.1. Peripheral T-cell lymphoma, NOS <sup>c</sup>	0.3	1,449	2.1*	0.0	29	-	0.2	626	2.0	1.3	794	1.9
2(b) 2.2.2. Angioimmunoblastic T-cell lymphoma	0.1	610	-1.0	-	-	-	0.1	213	-4.8*	0.7	397	1.0
2(b) 2.2.3. Subcutaneous panniculitis-like T-cell lymphoma	0.0	88	-	0.0	17	-	0.0	54	-	0.0	17	-
2(b) 2.2.4. Anaplastic large cell lymphoma, T-cell or null-cell type	0.2	737	-2.4*	0.1	86	-	0.2	411	-1.9	0.4	240	-5.9*
2(b) 2.2.5. Hepatosplenic T-cell lymphoma	0.0	28	-	-	-	-	0.0	22	-	-	-	-
2(b) 2.2.6. Enteropathy-type T-cell lymphoma	0.0	53	-	-	-	-	0.0	23	-	0.1	30	-
2(b) 2.2.7. Cutaneous T-cell lymphoma, NOS <sup>c</sup>	0.2	833	1.6*	0.0	16	-	0.2	446	2.5*	0.6	371	-0.4
2(b) 2.2.8. Primary cutaneous anaplastic large cell lymphoma	0.1	376	-3.3	-	-	-	0.1	222	-5.3	0.2	142	-2.3
2(b) 2.3. Adult T-cell leukemia/lymphoma	0.2	691	13.2*	0.2	256	-	0.1	294	9.4*	0.2	141	-
2(b) 2.4. NK <sup>c</sup> /T-cell lymphoma, nasal-type/aggressive NK <sup>c</sup> -cell leukemia	0.1	256	2.9	-	-	-	0.1	171	-	0.1	81	-
2(b) 2.5. T-cell large granular lymphocytic leukemia	0.0	82	-	-	-	-	0.0	37	-	0.1	44	-
2(b) 2.6. Prolymphocytic leukemia, T-cell	0.0	136	-	-	-	-	0.0	50	-	0.1	86	-
2(b) 3. Non-Hodgkin lymphoma, NOS <sup>c</sup> , T-cell	0.0	36	-	-	-	-	0.0	22	-	-	-	-
2(c) Non-Hodgkin lymphoma, unknown lineage	0.2	1,117	-	0.2	221	-	0.1	296	-	1.0	600	-
2(c) 1. Precursor lymphoblastic leukemia/lymphoma, unknown lineage	0.1	409	-	0.2	216	-	0.0	104	-	0.1	89	-
2(c) 2. Prolymphocytic leukemia, unknown lineage	0.0	37	-	-	-	-	-	-	-	0.0	25	-
2(c) 3. Non-Hodgkin lymphoma, NOS <sup>c</sup> , unknown lineage	0.1	671	-	-	-	-	0.1	180	-	0.8	486	-
2(*) 1. Total precursor lymphoma/leukemia <sup>d</sup>	1.5	6,203	-0.1	3.1	3,551	-0.2	0.7	1,875	0.8	1.3	777	-1.1
3 Composite Hodgkin lymphoma and Non-Hodgkin lymphoma	0.0	149	-	-	-	-	0.0	68	-	0.1	78	-
4 Lymphoid neoplasm, NOS <sup>c</sup>	0.8	3,779	-4.9*	0.0	51	-	0.4	1,082	-4.3*	4.2	2,646	-5.5*

The APC is the Annual Percent Change over the time interval.

<sup>a</sup> Morton LM, et al. Proposed classification of lymphoid neoplasms for epidemiologic research from the Pathology Working Group of the International Lymphoma Epidemiology Consortium (InterLymph). Blood. 2007;110:695-708.

<sup>b</sup> SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

<sup>c</sup> NOS = Not otherwise specified. MALT = Mucosa-associated lymphoid tissue. NK = Natural killer.

<sup>d</sup> Total precursor lymphoma/leukemia is comprised of categories 2(a)1., 2(b)1., and 2(c)1.

\* The APC is significantly different from zero (p<.05).

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Trend based on less than 10 cases for at least one year within the time interval.