

Table 20.22
Cancer of the Oral Cavity and Pharynx (Invasive)

Age-Adjusted SEER Incidence Rates^a
By Registry, Race and Sex

	All Races			Whites			Blacks		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
<u>SEER Incidence Rates^a, 2010-2014</u>									
California	10.31	15.57	5.73	10.89	16.49	5.90	8.36	12.79	4.72
Greater Bay Area	10.35	15.30	5.98	11.11	16.51	6.16	8.84	14.42	4.14
San Francisco-Oakland	10.68	15.90	6.08	11.36	16.95	6.20	9.01	14.91	4.22
San Jose-Monterey	9.76	14.21	5.82	10.66	15.72	6.08	8.02	12.19	-
Los Angeles	9.09	13.79	5.13	9.56	14.38	5.39	8.72	13.55	5.10
Greater California	10.86	16.46	5.91	11.36	17.33	6.03	7.85	11.39	4.65
Connecticut	11.40	16.65	6.88	11.74	17.07	7.06	7.87	12.82	4.29
Detroit	12.25	19.02	6.53	12.59	19.41	6.56	11.62	18.49	6.53
Georgia	12.20	18.47	6.93	13.42	20.14	7.46	8.83	13.57	5.30
Atlanta	10.94	16.60	6.34	12.76	19.08	7.18	7.81	12.52	4.50
Rural Georgia	13.41	21.97	6.54	15.63	26.16	6.22	9.55	12.26	-
Greater Georgia	12.83	19.30	7.30	13.65	20.45	7.62	9.66	14.45	5.93
Hawaii	12.38	18.08	7.22	17.03	24.48	8.82	-	-	-
Iowa	12.48	18.89	6.78	12.50	18.90	6.81	8.97	11.30	-
Kentucky	14.08	21.76	7.41	14.19	21.87	7.49	10.62	18.49	4.25
Louisiana	13.08	20.58	6.63	14.05	21.89	7.05	10.67	17.30	5.48
New Jersey	10.38	15.62	6.02	10.74	16.03	6.25	7.88	12.30	4.79
New Mexico	9.31	14.43	4.75	9.82	15.23	4.96	-	-	-
Seattle-Puget Sound	12.11	17.49	7.29	12.67	18.27	7.57	7.48	11.84	-
Utah	8.53	12.31	5.08	8.52	12.32	5.03	-	-	-
SEER 9 Areas ^b	11.26	16.81	6.41	11.78	17.53	6.59	8.98	14.31	4.89
SEER 13 Areas ^b	10.71	16.01	6.10	11.26	16.78	6.30	8.92	14.07	4.96
SEER 18 Areas ^b	11.16	16.85	6.23	11.70	17.64	6.42	9.08	14.20	5.10

^a Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130)

^b The SEER 9 areas are San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah and Atlanta.

The SEER 13 areas comprise the SEER 9 areas plus San Jose-Monterey, Los Angeles, the Alaska Native Registry and Rural Georgia.

The SEER 18 areas comprise the SEER 13 areas plus California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG.

- Statistic not shown. Rate based on less than 16 cases for the time interval.