


Table 20.25
Cancer of the Oral Cavity and Pharynx (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2010-2014

Males

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	3.83	0.02		
High Five States				
Mississippi	6.06	0.29	01(01,04)	58.01 ^c
Louisiana	5.44	0.22	02(01,08)	41.90 ^c
D.C.	5.27	0.64	03(01,38)	37.52 ^c
Tennessee	5.03	0.18	04(02,11)	31.19 ^c
Arkansas	5.02	0.26	05(01,13)	30.92 ^c
Low Five States				
Kansas	3.05	0.21	47(29,51)	-20.46 ^c
North Dakota	2.87	0.40	48(14,51)	-25.13 ^c
Rhode Island	2.73	0.32	49(27,51)	-28.75 ^c
Colorado	2.66	0.15	50(43,51)	-30.72 ^c
Utah	2.35	0.22	51(45,51)	-38.63 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	4.25	0.19	13(06,30)	10.91	Montana	3.43	0.34	39(09,50)	-10.48
Alaska	3.70	0.55	26(02,51)	-3.58	Nebraska	3.35	0.27	41(15,50)	-12.71
Arizona	3.07	0.13	46(34,50)	-19.81 ^c	Nevada	3.64	0.24	31(10,47)	-5.01
Arkansas	5.02	0.26	05(01,13)	30.92 ^c	New Hampshire	3.54	0.31	35(09,50)	-7.64
California	3.73	0.07	25(20,36)	-2.82	New Jersey	3.15	0.12	43(33,49)	-17.74 ^c
Colorado	2.66	0.15	50(43,51)	-30.72 ^c	New Mexico	3.52	0.26	37(11,49)	-8.07
Connecticut	3.12	0.18	44(28,50)	-18.61 ^c	New York	3.31	0.08	42(32,46)	-13.58
Delaware	4.45	0.43	10(02,43)	16.13 ^c	North Carolina	4.11	0.13	17(09,30)	7.15
D.C.	5.27	0.64	03(01,38)	37.52 ^c	North Dakota	2.87	0.40	48(14,51)	-25.13 ^c
Florida	4.10	0.08	18(10,25)	7.06	Ohio	3.93	0.11	21(12,34)	2.63
Georgia	4.14	0.14	16(08,29)	8.09	Oklahoma	4.45	0.22	09(04,25)	16.20 ^c
Hawaii	4.17	0.34	14(03,42)	8.70	Oregon	3.59	0.18	32(15,46)	-6.39
Idaho	3.54	0.30	34(10,49)	-7.53	Pennsylvania	3.38	0.10	40(29,46)	-11.93
Illinois	3.80	0.11	24(15,37)	-0.87	Rhode Island	2.73	0.32	49(27,51)	-28.75 ^c
Indiana	3.67	0.15	27(15,42)	-4.29	South Carolina	4.68	0.20	08(03,17)	22.16 ^c
Iowa	3.66	0.21	28(11,45)	-4.49	South Dakota	3.45	0.40	38(08,51)	-9.93
Kansas	3.05	0.21	47(29,51)	-20.46 ^c	Tennessee	5.03	0.18	04(02,11)	31.19 ^c
Kentucky	4.78	0.21	07(02,16)	24.62 ^c	Texas	4.15	0.09	15(09,24)	8.14
Louisiana	5.44	0.22	02(01,08)	41.90 ^c	Utah	2.35	0.22	51(45,51)	-38.63 ^c
Maine	4.39	0.33	11(02,38)	14.65	Vermont	4.28	0.50	12(02,48)	11.54
Maryland	3.53	0.16	36(18,46)	-7.83	Virginia	3.65	0.14	29(16,42)	-4.81
Massachusetts	3.83	0.15	23(11,39)	-0.12	Washington	3.97	0.15	20(10,35)	3.70
Michigan	4.03	0.12	19(09,32)	5.11	West Virginia	4.90	0.30	06(01,20)	27.70 ^c
Minnesota	3.11	0.15	45(32,50)	-18.77 ^c	Wisconsin	3.65	0.16	30(16,43)	-4.90
Mississippi	6.06	0.29	01(01,04)	58.01 ^c	Wyoming	3.57	0.50	33(03,51)	-6.88
Missouri	3.86	0.16	22(11,38)	0.70					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.