

Table 21.7  
Cancer of the Ovary (Invasive)

SEER Incidence<sup>a</sup> and U.S. Death<sup>b</sup> Rates, Age-Adjusted and Age-Specific Rates, by Race

	SEER Incidence <sup>a</sup>			U.S. Mortality <sup>b</sup>		
	All Races Females	White Females	Black Females	All Races Females	White Females	Black Females
<u>Age-Adjusted Rates, 2010-2014</u>						
Age at Diagnosis/Death:						
All ages	11.7	12.2	9.4	7.4	7.7	6.4
Under 65	7.3	7.6	5.7	2.9	3.0	2.5
65 and over	41.9	44.5	35.3	38.6	40.1	33.4
All ages (WHO world std) <sup>c</sup>	9.2	9.6	7.4	5.3	5.5	4.6
<u>Age-Specific Rates, 2010-2014</u>						
Age at Diagnosis/Death:						
<1	-	-	-	-	-	-
1-4	-	-	-	-	-	-
5-9	0.3	0.2	-	-	-	-
10-14	0.8	0.7	-	-	-	-
15-19	1.5	1.5	1.5	0.1	0.1	-
20-24	1.8	1.7	1.5	0.2	0.2	-
25-29	2.3	2.4	2.0	0.3	0.3	0.3
30-34	3.3	3.5	2.9	0.5	0.5	0.3
35-39	4.5	4.7	3.1	1.1	1.1	1.1
40-44	8.9	9.3	5.6	2.2	2.2	2.0
45-49	14.5	14.6	11.1	4.6	4.8	4.0
50-54	20.0	20.7	14.8	8.3	8.7	6.8
55-59	24.0	25.2	18.9	12.4	12.9	10.9
60-64	29.1	30.8	25.4	18.5	19.2	17.1
65-69	36.0	38.3	29.5	26.0	27.1	22.9
70-74	40.9	43.9	35.4	34.3	35.6	31.8
75-79	45.2	48.0	37.1	43.1	45.1	36.3
80-84	47.0	50.0	38.3	51.1	52.9	43.6
85+	45.6	47.4	41.6	52.8	54.7	43.1

- <sup>a</sup> Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473. SEER 18 areas. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.
- <sup>b</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.
- <sup>c</sup> Rates are per 100,000 and are age-adjusted to the world (WHO 2000-2025) standard million.
- Statistic not shown. Rate based on less than 16 cases for the time interval.