

Table 23.7  
Cancer of the Prostate (Invasive)

SEER Incidence<sup>a</sup> and U.S. Death<sup>b</sup> Rates, Age-Adjusted and Age-Specific Rates, by Race

	SEER Incidence <sup>a</sup>			U.S. Mortality <sup>b</sup>		
	All Races Males	White Males	Black Males	All Races Males	White Males	Black Males
<u>Age-Adjusted Rates, 2010-2014</u>						
Age at Diagnosis/Death:						
All ages	119.8	112.8	188.7	20.1	18.7	42.0
Under 65	47.7	43.9	87.3	1.7	1.4	4.1
65 and over	618.5	589.5	889.1	147.1	137.6	303.9
All ages (WHO world std) <sup>c</sup>	91.2	85.7	146.7	11.2	10.3	24.1
<u>Age-Specific Rates, 2010-2014</u>						
Age at Diagnosis/Death:						
<1	-	-	-	-	-	-
1-4	-	-	-	-	-	-
5-9	-	-	-	-	-	-
10-14	-	-	-	-	-	-
15-19	-	-	-	-	-	-
20-24	-	-	-	-	-	-
25-29	-	-	-	-	-	-
30-34	-	-	-	-	-	-
35-39	0.8	0.7	2.1	0.0	0.0	-
40-44	8.8	7.6	21.5	0.2	0.1	0.4
45-49	38.8	32.6	95.4	0.9	0.8	2.1
50-54	123.4	112.1	238.4	3.2	2.7	7.6
55-59	265.3	244.1	479.2	8.7	7.3	21.2
60-64	450.0	423.6	755.2	19.7	17.0	48.2
65-69	693.4	658.5	1,074.5	39.2	34.6	96.1
70-74	718.4	687.1	1,022.6	72.6	65.4	172.0
75-79	632.1	600.2	868.3	129.6	119.1	283.7
80-84	468.8	448.2	614.3	226.1	212.3	475.6
85+	396.4	381.4	558.3	477.7	459.6	870.9

<sup>a</sup> SEER 18 areas. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.

<sup>b</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

<sup>c</sup> Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.

- Rates are per 100,000 and are age-adjusted to the world (WHO 2000-2025) standard million.

- Statistic not shown. Rate based on less than 16 cases for the time interval.