Table 23.11 Cancer of the Prostate (Invasive)

SEER Incidence and U.S. Mortality Age-Adjusted Rates and Trends^a By Race/Ethnicity

SEER Incidence	SEER 18 Areas ^b Rate 2010-2014 Rate per 100,000 persons	SEER 13 Areas ^b Trend 2005-2014 [°] AAPC (%)	SEER 9 Areas ^b Trend 2005-2014 ^d AAPC (%)
	Males	Males	Males
RACE/ETHNICITY			
All Races	119.8	-5.9*	-5.8*
White	112.8	-6.2*	-6.1*
White Hispanic ^e	95.6	-6.8*	-
White Non-Hispanic ^e	116.0	-6.1*	-
Black	188.7	-5.6*	-5.5*
Asian/Pacific Islander	62.9	-7.2*	-
Amer Ind/Alaska Nat ^f	59.9	-6.2*	-
Hispanic ^e	98.3	-6.3*	-

	Rate 2010-2014 Rate per 100,000 persons	Trend 2005-2014 ^h AAPC (%)
	Males	Males
RACE/ETHNICITY		
All Races	20.1	-3.5*
White	18.7	-3.3*
White Hispanic ^e	17.2	-3.3*
White Non-Hispanic ^e	18.7	-3.0*
Black	42.0	-4.3*
Asian/Pacific Islander	8.8	-3.4*
Amer Ind/Alaska Nat		
Total U.S.	14.9	-2.6*
CHSDA Counties	19.4	-1.7*
Non-CHSDA Counties	10.0	-3.8*
Hispanic ^e	16.5	-2.8*

The AAPC is the Average Annual Percent Change over the time interval. The AAPCs are calculated by the Joinpoint Regression Program Version 4.3.0.0, April 2016, National Cancer Institute. Statistic not shown. Rate based on less than 16 cases for the time interval.

- Trend based on less than 10 cases for at least one year within the time interval. Rates are age-adjusted to the 2000 US Std Population (19 age groups Census P25-1130). Trends a
- are based on rates age-adjusted to the 2000 US Std Population (19 age groups Census P25-1130). The SEER 9 areas are San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, b Utah, and Atlanta.

The SEER 13 areas comprise the SEER 9 areas plus San Jose-Monterey, Los Angeles, the Alaska Native Registry, and Rural Georgia.

The SEER 18 areas comprise the SEER 13 areas plus California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, and Georgia excluding ATL/RG. The 2005-2014 AAPC estimates are based on a Joinpoint analysis with up to 4 Joinpoints over

С diagnosis years 1992-2014.

The 2005-2014 AAPC estimates are based on a Joinpoint analysis with up to 5 Joinpoints over d diagnosis years 1975-2014.

e Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry. The 2005-2014 Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma. f

Incidence data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

q US Mortality Files, National Center for Health Statistics, CDC. h

The 2005-2014 mortality AAPCs are based on a Joinpoint analysis using years of death 1992-2014. The APC is significantly different from zero (p<.05).

U.S. Mortality^g