


Table 23.14  
Cancer of the Prostate (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2010-2014

Males

State	Rate	SE	Rank(CI) <sup>d</sup>	PD
TOTAL U.S.	20.07	0.05		
High Five States				
D.C.	33.70	1.77	01(01,01)	67.94 <sup>c</sup>
Mississippi	26.17	0.67	02(02,03)	30.44 <sup>c</sup>
Alabama	23.85	0.49	03(02,08)	18.87 <sup>c</sup>
South Carolina	23.46	0.50	04(03,10)	16.94 <sup>c</sup>
Georgia	23.39	0.39	05(03,09)	16.56 <sup>c</sup>
Low Five States				
Arizona	18.24	0.34	47(39,50)	-9.11
West Virginia	18.22	0.62	48(29,50)	-9.19
Missouri	18.19	0.36	49(38,50)	-9.34
Florida	17.61	0.17	50(46,50)	-12.25 <sup>c</sup>
Hawaii	13.72	0.62	51(51,51)	-31.62 <sup>c</sup>

State	Rate	SE	Rank(CI) <sup>d</sup>	PD	State	Rate	SE	Rank(CI) <sup>d</sup>	PD
Alabama	23.85	0.49	03(02,08)	18.87 <sup>c</sup>	Montana	21.40	0.90	12(03,44)	6.64
Alaska	20.50	1.57	25(03,50)	2.17	Nebraska	20.87	0.69	19(06,44)	3.99
Arizona	18.24	0.34	47(39,50)	-9.11	Nevada	21.14	0.63	14(06,41)	5.34
Arkansas	20.78	0.56	20(07,42)	3.57	New Hampshire	19.84	0.81	30(08,49)	-1.11
California	20.02	0.16	28(22,38)	-0.23	New Jersey	19.44	0.31	40(24,46)	-3.12
Colorado	21.67	0.48	10(05,28)	7.99	New Mexico	20.69	0.67	23(07,44)	3.13
Connecticut	18.60	0.46	45(30,50)	-7.32	New York	19.49	0.21	39(27,44)	-2.87
Delaware	19.21	0.95	42(08,50)	-4.27	North Carolina	21.56	0.34	11(07,25)	7.46
D.C.	33.70	1.77	01(01,01)	67.94 <sup>c</sup>	North Dakota	19.83	1.06	31(05,50)	-1.18
Florida	17.61	0.17	50(46,50)	-12.25 <sup>c</sup>	Ohio	19.95	0.27	29(20,42)	-0.55
Georgia	23.39	0.39	05(03,09)	16.56 <sup>c</sup>	Oklahoma	20.78	0.50	21(08,40)	3.55
Hawaii	13.72	0.62	51(51,51)	-31.62 <sup>c</sup>	Oregon	21.23	0.47	13(06,34)	5.80
Idaho	23.32	0.83	06(02,20)	16.21 <sup>c</sup>	Pennsylvania	19.76	0.24	36(23,43)	-1.53
Illinois	20.91	0.28	18(10,31)	4.21	Rhode Island	19.79	0.87	34(07,50)	-1.39
Indiana	20.99	0.39	17(08,34)	4.59	South Carolina	23.46	0.50	04(03,10)	16.94 <sup>c</sup>
Iowa	19.78	0.50	35(14,47)	-1.40	South Dakota	19.52	0.96	38(07,50)	-2.70
Kansas	19.11	0.54	43(21,50)	-4.74	Tennessee	20.74	0.40	22(09,37)	3.35
Kentucky	19.82	0.48	32(15,46)	-1.24	Texas	18.68	0.21	44(38,49)	-6.90
Louisiana	22.46	0.51	07(03,18)	11.94 <sup>c</sup>	Utah	22.14	0.71	08(03,29)	10.36 <sup>c</sup>
Maine	19.80	0.75	33(08,49)	-1.34	Vermont	21.09	1.17	15(03,49)	5.13
Maryland	20.31	0.42	27(12,42)	1.23	Virginia	21.06	0.37	16(08,32)	4.94
Massachusetts	19.43	0.36	41(23,47)	-3.17	Washington	20.45	0.38	26(12,40)	1.89
Michigan	19.54	0.29	37(24,45)	-2.61	West Virginia	18.22	0.62	48(29,50)	-9.19
Minnesota	20.65	0.41	24(09,38)	2.90	Wisconsin	21.80	0.40	09(06,24)	8.64
Mississippi	26.17	0.67	02(02,03)	30.44 <sup>c</sup>	Wyoming	18.53	1.23	46(08,50)	-7.65
Missouri	18.19	0.36	49(38,50)	-9.34					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p < .0002$ ).

<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 10% or more.

<sup>d</sup> Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.