


Table 24.18
Cancer of the Stomach (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2010-2014

Males and Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	3.23	0.01		
High Five States				
D.C.	5.01	0.42	01(01,09)	55.04 ^c
Hawaii	4.84	0.24	02(01,04)	49.79 ^c
Alaska	4.15	0.41	03(01,24)	28.42 ^c
Louisiana	4.15	0.13	04(02,09)	28.30 ^c
California	4.12	0.05	05(02,07)	27.36 ^c
Low Five States				
Iowa	2.13	0.11	47(40,51)	-34.28 ^c
Wyoming	2.10	0.27	48(24,51)	-35.09 ^c
Maine	2.09	0.15	49(39,51)	-35.31 ^c
Utah	2.04	0.14	50(40,51)	-36.80 ^c
Nebraska	1.95	0.14	51(42,51)	-39.69 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	3.40	0.11	16(09,22)	5.08	Montana	2.19	0.19	45(34,51)	-32.15 ^c
Alaska	4.15	0.41	03(01,24)	28.42 ^c	Nebraska	1.95	0.14	51(42,51)	-39.69 ^c
Arizona	2.79	0.09	33(22,41)	-13.75	Nevada	2.94	0.15	28(15,40)	-9.06
Arkansas	2.95	0.13	26(16,40)	-8.94	New Hampshire	2.15	0.17	46(36,51)	-33.54 ^c
California	4.12	0.05	05(02,07)	27.36 ^c	New Jersey	3.50	0.08	13(08,18)	8.20
Colorado	2.61	0.10	38(25,44)	-19.16 ^c	New Mexico	3.85	0.19	07(03,16)	19.02 ^c
Connecticut	3.27	0.12	17(10,29)	1.04	New York	3.84	0.06	08(05,11)	18.57 ^c
Delaware	3.47	0.26	14(03,34)	7.19	North Carolina	3.03	0.08	22(16,33)	-6.17
D.C.	5.01	0.42	01(01,09)	55.04 ^c	North Dakota	2.62	0.26	37(15,51)	-18.91 ^c
Florida	2.89	0.05	30(22,36)	-10.79	Ohio	2.77	0.06	34(25,39)	-14.24
Georgia	3.44	0.09	15(09,19)	6.25	Oklahoma	2.95	0.12	27(16,38)	-8.94
Hawaii	4.84	0.24	02(01,04)	49.79 ^c	Oregon	2.68	0.11	35(23,44)	-17.11 ^c
Idaho	2.26	0.17	43(32,51)	-30.05 ^c	Pennsylvania	2.90	0.06	29(20,36)	-10.36
Illinois	3.50	0.07	12(09,17)	8.30	Rhode Island	3.55	0.24	11(03,31)	9.89
Indiana	2.64	0.09	36(27,43)	-18.42 ^c	South Carolina	3.82	0.12	09(03,14)	18.23 ^c
Iowa	2.13	0.11	47(40,51)	-34.28 ^c	South Dakota	2.40	0.22	42(19,51)	-25.73 ^c
Kansas	2.24	0.12	44(38,51)	-30.59 ^c	Tennessee	3.05	0.09	21(15,33)	-5.64
Kentucky	2.95	0.11	25(16,38)	-8.84	Texas	3.66	0.06	10(07,14)	13.04
Louisiana	4.15	0.13	04(02,09)	28.30 ^c	Utah	2.04	0.14	50(40,51)	-36.80 ^c
Maine	2.09	0.15	49(39,51)	-35.31 ^c	Vermont	2.49	0.25	41(16,51)	-22.93 ^c
Maryland	3.18	0.10	18(13,29)	-1.67	Virginia	3.03	0.09	23(16,34)	-6.28
Massachusetts	3.12	0.09	19(15,31)	-3.42	Washington	2.85	0.09	31(19,39)	-11.86
Michigan	2.97	0.07	24(17,34)	-8.16	West Virginia	3.09	0.16	20(11,37)	-4.37
Minnesota	2.54	0.09	40(31,45)	-21.52 ^c	Wisconsin	2.81	0.09	32(20,40)	-12.98
Mississippi	3.99	0.16	06(02,12)	23.46 ^c	Wyoming	2.10	0.27	48(24,51)	-35.09 ^c
Missouri	2.59	0.09	39(30,44)	-20.04 ^c					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.