


Table 24.19
Cancer of the Stomach (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2010-2014

Males

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	4.38	0.02		
High Five States				
D.C.	7.37	0.81	01(01,08)	68.35 ^c
Hawaii	6.28	0.42	02(01,10)	43.38 ^c
Louisiana	5.74	0.24	03(01,10)	31.01 ^c
South Carolina	5.42	0.22	04(02,14)	23.91 ^c
California	5.39	0.08	05(03,09)	23.09 ^c
Low Five States				
South Dakota	2.89	0.36	47(26,51)	-34.05 ^c
Nebraska	2.73	0.25	48(39,51)	-37.61 ^c
Maine	2.64	0.26	49(40,51)	-39.69 ^c
Montana	2.62	0.31	50(37,51)	-40.19 ^c
Utah	2.61	0.23	51(40,51)	-40.32 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	4.70	0.21	15(06,25)	7.29	Montana	2.62	0.31	50(37,51)	-40.19 ^c
Alaska	5.31	0.68	06(01,41)	21.35 ^c	Nebraska	2.73	0.25	48(39,51)	-37.61 ^c
Arizona	3.68	0.15	36(22,43)	-15.98 ^c	Nevada	3.80	0.25	33(15,44)	-13.20
Arkansas	3.83	0.23	31(16,43)	-12.40	New Hampshire	3.09	0.30	44(27,51)	-29.31 ^c
California	5.39	0.08	05(03,09)	23.09 ^c	New Jersey	4.85	0.15	12(06,19)	10.88
Colorado	3.55	0.18	38(23,46)	-18.84 ^c	New Mexico	4.95	0.31	09(03,25)	13.00
Connecticut	4.56	0.23	17(07,31)	4.06	New York	5.29	0.11	07(03,11)	20.76 ^c
Delaware	4.60	0.45	16(02,42)	5.17	North Carolina	4.00	0.14	27(17,38)	-8.71
D.C.	7.37	0.81	01(01,08)	68.35 ^c	North Dakota	3.23	0.42	41(15,51)	-26.29 ^c
Florida	3.80	0.08	32(24,39)	-13.12	Ohio	3.74	0.11	35(24,41)	-14.53
Georgia	4.70	0.16	14(07,22)	7.34	Oklahoma	4.31	0.22	19(09,37)	-1.47
Hawaii	6.28	0.42	02(01,10)	43.38 ^c	Oregon	3.55	0.19	39(22,46)	-18.98 ^c
Idaho	3.16	0.29	43(24,51)	-27.72 ^c	Pennsylvania	4.08	0.11	26(17,35)	-6.70
Illinois	4.92	0.13	10(06,17)	12.39	Rhode Island	4.84	0.42	13(02,37)	10.53
Indiana	3.78	0.16	34(20,42)	-13.59	South Carolina	5.42	0.22	04(02,14)	23.91 ^c
Iowa	2.94	0.19	46(38,51)	-32.87 ^c	South Dakota	2.89	0.36	47(26,51)	-34.05 ^c
Kansas	3.18	0.22	42(30,50)	-27.25 ^c	Tennessee	4.26	0.17	20(12,34)	-2.71
Kentucky	4.19	0.20	22(11,38)	-4.26	Texas	4.88	0.10	11(07,17)	11.52
Louisiana	5.74	0.24	03(01,10)	31.01 ^c	Utah	2.61	0.23	51(40,51)	-40.32 ^c
Maine	2.64	0.26	49(40,51)	-39.69 ^c	Vermont	3.92	0.49	29(05,50)	-10.42
Maryland	4.39	0.19	18(09,32)	0.20	Virginia	4.16	0.15	23(14,35)	-4.96
Massachusetts	4.12	0.16	24(15,37)	-5.95	Washington	3.98	0.16	28(17,40)	-9.07
Michigan	4.09	0.13	25(16,36)	-6.50	West Virginia	4.20	0.28	21(09,41)	-4.04
Minnesota	3.50	0.17	40(26,46)	-20.11 ^c	Wisconsin	3.84	0.16	30(19,42)	-12.24
Mississippi	5.17	0.28	08(02,19)	18.15 ^c	Wyoming	2.94	0.48	45(17,51)	-32.82 ^c
Missouri	3.66	0.16	37(23,44)	-16.44 ^c					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.