


Table 24.20  
Cancer of the Stomach (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2010-2014

Females

State	Rate	SE	Rank(CI) <sup>d</sup>	PD
TOTAL U.S.	2.34	0.02		
High Five States				
Hawaii	3.67	0.28	01(01,07)	56.77 <sup>c</sup>
D.C.	3.47	0.46	02(01,25)	48.01 <sup>c</sup>
California	3.13	0.06	03(01,07)	33.87 <sup>c</sup>
Mississippi	3.03	0.18	04(01,14)	29.19 <sup>c</sup>
Alaska	2.99	0.48	05(01,41)	27.54 <sup>c</sup>
Low Five States				
Wyoming	1.46	0.31	47(13,51)	-37.50 <sup>c</sup>
Idaho	1.46	0.18	48(28,51)	-37.80 <sup>c</sup>
Vermont	1.43	0.26	49(17,51)	-39.07 <sup>c</sup>
New Hampshire	1.41	0.18	50(30,51)	-39.80 <sup>c</sup>
Nebraska	1.31	0.15	51(39,51)	-44.20 <sup>c</sup>

State	Rate	SE	Rank(CI) <sup>d</sup>	PD	State	Rate	SE	Rank(CI) <sup>d</sup>	PD
Alabama	2.43	0.13	15(06,30)	3.91	Montana	1.80	0.24	40(11,51)	-22.92 <sup>c</sup>
Alaska	2.99	0.48	05(01,41)	27.54 <sup>c</sup>	Nebraska	1.31	0.15	51(39,51)	-44.20 <sup>c</sup>
Arizona	2.04	0.10	29(17,41)	-12.89	Nevada	2.22	0.18	22(08,42)	-5.17
Arkansas	2.25	0.16	21(08,40)	-3.96	New Hampshire	1.41	0.18	50(30,51)	-39.80 <sup>c</sup>
California	3.13	0.06	03(01,07)	33.87 <sup>c</sup>	New Jersey	2.51	0.09	12(06,23)	7.13
Colorado	1.86	0.12	37(21,47)	-20.48 <sup>c</sup>	New Mexico	2.95	0.22	06(01,19)	25.99 <sup>c</sup>
Connecticut	2.26	0.14	20(09,38)	-3.47	New York	2.77	0.07	08(04,12)	18.37 <sup>c</sup>
Delaware	2.51	0.29	13(02,43)	7.09	North Carolina	2.28	0.09	18(11,32)	-2.58
D.C.	3.47	0.46	02(01,25)	48.01 <sup>c</sup>	North Dakota	2.09	0.31	28(04,50)	-10.71
Florida	2.13	0.06	25(18,35)	-9.10	Ohio	2.01	0.07	30(20,41)	-14.19
Georgia	2.49	0.10	14(07,24)	6.21	Oklahoma	1.86	0.13	38(20,48)	-20.57 <sup>c</sup>
Hawaii	3.67	0.28	01(01,07)	56.77 <sup>c</sup>	Oregon	1.95	0.13	34(17,46)	-16.86 <sup>c</sup>
Idaho	1.46	0.18	48(28,51)	-37.80 <sup>c</sup>	Pennsylvania	2.00	0.07	32(21,40)	-14.50
Illinois	2.43	0.08	16(09,24)	3.77	Rhode Island	2.62	0.27	10(02,38)	11.87
Indiana	1.77	0.10	41(28,48)	-24.41 <sup>c</sup>	South Carolina	2.52	0.13	11(05,27)	7.77
Iowa	1.52	0.12	46(34,51)	-35.11 <sup>c</sup>	South Dakota	1.93	0.27	35(07,51)	-17.48 <sup>c</sup>
Kansas	1.53	0.13	45(32,51)	-34.59 <sup>c</sup>	Tennessee	2.11	0.10	26(14,40)	-9.89
Kentucky	1.97	0.12	33(17,45)	-15.84 <sup>c</sup>	Texas	2.70	0.07	09(05,14)	15.37 <sup>c</sup>
Louisiana	2.95	0.15	07(02,14)	25.85 <sup>c</sup>	Utah	1.55	0.16	44(28,51)	-33.88 <sup>c</sup>
Maine	1.61	0.18	43(24,51)	-31.38 <sup>c</sup>	Vermont	1.43	0.26	49(17,51)	-39.07 <sup>c</sup>
Maryland	2.26	0.11	19(10,36)	-3.37	Virginia	2.17	0.10	24(13,37)	-7.41
Massachusetts	2.38	0.10	17(08,29)	1.55	Washington	1.92	0.10	36(21,44)	-17.98 <sup>c</sup>
Michigan	2.11	0.08	27(17,38)	-10.08	West Virginia	2.18	0.19	23(07,43)	-6.72
Minnesota	1.81	0.11	39(24,48)	-22.88 <sup>c</sup>	Wisconsin	2.01	0.11	31(17,42)	-14.27
Mississippi	3.03	0.18	04(01,14)	29.19 <sup>c</sup>	Wyoming	1.46	0.31	47(13,51)	-37.50 <sup>c</sup>
Missouri	1.74	0.10	42(30,49)	-25.90 <sup>c</sup>					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p < .0002$ ).

<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.

<sup>d</sup> Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.