

CANCER STATISTICS REVIEW 1975-2015: INTRODUCTION

The annual *SEER Cancer Statistics Review (CSR)* contains incidence, mortality, prevalence, and survival statistics from 1975 through the most recent year for which data are available. This report is published by the Surveillance Research Program of the National Cancer Institute, which manages the Surveillance, Epidemiology, and End Results (SEER) Program. The scope and purpose of the **CSR** follow a report to the Senate Appropriations Committee (Breslow, 1988), which recommended that a broad profile of cancer be presented regularly to the American public.

The SEER program is an authoritative source of information on cancer incidence and survival in the United States. SEER collects and publishes these statistics from population-based registries covering 30% of the US population. The 18 SEER registries routinely collect data on patient demographics, primary tumor site, tumor morphology, extent of disease, first course of treatment, and active follow-up for vital status. Detailed information describing these fields can be found at <https://seer.cancer.gov/resources/>.

This report presents statistics on 29 primary sites and subsites, organized into site-specific chapters. Detailed statistics on cancer incidence, mortality, survival, and prevalence are reported by sex, race and ethnicity, age, stage at diagnosis, and geographic area. Information on tumor morphology is also presented. In addition, the **CSR** features a chapter on adolescent and young adult cancers and a chapter on childhood cancers. Information on some rare cancers can be found in the summary tables of Section 1. For a detailed list of primary sites, the summary tables provide incidence and death rates for the most recent 5-year period, trends from 1975 to the most recent year, median age at diagnosis, median age at death, and survival rates.

Delay-adjusted cancer incidence rates are a distinctive feature of the **CSR**. Delay-adjustment corrects the current case count to account for underreporting and corrections to the data. The final delay-adjusted rates are valuable in more precisely estimating trends.

Changes in methodology to **CSR** include:

- Relative survival statistics using the SEER-18 registries are now calculated using lifetables based on socio-economic status and geography. A comparison of these statistics using the new and previous method is in Section 35: Survival Rate Comparison (https://seer.cancer.gov/csr/1975_2015/results_merged/sect_35_expected_rates.pdf).

The **CSR** files are provided in both PDF and HTML formats. The HTML format is provided as an alternative and accessible version of the *SEER Cancer Statistics Review*. The current edition of the **CSR** is available on the web at <https://seer.cancer.gov/csr/>. Statistics from SEER may also be obtained via SEER*Explorer (<https://seer.cancer.gov/explorer/>) **FastStats** (<https://seer.cancer.gov/faststats/>) or **Cancer Query Systems**

(<https://seer.cancer.gov/canques/>), which allow the user to access over 10,000,000 cancer statistics. The SEER Research Data file (<https://seer.cancer.gov/data/>) may be accessed by the public, either through **SEER*Stat** software or in an ASCII text format that can be analyzed with standard statistical software.

While most of the rates in this publication have been age-adjusted to the 2000 US standard population, some previous SEER publications have used the 1970 US standard million population. Therefore, rates given in this publication cannot be compared to rates given in those publications. This change conforms to a federal policy for reporting disease rates; it allows for the age-adjusted rate to more accurately reflect the current age distribution and burden of cancer.

INTERPRETATION OF CANCER STATISTICS

A number of factors may affect the interpretation of cancer incidence, mortality, and survival statistics provided in this report.

Survival rates for all cancers combined: The mix of cancers changes over time as the incidence of some cancers increases and the incidence of others decreases. The overall cancer survival rate can fluctuate even when the survival rates for site-specific cancers remain unchanged. (While it is possible to adjust the survival rate for all cancers combined on the basis of the relative frequencies of the component cancers, rates adjusted in this manner differ by only a small amount from unadjusted rates. In the future, such an adjustment may become more important if there are substantial changes in the incidence of various cancers.)

Early detection/screening: The improved earlier detection and diagnosis of cancers caused by new screening procedures may produce an *increase* in both incidence rates and survival rates. These increases can occur as a result of the introduction of a new procedure to screen subgroups of the population for a specific cancer; they need not be related to whether use of the screening test results in a decrease in mortality from that cancer. As the proportion of cancers detected at screening increases, presumably as a result of increased screening of the population, patient survival rates will *increase*, because they are based on survival time *after diagnosis*. The interval between the time a cancer is diagnosed by a screening procedure and the time when the cancer would have been diagnosed in the absence of screening is called **lead-time** (Zelen, 1976). (Screening for breast cancer has been demonstrated to result in increased survival over and above that resulting from lead-time alone and to reduce breast cancer mortality. The benefit of screening is being studied for some other cancers.)

If a new screening procedure consistently detects cancer in a *preinvasive* phase, it may result in a *decrease* in survival rates for *invasive* cancer. In this case, **length-biased sampling** (Zelen, 1976) may be operating. Length-biased sampling would result in the preferential detection—in a preinvasive phase—of those cancers that would have had a relatively good prognosis had they progressed to invasive disease; these potentially invasive cancers would be systematically

eliminated. If this occurs, the mix of cancers that are not detected at screening and then progress to invasive behavior may become less prognostically favorable, resulting in a *decrease* in survival rates for patients with invasive cancers. (Length-biased sampling may at least partially explain survival trends for cervical cancer. Other cancers possibly affected include breast, colon, rectum, and prostate.)

Changes in diagnostic criteria: Early detection of cancer resulting from either screening or earlier response to symptoms may result in the increasing diagnosis of small tumors that are not yet life-threatening. This may have the effect of raising the incidence rates and survival estimates without changing the mortality rates. Breast, colon, prostate, cervix uteri, bladder, and skin (melanoma) are the cancer sites most likely to be affected.

Technological advances in diagnostic procedures: In this report, trends in survival by stage at diagnosis for specific cancers are not presented; trends in stage distributions are presented rarely. However, it is possible to compare survival by stage.

The assignment of a given stage to a particular cancer may change over time due to advances in diagnostic technology. Introduction of new technology can give rise to a phenomenon known as **stage migration**. Stage migration occurs when diagnostic procedures change over time, resulting in an *increase* in the probability that a given cancer will be diagnosed in a *more advanced* stage. For example, certain distant metastases that would have been undetectable a few years ago can now be diagnosed by a computer tomography (CT) scan or by magnetic resonance imaging (MRI). Therefore, some patients who would have been diagnosed previously as having cancer in a *localized or regional* stage are now diagnosed as having cancer in a *distant* stage. The likely result would be to remove the worst survivors, those with previously undetected distant metastases, from the localized and regional categories and put them into the distant category. As a result, the stage-at-diagnosis distribution for a cancer may become less favorable over time, but the survival for each stage may improve: The early stage will *lose* cases that will survive *shorter* than those remaining in that category, while the advanced stage will *gain* cases that will survive *longer* than those already in that category. However, *overall survival would not change* (Feinstein et al., 1985). Stage migration is an important concept to understand when examining temporal trends in survival by stage at diagnosis as well as temporal trends in stage distributions; it could affect the analysis of virtually all solid tumors.

Evolution of stage classifications: Every few years, the American Joint Committee on Cancer produces a new cancer-staging manual; the seventh edition is the most recent (Edge et al., 2010). The evolution of such classifications reflects the identification of new prognostic factors that may influence choice of treatment. Historically, the SEER Program has only collected data on **extent of disease (EOD)**, rather than stage. EOD is *more specific* than stage and usually determines stage, even when stage definitions change. Thus, SEER easily adapts to changes in stage definitions; moreover, trends in a newly redefined stage can usually be calculated. Recently the SEER Program has begun collecting **Collaborative Stage**. Collaborative Stage has the advantage of being a consolidated data collection system of three

main staging systems (TNM, EOD, and Summary Stage) and allows combined pathological and clinical stage to be captured. New prognostic variables are introduced into staging for some cancers and so previously collected EOD data cannot determine new stage categories. There can be problems in assessing trends in stage of disease for these cancers. Only by reviewing the evolution of staging for a given cancer is it possible to determine what effects changes in stage definitions have had on stage-specific survival and on stage-at-diagnosis distributions. Stage migration (mentioned above) and EOD migration need also be taken into account. For some sites, the historic stage (*localized, regional, or distant*) is not shown, either because of inconsistencies in its definition over time or because stage is not appropriate (such as for leukemias, which are all considered to be distant at diagnosis).

Interpreting relative survival: The relative survival estimate is the ratio of observed survival to expected survival for a given patient cohort. Expected survival is based on mortality rates for the entire population, taking into account, as appropriate, the age, sex, race, and year of diagnosis of the patients. Assuming that the presence of cancer is the only factor that distinguishes the cancer patient cohort from the general population, relative survival estimates the probability that a patient will *not* die of the diagnosed cancer within the given time interval. This is the same as the probability that the patient will either survive the interval or die of a different cause.

A factor related to the risk of a cancer may also be related to the risk of dying from causes unrelated to the cancer. An example of such a factor is smoking. Smoking is a major risk factor for lung cancer; therefore, a cohort of lung cancer patients will contain a much higher proportion of smokers than the general population. However, smoking is also a risk factor for other diseases so smokers have a shorter life expectancy than nonsmokers. For this reason, expected survival estimates for lung cancer patients based on life tables for the general population will be unrealistically high; since relative survival = observed / expected, this will result in relative-survival estimates that are *lower* than they would be if the population consisted only of smokers. The problem cannot be easily corrected because separate life tables for smokers and nonsmokers are not available. Moreover, amount of smoking (usually measured in pack-years) is an important variable and cannot be easily quantified. In addition, expected survival may not be appropriate for patients with cancers of the cervix uteri or breast because the risk of these cancers has been associated with socioeconomic status (Baquet et al., 1991) which may be related to life expectancy. This should be considered when interpreting relative survival for these cancers.

Previous to the CSR for 1973–1996, the expected survival tables used were for 1970 and 1980; there were separate tables for whites, blacks, American Indians, Chinese, Japanese, Filipinos, white Hispanics, and Hawaiians. In updating the tables for 1990, several problems emerged. The US life tables are based on age, race, and sex information from death certificates. The information on race on the death certificate may not be accurate (Rosenberg et al., 1999). One reason is that funeral directors may inaccurately report race on a death certificate. Also, reported age at death, especially for those older than 85, may not be accurate because birth

certificates were not issued with as much regularity in the early 1900s as they are today. Although race misclassification and age-at-death misreporting exist across all races, they may be more problematic for races other than white or black because of those races' smaller population sizes. Therefore, life tables were generated for 1970, 1980, 1990, and 2000 only for white, black, and other; these life tables were used to produce the relative survival estimates in this review. There may be small variations among survival estimates calculated in this CSR and those in CSRs prior to 1973–1996.

Comparison with other databases: The SEER data are obtained from population-based cancer registries covering about 28 percent of the US population. It is sometimes of interest to compare cancer statistics for SEER areas with those from other registries both in the US and worldwide. In making such comparisons, one must carefully consider the factors mentioned above for both data sources. In addition, one should assess all of the following: (1) completeness of case ascertainment, (2) rules used to determine multiple primaries, (3) follow-up, (4) rules used in assigning and coding cause of death, and (5) the sources and procedures used in obtaining population estimates. Depending on the rates being compared, there could be other confounding factors which should be considered. The same standard or standard million population should be used for the age-adjustment of each group being compared; most statistics from outside the US are based on the 2000 world standard million population. Examples of other databases are US Cancer Statistics (<https://nccd.cdc.gov/uscs>) and CINA+ Online (<https://www.cancer-rates.info/naaccr/>).

It is sometimes of interest to compare survival for cancer patients in SEER areas with data from clinical trials. *This must be done with great caution.* Survival data from clinical trials may have been obtained from a patient population that differs from that of SEER patients in prognostic factors for the given cancer; any survival comparisons would have to adjust for such differences. Also, it is necessary to verify that the methodology used in computing survival is the same for both data sources. Furthermore, patients on clinical trials may differ from SEER patients in characteristics that may be related to survival but are not recorded in either database. If this were true for a given cancer, it would not be possible to make valid comparisons of this type.

Errors in data collection: In the process of registering cancer patients, errors may be made in abstracting and coding the data, which include demographic information, cancer site, histology, extent of disease, treatment, and patient survival. Quality control studies are periodically carried out to detect and correct this type of error, but no attempt is made to incorporate this source of error into the variance estimates of cancer rates reported here.

Comparison of this report with previous reports: The cancer registries that participate in the SEER Program submit data on all cancers diagnosed in their coverage areas to the NCI each year. Because of the dynamic nature of the registries' databases, *the reported number of new cancer cases in a particular race, sex, age, cancer category in a given calendar year may change from that which has been reported in a previous publication.* For a given diagnosis year, additional cancer cases that were previously overlooked may have been found

and reported to the central registry. There may have been follow-back of cancers diagnosed by death certificate only; successful efforts to establish the dates of diagnosis for such patients will change the number of patients reported for a given diagnosis year. Code changes may occur when a patient dies; for example, information on race is generally available on the death certificate and may be used to update a previously unknown value. There may have been elimination of duplicate records for the same patient, often due to name changes or misspellings.

Thus, a recent report may have a different number of cases for a given diagnosis year than an earlier report, with resulting effects on incidence and possibly survival. Population estimates may also change from one report to another for some calendar years. This occurs because the NCI receives population estimates that are regularly revised and updated by the Bureau of the Census (**BOC**). Such changes may result in some differences between incidence and mortality rates for a given calendar period as published in different reports. See our website for the most current information about the population estimates (<https://seer.cancer.gov/popdata/>).

REFERENCES

Baquet CR, Horm JW, Gibbs T, Greenwald P. Socioeconomic factors and cancer incidence among blacks and whites. *J Natl Cancer Inst* 1991; 83:551-557.

Breslow L (Chairman, Extramural Committee to Assess Measures of Progress Against Cancer). Measurement of progress against cancer: Final report to the Senate Appropriations Committee. Bethesda: National Cancer Institute; 1988.

Edge SB, Byrd DR, Compton CC, Fritz AG, Greene FL, Trotti A, editors. *AJCC Cancer Staging Manual*, 7th ed. New York (NY): Springer; 2010.

Feinstein AR, Sosin DM, Wells CK. The Will Rogers phenomenon: Stage migration and new diagnostic techniques as a source of misleading statistics for survival of cancer. *New Engl J Med* 1985;312:1604-1608.

Lewis DR, Check DP, Caporaso NE, Travis WD, Devesa SS. U.S. Lung Cancer Trends by Histologic Type. *Cancer* 2014; 120(18):2883-92.

Rosenberg HM, Maurer JD, Sorlie PD, Johnson NJ, MacDorman MF, Hoyert DL, Spitler JF, Scott C. Quality of death rates by race and Hispanic origin: A summary of current research. Hyattsville (MD): National Center for Health Statistics; Vital and Health Statistics, Series 2, No. 128, 1999.

Zelen M. Theory of early detection of breast cancer in the general population. In: Heuson J-C, Matthei WH, Rozenzweig M, editors. *Breast Cancer: Trends in Research and Treatment*. New York (NY): Raven Press; 1976. p. 287-299.

Zhang S, Luo J, Zhu L, Stinchcomb D, Campbell D, Carter G, Gilkeson S, Feuer EJ. (2014), Confidence intervals for ranks of age-adjusted rates across states or counties. *Statistics in Medicine*. doi: 10.1002/sim.6071

TECHNICAL NOTES

There are four measures commonly used to assess the impact of a cancer in the general population and are reported in this review. The **incidence rate** is the number of new cases per year per 100,000 persons. The **death (or mortality) rate** is the number of deaths per year per 100,000 persons. The survival estimate is the proportion of patients alive at some point subsequent to the diagnosis of their cancer. The **prevalence count** is the number of people alive that have ever been diagnosed with a cancer. The Surveillance, Epidemiology, and End Results (**SEER**) Program (<https://seer.cancer.gov/>) (based within the Surveillance Research Program (**SRP**) at the National Cancer Institute (**NCI**) collects incidence and survival data for all areas that participate in the Program. The National Center for Health Statistics (**NCHS**) provides mortality data for the entire United States (**US**). All incidence and mortality rates in this report are age-adjusted (see below) to the 2000 US standard population (see Appendix) unless otherwise specified. Age-adjustment minimizes the effect of a difference in age distributions when comparing rates.

THE SEER PROGRAM

The National Cancer Act of 1971 mandated the collection, analysis, and dissemination of data useful in the prevention, diagnosis, and treatment of cancer. This mandate led to the establishment of the SEER Program. The population-based cancer registries participating in NCI's SEER Program routinely collect data on all cancers occurring in residents of the participating areas. Trends in cancer incidence and patient survival in the US are derived from this database. See the SEER Research Data (<https://seer.cancer.gov/data/>) for more information.

The SEER Program is a sequel to two earlier NCI programs—the End Results Program and the Third National Cancer Survey. The initial SEER reporting areas were the States of **Connecticut, Iowa, New Mexico, Utah, and Hawaii**; the metropolitan areas of **Detroit, Michigan, and San Francisco-Oakland, California**; and the Commonwealth of Puerto Rico. Case ascertainment began with January 1, 1973, diagnoses.

In 1974-1975, the program was expanded to include the metropolitan area of New Orleans, Louisiana, the thirteen-county **Seattle-Puget Sound** area in the State of Washington, and the metropolitan area of **Atlanta, Georgia**. New Orleans participated in the program only through the 1977 data collection year. In 1978, ten predominantly African-American counties in **rural Georgia** were added. **American Indian residents of Arizona** were added in 1980. In 1983, four counties in New Jersey were added with coverage retrospective to 1979. New Jersey and Puerto Rico participated in the program until the end of the 1989 reporting year. The National Cancer Institute also began funding a cancer registry that, with technical assistance from SEER, collects information on cancer cases among **Alaska Native** populations residing in Alaska. In 1992, the SEER Program was expanded to increase coverage of minority populations,

especially Hispanics, by adding **Los Angeles County** and four counties in the **San Jose-Monterey** area south of San Francisco. In 2001, the SEER Program expanded coverage to include **Kentucky, Greater California** (the counties of California that were not already covered by SEER), **New Jersey**, and **Louisiana**. In 2012, **Greater Georgia** (the parts of Georgia not included in Atlanta and Rural Georgia) was added to the SEER Program, with data retroactive to 2000.

The long-term incidence trends and survival data for this report are from five states (Connecticut, Hawaii, Iowa, New Mexico, and Utah) and four metropolitan areas (Detroit, Atlanta, San Francisco-Oakland, and Seattle-Puget Sound) (Fig. I-1); this set of registries is called the **SEER 9**. Additional tables show more recent incidence trends for the **SEER 13** areas (the 9 areas above plus Los Angeles, San Jose-Monterey, Alaska Native Registry, and rural Georgia) since 1992 and additional information on race and ethnicity. Other tables give statistics for the **SEER 18** areas; these are the SEER 13 plus Kentucky, Greater California, New Jersey, Louisiana, and Greater Georgia.

The participating regions were selected principally for their ability to operate and maintain a population-based cancer reporting system and for their epidemiologically significant population subgroups. With respect to selected demographic and epidemiologic factors, they are when combined a reasonably representative subset of the US population. Data from the 9, 13, or 18 SEER geographic areas are used in this report; the given groups contain, respectively, approximately 9, 14, or 28 percent of the US population. By the end of the 2012 diagnosis year, the database of the 18 SEER registries (plus Arizona Indians) contained information on over 7 million cases diagnosed since 1973. New cases added in the most recent data year numbered over 449,000.

The goals of the SEER Program are:

- 1) to assemble and report, on a periodic basis, estimates of cancer incidence, mortality, survival, and prevalence in the US;
- 2) to monitor annual cancer incidence trends to identify unusual changes in specific forms of cancer occurring in population subgroups defined by geographic and demographic characteristics;
- 3) to provide continuing information on trends over time in the extent of disease at diagnosis, trends in therapy, and associated changes in patient survival; and
- 4) to promote studies designed to identify factors amenable to cancer control interventions, such as: (a) environmental, occupational, socioeconomic, dietary, and health-related exposures; (b) screening practices, early detection and treatment; and (c) determinants of the length and quality of patient survival.

DATA SOURCES

INCIDENCE AND SURVIVAL DATA

The SEER Program contracts with nonprofit, medically-oriented organizations having statutory responsibility for registering diagnoses of cancer among residents of their respective geographic coverage areas. Each SEER contractor:

- 1) maintains a cancer information reporting system;
- 2) abstracts records for *resident* cancer patients seen in every hospital both inside and outside the coverage area;
- 3) abstracts all death certificates of *residents* (dying both inside and outside the coverage area) on which cancer is listed as a cause of death;
- 4) strives for complete ascertainment of cases by searching records of private laboratories, radiotherapy units, nursing homes, and other health services units that provide diagnostic service;
- 5) registers all in situ and malignant neoplasms (with the exceptions of certain histologies for cancer of the skin and—beginning in 1996—in situ neoplasms of the cervix uteri);
- 6) records data on all newly diagnosed cancers, including selected patient demographics, primary site, morphology, diagnostic confirmation, extent of disease, and first course of cancer-directed therapy;
- 7) provides active follow-up on all living patients (except for those with in situ cancer of the cervix uteri);
- 8) maintains confidentiality of patient records;
- 9) at least annually submits electronically to NCI data on all reportable diagnoses of cancer made in residents of the coverage area.

For 1992 to 2000 diagnoses, the SEER program codes site and histology by the *International Classification of Diseases for Oncology*, second edition (**ICD-O-2**) (Percy et al., 1990). All cases before 1992 were machine-converted to ICD-O-2. Cases diagnosed 2001-2009 have been coded according to the third edition (**ICD-O-3**) (Fritz et al., 2000). Starting with patients diagnosed in 2007, the new multiple primary and histology coding rules may impact their incidence data for some cancer sites (e.g., female breast). However, the impact of the new rule on observed incidence is negligible for a majority of the cancer sites. To learn more about the multiple primary rules, visit: <https://seer.cancer.gov/tools/mphrules/>. Beginning with 2010 diagnoses, cases are coded based on ICD-O-3 updated for hematopoietic codes based on *WHO Classification of Tumours of Haematopoietic and Lymphoid Tissues* (2008). The primary site groupings used for incidence are found in the Appendix. Changes were made to the site recode for ICD-O-2 for comparability with cases coded to ICD-O-3. Follow-up rates are also in the Appendix.

MORTALITY DATA

The SEER Program annually obtains from the National Center for Health Statistics (NCHS) a file containing information on all deaths occurring in the US by calendar year. Information on each death includes age at death, sex, geographic area of residence, and underlying and contributing causes of death. For this publication, only the underlying cause of death is used in the calculation of death rates. Cause of death for 1969-1978 was coded according to ICD-8; for 1979-1998, ICD-9 was used; beginning with deaths in 1999, ICD-10 was used. Mortality rates for the SEER geographic areas, for each state, and for the entire US are obtained from these data. A list of the mortality site groupings used in this publication is in the Appendix and reflects updates made in 2004.

POPULATION DATA

The population estimates used in the SEER*Stat software to calculate cancer incidence and mortality rates for this report are a modified version of the intercensal and Vintage 2015 annual time series of July 1 county population estimates by age, sex, race, and Hispanic origin that are produced by the Population Estimates Program of the US Census Bureau (<https://www.census.gov/programs-surveys/popest.html>) with support from the NCI through an interagency agreement. Descriptions of the methodologies employed by the Census Bureau for various sets of estimates may be found on the same website. Vintage 2015 population estimates were used; these estimates were developed from the actual 2010 census results.

County population estimates for 2000 and later years must be bridged from 31 race categories used in Census 2000 to the four race categories specified under the 1997 OMB standards in order to report long-term cancer trends. The bridging methodology was developed by the National Center for Health Statistics and is described in a report (Ingram et al., 2003) and on their website http://www.cdc.gov/nchs/nvss/bridged_race.htm

Modifications made by the NCI to the population estimates are documented in "Population Estimates Used in NCI's SEER*Stat Software" (<https://seer.cancer.gov/popdata/methods.html>) and the population data files are available for download (see "Download US Population Data" from <https://seer.cancer.gov/popdata/download.html>). Several of the modifications pertaining to the grouping of specific counties needed to assure the compatibility of all incidence, mortality and population datasets. Another modification affects only population estimates for the State of Hawaii. The Epidemiology Program of the Hawaii Cancer Research Center has developed its own set of population estimates, based on sample survey data collected by the Hawaii Department of Health. This effort grew out of a concern that the native Hawaiian population has been vastly undercounted in previous censuses. The "Hawaii adjustment" to the Census Bureau's estimates has the net result of reducing the estimated white population and increasing the estimated Asian and Pacific Islander population for the state. The estimates for the total population, black population, and American Indian and Alaska Native populations in Hawaii are not modified.

The cancer incidence and mortality rates for American Indians and Alaska Natives (AI/AN) are based on the geographic areas (counties) included in the Indian Health Service's Contract Health Service Delivery Area (CHSDA). This reflects a concern that previously reported AI/AN rates were underestimated due to racial/ethnic misclassification of American Indian cases in geographic areas outside of CHSDA. This change has the net effect of higher, and more accurate, incidence and mortality rates for this population. Beginning in 2013, CSR reporting diagnoses 1975-2010, CHSDA counties were updated with 9 new counties designated as CHSDA. Four of these are in SEER areas. This addition was made to better reflect AI/AN populations that had been living in these counties.

Usually the use of a population estimate for July 1 of a particular year reflects the average population of that area for the year. Both Hurricane Katrina and Hurricane Rita struck the Gulf Coast area of the United States in 2005. This had the effect of displacing large populations. Since there weren't any population estimates by age, race, sex, and county for time periods just after the hurricanes, it is very difficult to estimate the actual population at risk for certain areas along the Gulf Coast for 2005. For Louisiana, only the first six months of incidence data for 2005 coupled with ½ of the population estimate for July 1, 2005, were used to calculate cancer incidence. For death rate calculations, no adjustments were made to the total US population, but for the Gulf area, an adjustment for displaced populations was made for 2005 state rates. For more details, see <https://seer.cancer.gov/popdata/methods.html>.

2000 US STANDARD POPULATION

Starting with the November 2004 SEER submission of data (diagnoses through 2002), the SEER Program age-adjusts using the 2000 US standard population based on single years of age from the Census P25-1130 series estimates of the 2000 US population (Day, 1996). For the CSR, 19 age groupings were used for age-adjustment: <1, 1-4, 5-9, ... , 80-84, 85+.

STATISTICAL METHODS

ESTIMATED CANCER CASES AND DEATHS IN 2018

The American Cancer Society (**ACS**) projects the numbers of new cancer cases and cancer deaths in the US in 2018 (Cancer Facts & Figures – 2018, American Cancer Society). The ACS projects incidence in 2018 based on incidence rates for 2000-2014 from 50 states and the District of Columbia, representing about 98% of the US population. These high-quality incidence data were submitted to the North American Association of Central Cancer Registries (NAACCR) by 50 states (and District of Columbia) belonging to the SEER Program and/or the National Program of Cancer Registries (NPCR).

LONG-TERM TRENDS, 1950-2015

Trends in cancer mortality from 1950 to 2015 are summarized by age both for all cancers combined and for lung cancer (Table 1-2). These cancer mortality trends are based on the mortality experience in the entire US. Summaries of long-term trends back to 1950 in cancer survival are also shown for whites. Use caution when interpreting these statistics. Evaluating trends over a long period of time may hide recent changes in the trends.

YEARS OF LIFE LOST DUE TO PREMATURE DEATH FROM VARIOUS CAUSES

Death rates alone give an incomplete picture of the burden that deaths impose on the population. Another measure is the years of life lost due to premature death. This shows the extent to which life is cut short by a particular cause or disease.

This measure is estimated by linking life table data to each death of a person of a given age and sex. The life table permits a determination of the number of additional years an average person of that age, race, and sex would be expected to live. In this report, the age groups used in the calculation were 1-year intervals. These remaining years of life left are summed over all deaths due to a particular cause, yielding the estimate of the number of person-years of life lost (**PYLL**). The average years of life lost (**AYLL**) is obtained by dividing the PYLL by the number of deaths. Both of these measures can be calculated for any cause of death.

RELATIVE SURVIVAL

Relative survival (Ederer, 1961) was developed to provide an objective measure of the probability of survival of cancer in the absence of other causes of death. It is a measure that is not influenced by changes in mortality from other causes and, therefore, provides a useful measure for both tracking survival across time and comparisons between racial/ethnic groups or between registries. Cause-of-death information obtained from death certificates can be unreliable due to misclassification error (e.g. the site of recurrence being classified as the cause of death). Therefore, instead of calculating the probability of surviving cancer in the usual (cause-specific) way, considering deaths from other causes as censoring events, relative survival compares the observed survival proportion of a group of cancer patients with the survival of a “similar” theoretical cancer-free group. Relative survival is formally defined as the ratio of the observed survival (all causes of death) of a cohort of cancer patients to the expected survival of a comparable set of cancer-free individuals. Since a cohort of cancer-free individuals is difficult to obtain, life tables representing survival of the general population are used instead. The underlying assumption is that the cancer deaths are a negligible proportion of all deaths. To learn more on this topic, visit: <https://surveillance.cancer.gov/survival/measures.html>.

Expected survival can be calculated using different methods which vary with respect to the definition of the matching group. The most common methods are: Ederer I (Ederer, et al., 1961), Ederer II (Ederer and Heise, 1959) and Hakulinen (Hakulinen, 1982) and Pohar-Perme. Since

2012, we use the Ederer II (instead of Ederer I used previously) method to estimate the expected rate in SEER*Stat and the CSR. This method has shown to be a less biased estimate of net survival. For more detail regarding this topic, read Cho et al., 2012 at: <https://surveillance.cancer.gov/reports/>. As of 2013, Survival time was calculated using pre-calculated months based on the exact day information. See <https://seer.cancer.gov/survivaltime/>. As of 2014, the default censoring age for survival calculations has changed from 199 to 99 year when using newly available expected survival tables. Minimal changes may occur in survival for older age groups. See <https://seer.cancer.gov/expsurvival/> for more information. We use national life tables by age, calendar year and race (whites, black, other races) to estimate expected survival. Other races include both Asian or Pacific Islanders (API), and lowest for American Indians and Alaskan Natives (AIAN). In 2017 we constructed state and race specific life tables by county of residence socio-economic status from 1992 forward. As of 2018 these life tables will be used as the default to estimate expected survival for that only include cases diagnosed after 1992 (for example SEER (2000+)).

The state/race/SES life table were constructed using counts of deaths and populations by county, single year age at death (30 to 84 years), race/ethnicity, sex, and calendar year 1992-2013. We used mutually exclusive race/ethnicity groups: Non-Hispanic (NH) White, NH Black, NH AIAN, NH API, and Hispanics (hereafter we exclude the NH prefix when referencing race/ethnicity). Hispanic ethnicity includes all race categories. Because of misclassification errors of AIAN race in death certificates, we restricted the AIAN data to mortality rates from Contract Health Service Delivery Area (CHSDA) counties. We fit Poisson regression models to the log of mortality rates to estimate the life tables separately for men and women and each race/ethnicity. Age and calendar year were modeled as spline functions to capture non-linear effects. The models varied by geographic area (state, region, and national) and the inclusion or not of the SES index as a covariate depending on sufficient numbers of deaths and population counts for each race-ethnicity. For more details on the methods and data to estimate life tables a technical is available on request.

CAUSE-SPECIFIC SURVIVAL

Cause-specific survival is a net-survival measure representing survival of a specified cause of death in the (theoretical) absence of other causes of death. Estimates are calculated by specifying the cause of death. Individuals who die of causes other than the specified cause are censored. This requires a cause-of-death variable that accurately captures all causes related to the specific cause. Cancer registries use algorithms to process causes of death from death certificates in order to identify a single, disease-specific, underlying cause of death. In some cases, attribution of a single cause of death may be difficult and misattribution may occur. For example, a death may be attributed to the site of metastasis instead of the primary site (Percy et al., 1981).

To capture deaths related to the specific cancer but not coded as such, the SEER cause-specific death classification variable is defined by taking into account causes of deaths in

conjunction with tumor sequence (i.e., only one tumor or the first of subsequent tumors), site of the original cancer diagnosis, and comorbidities (e.g., AIDS and/or site-related diseases). To learn more on this topic, please read the recent article published at the Journal of National Cancer Institute (Howlader et al., 2010) or visit: <https://seer.cancer.gov/causespecific/>.

CANCER PREVALENCE

Methods: In this report prevalence is calculated at 1/1/2014. Limited-duration prevalence is calculated using the counting method implemented in the SEER*Stat software. This method calculates the number or proportion of people alive at the prevalence date who had a diagnosis of the disease within the past x years (e.g., $x = 5, 10, 20$, or the full history of the registry). Because SEER has available information for the various racial/ethnic groups for different numbers of years, different years and registries were used to estimate limited-duration prevalence. Prevalence estimates for all races combined, for whites, and for blacks use cases from 1975 through 2015 from the SEER 9 registries; prevalence estimates for Asian Pacific Islanders and Hispanics use cases diagnosed from 1990 through 2015 from the SEER 11 areas and rural Georgia.

The limited-duration prevalence method includes a correction for people lost to follow-up. For each individual lost to follow-up, a probability of being alive at the prevalence date is estimated from an appropriate survival function stratified by age at diagnosis (0–59, 60–69, 70+), sex, cancer site, year of diagnosis, and race, conditional on being alive at the time of loss to follow-up. Year of diagnosis is stratified into 5-year groups from the prevalence date, with the least recent interval being of varying length (4–8 years), depending on the length of years used to calculate prevalence. Race is stratified into white, black, other (American Indian/Alaska Native, Asian/Pacific Islander), and unknown/other-unspecified. When we use the SEER 11 registries, the same stratification as before is used, with American Indian/Alaska Native separated from Asian/Pacific Islander. Prevalence calculations for Hispanics use race stratified into: white, non-white, and unknown.

Different methods can be used to determine which tumors are to be included for people diagnosed with multiple tumors. Unless otherwise specified, prevalence calculations include only the *first malignant tumor per person*; that is, in situ cancers and second-or-later primary cancers were not included. Thus, if a woman had a melanoma prior to a breast cancer diagnosis, her melanoma would contribute to the prevalence of melanoma and to the prevalence of all sites, but the breast cancer would not contribute to the prevalence of breast cancer. Counting only one cancer per individual avoids some ambiguity in prevalence counts, and allows the counts for individual sites to sum to the all sites total. Table 1.22 in the Overview Chapter compares 5-Year Limited Duration Prevalence using different selection criteria: A) 1st Invasive Tumor Ever, B) 1st Per Site in Previous 38 Years and C) 1st Per Site in Previous 5 years. A female breast cancer to be included in the 5-Year Limited Duration Prevalence needs to be diagnosed in in the 5 years prior to the prevalence date and (A) be the first tumor ever of

the woman; (B) the first breast cancer of the women in the prior 38 years, the women could not have had other breast cancers between 6 and 38 years prior to the prevalence date, and (C) be the first breast cancer in the prior 5 years, i.e., the women could have had other breast cancer 6 or more years prior to the prevalence date, and if she had 2 breast cancers between 2008 and 2012 only the first can be counted. For more information on tumor selection criteria refer to <http://surveillance.cancer.gov/prevalence/methods.html>.

Complete prevalence is an estimate of the number of persons (or the proportion of population) alive on a specified date who had been diagnosed with the given cancer, no matter how long ago that diagnosis was. It was estimated for all races, whites, and blacks by applying the *completeness index method* (Capocaccia & De Angelis, 1997; Merrill et al., 2000; Mariotto et al., 2002) to limited-duration prevalence. The completeness index method is implemented in the COMPREV software, which can be found at <https://surveillance.cancer.gov/comprev/>. Validation of the completeness index for all races and for whites was made by using data from the Connecticut Tumor Registry (CTR) beginning with 1940. For blacks, SEER 9 data beginning with 1975 were used; identification of blacks is not possible in the CTR data prior to 1970. To validate the completeness index for blacks, we have compared the performance of the method to obtain 24-year prevalence from 10-year limited-duration prevalence. For all races combined and for whites, in cases where the validation indicated some lack of fit of the model, an approximation to the completeness index was derived from the CTR data. If there was a lack of fit for blacks, no estimate of complete prevalence was reported. Complete prevalence for Asian/Pacific Islanders and Hispanics is not available at this time. Complete prevalence by age for all races combined was validated by comparing estimated 10-year complete prevalence with observed prevalence from the CTR data. Prevalence by age is reported for the sites that validated well.

The US cancer prevalence counts at 1/1/2014 *were estimated* by multiplying the SEER age- and race-specific prevalence proportions by the corresponding US population estimates based on the average of 2013 and 2014 population estimates from the US Census Bureau. US cancer prevalence counts for all races were estimated by summing the US estimated counts for whites/unknown, blacks, and other races. For Hispanics, the estimates for Hispanics of white or unknown race and for Hispanics of other races were summed.

Complete prevalence estimates of the number of individuals in the US diagnosed with cancer as children (ages 0-19), including those surviving for more than 38 years, is calculated using a statistical method that estimates the number of childhood survivors diagnosed before 1975 (Simonetti et al., 2008; Mariotto et al., 2009). Limited-duration prevalence proportions by age at prevalence are not shown for childhood cancers (age at diagnosis 0-19) since many of these estimates are not informative. For example, the number of people diagnosed with childhood cancers in the last 25 years and who are currently age 50-59 is zero by definition. For more details on available prevalence estimates, see <https://surveillance.cancer.gov/prevalence/>.

PROBABILITY OF BEING DIAGNOSED WITH OR DYING FROM CANCER

Lifetime and interval risks of being diagnosed with cancer: The probability of being diagnosed with cancer is computed by applying cross-sectional age-specific 2012-2014 incidence rates from the SEER 18 areas and death rates from those same areas to a hypothetical cohort of 10,000,000 live births. This cohort is considered to be at risk for two mutually exclusive events: (1) developing the specified cancer, and (2) dying of other causes without developing the specified cancer. Using these two types of events, a standard **multiple decrement life table** (with 20 age groups from 0-4 to 90-94 and 95+) is derived. For each age interval, the number alive and free of the specified cancer at the beginning of the interval is decremented by the number who develop the specified cancer and the number who die of other causes. The lifetime risk of being diagnosed with the specified cancer is derived by summing all cancer cases from age 0-4 through age 95+ and dividing by 10,000,000. This calculation does not assume that an individual lives to any particular age; rather, it is the sum over all age intervals of the probability of living to the beginning of that interval without developing the given cancer times the probability of developing the cancer in that interval. The probability of developing cancer during any time period (e.g., between age 50 and age 60) is calculated by adding up all the cancers in the life table over the specified age range and dividing by the number of individuals alive and free of the specified cancer at the beginning of the period. The methodology is described in detail in (Fay et al., 2003) and (Fay, 2004). To improve the precision of the calculations, rates were calculated beyond the usual last open ended age interval (i.e. 85+) for the age groups 85-89, 90-94, and 95+.

Lifetime risk of dying from cancer: The lifetime risk of dying from a specified cancer is derived using a standard multiple decrement life table (Elandt-Johnson & Johnson, 1980). For each age, the risks of dying of the specified cancer and of all other causes are calculated, based on mortality data from the entire United States.

Detailed methodology and software: The estimates of developing and dying from cancer are implemented in DevCan (Probability of DEveloping or dying from CANcer software). More details on the software, various databases, and the methodology can be found at <https://surveillance.cancer.gov/devcan/>.

US CANCER DEATH RATES BY STATE

Each cancer-site-specific section presents the death rate for the given cancer for each state and the District of Columbia, specifying the five highest and the five lowest death rates by state for the most recent 5-year period for all persons, males only, and females only. The rates are per 100,000 persons; they are age-adjusted to the 2000 US standard population. (In some previous editions of the CSR, the 1970 US standard million population was used; *death rates standardized to the 2000 US standard million population cannot be compared to death rates standardized to the 1970 US standard million population.*)

The **percent difference (PD)** between a state rate and the rate for the total US is given by the formula:

$$PD = [(State\ Rate - Total\ US\ Rate)/Total\ US\ Rate] * 100$$

The **standard error** for each age-adjusted state death rate is calculated, based on the assumptions that (1) for each age-specific rate, the number of deaths is a Poisson random variable (Keyfitz, 1966) and (2) the variance of the age-adjusted rate is a linear combination of the variances of the age-specific rates (Snedecor & Cochran, 1980; pp. 188-9).

The **standard error of the difference (SE_d)** between a state rate and the total US rate is given by the formula

$$SE_d = \text{Square Root of } [SE_s^2 + SE_U^2 - 2 * Cov_{s,U}]$$

where SE_s and SE_U are the standard errors of a state rate and of the total US rate, respectively, and Cov_{s,U} is the covariance between the two rates. The variance of each rate (i.e., the square of the standard error) and the covariance between the two rates are based on the Poisson assumption. The standard error does not represent the total error that may be present in the age-adjusted rate; it is merely the square root of the variance associated with the rates. In addition to this variance, there also exist potential biases and errors in the measurement of the rate that are difficult to assess accurately and probably impact differently on the error calculations for different states.

The difference between each age-adjusted state rate and the age-adjusted US rate is tested for statistical significance (see below) by calculating a **Z** (standard normal) statistic from the formula:

$$Z = (State\ rate - Total\ US\ rate) / SE_d$$

Although the rates being compared are not independent because each state is part of the US, the statistical test may not be substantially affected if the state represents a small proportion of the total US. There is also an adjustment for multiple comparisons; see below under *Statistical Significance*.

The states are ranked according to the death rate, with 1 indicating the highest and 51 the lowest rate in the US. 95% confidence intervals for the rank are shown in parentheses () after the rank. The confidence intervals of ranks of age-adjusted rates are calculated using a simulation-based method (Zhang, 2014) implemented in the CI*Rank tool <https://surveillance.cancer.gov/cirank/>.

JOINPOINT REGRESSION ANALYSIS OF CANCER TRENDS

Joinpoint regression is a useful way to characterize trends in cancer rates and other health indices (Kim et al., 2000). It characterizes segments using connected linear segments on a log scale (i.e. constant annual percent changes (APC's) between changepoints. The locations of the changepoints are optimally determined using by the data using a statistical algorithm. To achieve greater descriptive accuracy, a statistical algorithm finds the optimal number and location of places where a trend changes. The point (in time) when a trend changes is called a **joinpoint**. Trends may change in different ways at a joinpoint: from up to down, from down to up, from up to up at a different rate, or from down to down at a different rate. A **joinpoint regression model** describes the trends by a continuous, piecewise-exponential function. Adjacent segments are connected at a joinpoint. The segments are connected because we assume that rates generally change smoothly, rather than “jump” abruptly. In each segment, the rates are assumed to grow or decay exponentially ($y = e^{mx+b}$), i.e., to change by a constant percentage each year. Thus the “slope” m in each segment can be associated with a fixed annual percent change (**APC**) by $APC = 100(e^m - 1)$.

Joinpoint analysis first assumes no joinpoints are needed to describe the data accurately, i.e., the trend over the entire interval 1975-2014 does not change. Joinpoints are added in turn if they are statistically significant. Thus, in the final model, each joinpoint represents a significant change in trend. Smoother polynomial models may provide a good fit overall, but are less sensitive to what is occurring at the ends of the data.

In running the Joinpoint program, we set the program parameters as follows:

- (1) Joinpoints occur only at exact years; the joinpoint is not necessarily the same as the data point for that year;
- (2) The minimum time interval between consecutive joinpoints is three years;
- (3) The first joinpoint is not earlier than two years after the first year of data;
- (4) The last joinpoint is not later than two years before the last year of data;
- (5) The maximum number of joinpoints is five for 1975-2013 (SEER 9) data and three for 1992-2014 (SEER 13) data.

These restrictions provide some added stability to the resultant models. Different values for these parameters may yield a different joinpoint model. Since the test statistic to determine if additional joinpoints are necessary cannot be compared against any known standard distribution to determine significance (e.g., the normal, t, or f), a permutation test is used which simulates the distribution of the test statistic under the null hypothesis. Thus an element of randomness is introduced by the random number stream used. However, for greater consistency in the p-values obtained if one were to change the random seed for each run, we run the program for 4499 permutations.

A Windows-based program, *Joinpoint*, is freely available at <https://surveillance.cancer.gov/joinpoint/>; it accepts data from the *SEER*Stat* program, as well as user-defined data. Further details on joinpoint regression may be found at the website. Starting with the 2012 edition of CSR, we have generated all our cancer trend statistics using a Linux-based *Joinpoint* program as opposed to the downloadable Windows-based program. As a result of using a different platform, in rare instances the results (e.g., # of joinpoints) may differ.

Average Annual Percent Change (AAPC) is a summary measure of a trend over a pre-specified fixed interval based on an underlying joinpoint model. It allows us to use a single number to describe the average trend over a period of multiple years. It can be estimated even if the joinpoint model indicates that there were changes in trends during those years, since it is estimated as a geometric weighted average of the joinpoint APCs, with the weights equal to the lengths of each segment over the pre-specified fixed interval. In this report, we have included AAPCs as an addendum to the underlying joinpoint trends, and as a summary measure to compare fixed interval trends by race/ethnicity. For more information on how the AAPC is calculated and the advantages of reporting an AAPC over APCs, see <https://surveillance.cancer.gov/help/joinpoint/setting-parameters/advanced-tab/average-annual-percent-change-aapc>.

Jump Model/Comparability Ratio Model

The Jump Model / Comparability Ratio Model in the Joinpoint software provides a direct estimation of trend data (e.g. cancer rates) where there is a coding, which causes a “jump” in the rates, but is assumed not to affect the underlying trend. To account for ICD-9 to ICD-10 coding change, occurred in 1998, alternative trends estimated from Jump model and Comparability Ratio Model are obtained for Melanoma. Those trends and more information can be found in <https://surveillance.cancer.gov/joinpoint/jump.html> .

REPORTING DELAY

Timely and accurate calculation of cancer incidence rates is hampered by **reporting delay**, the time lapse before a diagnosed cancer case is reported to the NCI or the delay in receiving updated information for an existing case. Currently, NCI allows a standard delay of 22 months between the end of the diagnosis year and the time the cancers are reported to the NCI in November, almost two years later. The data are released to the public in the spring of the following year. For example, cases diagnosed in 2014 were first reported to the NCI in November 2016 and released to the public in April 2017. However, in each subsequent release of the SEER data, *records from all prior diagnosis years* (e.g., diagnosis years 2014 and earlier in the 2016 submission to the NCI) *are updated* as either new cases are found or new information is received about previously submitted cases.

The submissions for the most recent diagnosis year are, in general, about two percent below the total number of cancers that will eventually be submitted for that year, although this varies by cancer site and other factors. To adjust for this, statistical models have been developed to

estimate “reporting delay-adjusted rates” for the SEER 9 since 2003 and SEER 13 registries since 2010 and the delay adjusted rates are reported.

The idea behind modeling reporting delay is *to adjust the recent rates to anticipate future corrections (additions, changes, and deletions) to the data*. These adjusted rates and the associated delay model are valuable in more precisely determining current cancer trends, as well as in monitoring the timeliness of data collection—an important aspect of quality control (Clegg et al., 2002).

In addition to registries funded by NCI-SEER, registries for the remainder of the U.S. are funded by the Centers for Disease Control and Prevention National Program of Cancer Registries ([CDC-NPCR](#)). (Some registries are co-funded by both NCI and CDC). Annual cancer incidence and survival data are reported by U.S. registries to NCI-SEER and CDC-NPCR, while registries throughout the US and Canada are report annually to the North American Association of Central Cancer Registries (NAACCR), a registry member organization. A coordinated effort by NCI, CDC and NAACCR has led to a unified approach to estimate and report delay adjusted rates.

Starting with data released in 2015, for the first time, delay adjustment factors is produced based on December 2014 data submitted to the NAACCR. The delay adjusted rates are then estimated from the delay adjustment factors by cancer site, registry, age group, gender, race, and year of diagnosis and linked to the appropriate cases (based on cancer site, registry, age group, gender, race, and year of diagnosis), to data submissions for each of the three partners in this joint effort (NCI-SEER, NAACCR, and CDC-NPCR). Starting from 2017 release, delay adjustment factors for Ethnicity (Hispanic and Non-Hispanic) and Race x Ethnicity combination are also estimated. This will allow all the partners and users of these data to produce delay adjusted rates. See Appendix for details.

In this report, we show SEER age-adjusted incidence rates and trends, along with their calculated delay adjustments for SEER 9 and SEER 13 areas. The adjusted rates, factors, and trends are available for all cancers combined (malignant only except for urinary bladder), for female breast in situ, for urinary bladder (in situ and malignant combined), and for 22 malignant cancer sites: melanoma (for all races combined and whites only), lung/bronchus, colon/rectum, prostate, female breast, liver and intrahepatic bile duct, pancreas, cervix uteri, corpus and uterus, ovary, testis, kidney and renal pelvis, brain and other nervous system, Hodgkin lymphoma, non-Hodgkin lymphoma, all leukemia, esophagus, larynx, myeloma, oral cavity and pharynx, thyroid, and stomach.

For more information on cancer incidence rates adjusted for reporting delay, see <https://surveillance.cancer.gov/delay/>.

STATISTICAL SIGNIFICANCE

Errors may be made in the estimation of a given statistic. In order to test whether two groups (such as the populations of a state and the entire US) have the same or different *actual* rates, the *observed* rates for the groups are compared. Statisticians consider that a difference in observed rates can be explained by one of two hypotheses: (H_0) The actual rates are really the same, but the observed rates are different because of some combination of error-causing factors, or (H_1) the actual rates of the groups are really different. H_0 is called the **null hypothesis** (because it says there is *no* real difference); H_1 is called the **alternate hypothesis**. Typically, H_0 is rejected only if there is strong evidence in favor of H_1 . (Thus, if the observed rates are equal, we cannot reject H_0 .)

Using statistical theory, one can determine the distribution of the rate difference under the assumption that H_0 is true. Then values of the rate difference that are very unlikely to occur if H_0 is true are identified. More specifically, a small positive number, called **alpha** (α), is chosen; usually, α is 0.05 or 0.01. (Alpha is called the **significance level** of the hypothesis test.) One can then identify limits for the difference in rates such that, if H_0 is true, the probability of the difference being outside of those limits is α . If the observed difference is *outside* of these limits, then the observed result is *very unlikely* to happen if H_0 is true, so H_0 is rejected.

Another way of looking at the same process is to calculate, assuming H_0 is true, the probability that the observed difference or any greater difference would occur; this number is called the **P-value** of the observed result. If the P-value of a comparison is less than α (that is, the observed difference is *very unlikely* to happen if the null hypothesis is true), H_0 will be rejected. If the P-value of a test is greater than the significance level α , H_0 will not be rejected. When a difference in rates is sufficiently large to cause the null hypothesis to be rejected for a given value of α (usually 0.05), it is called a **statistically significant** difference.

When a null hypothesis is rejected, there remains a small chance that a wrong decision has been made. If many statistical comparisons are done, even with $\alpha = 0.01$, the chance of making at least one wrong decision becomes a concern. In testing the differences between the total US rate and the rate for each state (or for the District of Columbia) for a given cancer, 51 statistical comparisons of the type described above are performed. Based on one of Bonferroni's inequalities (if there are n events and p_i is the probability of success in event i , then $P(\text{at least 1 success}) < p_1 + \dots + p_n$) (Snedecor & Cochran, 1980; p. 115-117), the significance level α for each individual comparison was set equal to $0.01/51 \approx 0.0002$. Thus, only individual-state-to-total-US comparisons with an associated P-value less than 0.0002 are considered to be statistically significant. That is, a *very small* significance level α (0.0002) is used in order to minimize the total risk (0.01) of falsely deciding that some pair of equal rates are unequal.

Use caution in assessing statistically significant differences. Population size has an important role in any calculation of statistical significance. Some states may have estimated rates that are very close to the estimated total US rate, but because of their large population, the difference

between their estimated rate and the estimated total US rate is found to be statistically significant. In this case, the true state rate and the true US rate are almost certainly different, because the observed difference, though small, is nearly impossible if the null hypothesis (equal rates) is true. A small difference in rates, however, may have no practical importance. On the other hand, some smaller states may have estimated rates that differ substantially from the estimated total US rate, but because of their relatively small population, the differences are found to be statistically nonsignificant. When this happens, if the true state rate and the true US rate were equal, the probability of obtaining a difference at least as large as what has been observed is greater than $\alpha \approx 0.0002$. Therefore, *because the evidence against it isn't strong enough, the null hypothesis (equal rates) is not rejected.*

If the percent difference (PD) between the two rates is small, there may be some question about the importance of the difference. It is difficult to specify a minimally significant absolute PD, below which the difference would always be unimportant, because the observed PD will depend on the populations of the areas involved. It may be of value to consider the size of the PD between a state rate and the US rate in assessing the importance of a statistically significant difference.

Comparing individual state rates with the US rate and assessing statistical significance is not an appropriate procedure for assessing geographic clustering of state rates. Identification of states which may represent regional clusters of high or low rates would require additional statistical and graphical analyses.

For a number of cancers, the District of Columbia has the highest death rates. *Use caution when comparing cancer rates for the District with those from the 50 states.* The District is an entirely urban area, whereas a state includes urban, suburban, and rural areas. Mortality rates for many cancers are higher in urban areas. Also, the District has a higher percentage of blacks—51% of the total population in 2010 (US Census Bureau, 2013)—than any state. In addition, their higher mortality rates for several types of cancer elevate the overall rate for the District.

STANDARD ERRORS OF RATES

Survival rates: In the tables presenting survival estimates, the magnitude of the standard error is given as a measure of the reliability of a given rate: the greater the standard error, the more uncertainty associated with the estimated rate. In addition, if there were fewer than 25 diagnoses in the first interval of the life table constructed to calculate survival, or if all cases became lost to follow-up within an interval, a valid survival estimate could not be calculated, as is noted in the table footnotes.

The **standard error (SE)** of a relative survival estimate is obtained as follows (Ederer et al., 1961):

$$SE(CR_t) = CR_t * \text{square root of } [q_1/(e_1-d_1) + q_2/(e_2-d_2) + \dots + q_t/(e_t-d_t)]$$

where CR_t is the t -year relative survival estimate, and for $i = 1, \dots, t$,
 q_i is the probability of dying in year i after diagnosis,
 e_i is the effective number of patients at risk in year i after diagnosis, and
 d_i is the number of deaths in year i after diagnosis.

Incidence and mortality rates: The standard errors of age-adjusted incidence and mortality rates are often not specified. However, the reader can approximate the SE of a particular incidence or mortality rate by the SE of a crude incidence or mortality rate (Keyfitz, 1966), that is, the SE can be approximated by the rate divided by the square root of the number of cancer cases (or the number of deaths).

Appendix tables provide numbers of cancer diagnoses within SEER areas and numbers of deaths in the entire US, respectively, by race and sex for the most recent 5-year period. These can be used to obtain approximations of the standard errors for associated age-adjusted rates for the same time period using the above formula. To approximate the standard error of a rate for a single year, use the formula but replace the number of cancer cases or deaths with the number of cancer cases or deaths divided by 5.

DEFINITIONS

Several technical terms are used in presenting the data in this report. Their definitions are presented here to clarify them for the reader.

Incidence rate: The cancer incidence rate is the number of new cancers of a specific site/type occurring in a specified population during a year, usually expressed as the number of cancers per 100,000 persons at risk. That is,

$$\text{Incidence rate} = (\text{New cancers} / \text{Population}) * 100,000.$$

The *numerator* of the incidence rate is the number of new cancers; the *denominator* of the incidence rate is the size of the population. The number of new cancers may include multiple primary cancers occurring in one patient. The primary site reported is the site of origin and not the metastatic site. In general, the incidence rate would not include recurrences. *The population used depends on the rate to be calculated.* For cancer sites that occur in only one sex, the sex-specific population (e.g., females for cervical cancer) is used.

The incidence rate can be computed for a given type of cancer or for all cancers combined. Except for 5-year age-specific rates, all incidence rates in this report are *age-adjusted* (see below) to the 2000 US standard population (or, where appropriate, to the world standard million population). (In some previous editions of the CSR, the 1970 US standard million population was used; therefore, *incidence rates in this edition cannot be compared to rates published in those editions.*) Incidence rates are for *invasive cancer only*, unless otherwise specified.

(Exceptions are the incidence rate for cancer of the urinary bladder (where both in situ and invasive cancers are counted) and breast cancer in situ, which is shown separately.)

Death rate: The cancer death (or mortality) rate is the number of deaths with cancer given as the underlying cause of death occurring in a specified population during a year, usually expressed as the number of deaths due to cancer per 100,000 persons. That is,

$$\text{Death Rate} = (\text{Cancer Deaths} / \text{Population}) * 100,000.$$

The *numerator* of the death rate is the number of deaths; the *denominator* of the death rate is the size of the population. As with the incidence rate, *the population used depends on the rate to be calculated*. The death rate can be computed for a given cancer site or for all cancers combined. Except for 5-year age-specific rates, all death rates in this report are *age-adjusted* (see below) to the 2000 US standard population (or, where appropriate, to the world standard million population). (In some previous editions of the *CSR*, the 1970 US standard million population was used; therefore, *death rates in this edition cannot be compared to rates published in those editions*.)

Age distribution: A table showing a partition of the entire lifespan into disjoint age intervals, along with the proportion of the population in each interval.

Median age: The age at which half of a population is younger and half is older.

Standard population: A **standard population** for a geographic area, such as the US or the world, is a table giving the proportions of the population falling into the age groups 0, 1-4, 5-9, ..., 80-84, and 85+. A **standard million population** for a geographic area is a table giving the number of persons in each age group 0, 1-4, ... , 85+ out of a theoretical cohort of 1,000,000 persons that is distributed by age in the same proportions as the standard population. Table A-7 shows the US 2000 standard population and the world standard million population. (Some World Health Organization mortality publications use a different world standard million population.)

Age-adjusted rate: An age-adjusted incidence or mortality rate is a weighted average of the age-specific incidence or mortality rates, where the weights are the counts of persons in the corresponding age groups of a standard population. The potential confounding effect of age is reduced when comparing age-adjusted rates based on the same standard population. For this report, the 2000 US standard population (or, where appropriate, the world standard million population) is used in computing age-adjusted rates, unless otherwise noted.

Percent change: The percent change (**PC**) in a statistic over a given time interval is

$$\text{Percent change} = (\text{Final value} - \text{Initial value}) / \text{Initial value} * 100.$$

A positive PC corresponds to an increasing trend, a negative PC to a decreasing trend.

Annual percent change: The annual percent change (**APC**) is calculated by first fitting a

regression line to the natural logarithms of the rates (r) using calendar year (x) as a regressor variable. In this report the method of *weighted least squares* is used to calculate the regression equation. If $\ln(r) = mx + b$ is the resulting regression equation (with slope m), then **APC = 100 * (e^m - 1)**. A positive APC corresponds to an increasing trend, a negative APC to a decreasing trend.

Because the methods used in their calculation are mathematically different, *the signs of the PC and the APC for a given statistic and time interval may differ*, as occurs in a few of the tables presented. That is, one of these statistics may show an increasing trend, the other a decreasing trend.

Testing the hypothesis that the actual mean annual percent change is 0 is equivalent to testing the hypothesis that the theoretical slope estimated by the slope m of the line representing the equation $\ln(r) = mx + b$ is 0. The latter hypothesis is tested using the t distribution of m / SE_m with $n - 2$ degrees of freedom. The standard error of m , called SE_m , is obtained from the fit of the regression (Kleinbaum et al., 1988). (This calculation assumes that the rates increased or decreased at a constant rate over the entire calendar year interval; the validity of this assumption was not assessed.) In those few instances where at least one of the rates was 0, the linear regression was not calculated.

Average Annual Percent Change: The average annual percent change (**AAPC**) is a summary measure of a trend over a pre-specified fixed interval based on an underlying joinpoint model. It allows us to use a single number to describe the average trend over a period of multiple years. It can be estimated even if the joinpoint model indicates that there were changes in trends during those years, since it is estimated as a weighted average of the joinpoint APCs, with the weights equal to the lengths of each subinterval over the pre-specified fixed interval.

Life table: A table for a given population listing, for each sex and each age from 0 to 120, how many members die at that age and how many survive one more year.

Observed survival: The observed survival estimate represents the proportion of cancer patients surviving for a specified time interval after diagnosis. Note that some of those not surviving died of the given cancer and some died of other causes.

Relative survival: The relative survival estimate is calculated using a procedure (Ederer et al., 1961; Ederer and Heise, 1959) whereby the observed survival estimate is adjusted for expected mortality. The relative survival estimate approximates the likelihood that a patient will not die from causes associated specifically with the given cancer before some specified time after diagnosis. It is always larger than the observed survival estimate for the same group of patients.

Standard error: The standard error of a rate is a measure of the sampling variability of the rate.

Person-years of life lost: The person-years of life lost (**PYLL**) is calculated as follows: For each

individual who dies of the cancer of interest, the number of years of expected additional life for an average person of that age, race, and sex is obtained from life tables for the US population (available from the NCHS). The PYLL in the general population associated with a particular cancer for a given year is simply the sum of this expectation over all those individuals who died of that cancer in that year.

Average years of life lost: The average years of life lost (**AYLL**) associated with a particular cancer for a given year is the PYLL associated with that cancer in the general population divided by the number of deaths from that cancer in the general population in that year.

Prevalence: Prevalence is defined as the number or percent of people alive on a certain date in a population who previously had a diagnosis of the disease. It includes new (incident) and pre-existing cases and is a function of past incidence, past survival, and the size and age structure of the population. *Limited-duration prevalence* represents the proportion of people alive on a certain day who had a diagnosis of the disease within the past x years (e.g. $x = 5, 10,$ or 20 years). *Complete prevalence* is an estimate of the number of persons (or the proportion of the population) alive on a specified date who had been diagnosed with the given disease, no matter how long ago that diagnosis was. For more details on cancer prevalence definitions and methods, refer to <https://surveillance.cancer.gov/prevalence/>.

Stage of disease at diagnosis: Extent-of-disease information determines stage of disease at diagnosis. The **SEER summary stage** presented has four levels. An invasive neoplasm confined entirely to the organ of origin is said to be **localized**. A neoplasm that has extended beyond the limits of the organ of origin, either directly into surrounding organs or tissues or into regional lymph nodes, is said to be **regional**. A neoplasm that has spread to parts of the body remote from the primary tumor, either by direct extension or by discontinuous metastasis, is said to be **distant**. When information is not sufficient to assign a stage, a neoplasm is said to be **unstaged**. In situ tumors (except those of the cervix uteri) are also collected by SEER but generally are not published in this series. For some cancers and diagnosis years, the extent of disease information can also be converted to Stages 0-IV as defined by the American Joint Committee on Cancer (Greene et al, 2002; Edge et al., 2010).

SOFTWARE USED TO GENERATE THE SEER CANCER STATISTICS REVIEW

The SEER Cancer Statistics Review includes statistics generated by a variety of statistical software including:

- [SEER*Stat](#), statistical software for the analysis of SEER and other cancer databases, was used to generate incidence, mortality, prevalence, and survival statistics presented in the CSR.
- Analysis generated by the [Joinpoint Regression Program](#) are presented to better describe trends that are not constant over time.
- The [DevCan](#) system generated the probability of developing cancer from twelve SEER areas and the probability of dying from cancer from the total United States.
- The [ComPrev](#) software was used to calculate complete prevalence estimates.

Additional statistics can be obtained via SEER's [Cancer Query Systems](#). These data retrieval applications provide access to pre-calculated cancer statistics stored in online databases.

REFERENCES

American Cancer Society. *Cancer Facts & Figures 2012*. Atlanta: American Cancer Society; 2012.

Baquet CR, Horm JW, Gibbs T, Greenwald P. Socioeconomic factors and cancer incidence among blacks and whites. *J Natl Cancer Inst* 1991; 83:551-557.

Breslow L (Chairman, Extramural Committee to Assess Measures of Progress Against Cancer). Measurement of progress against cancer: Final report to the Senate Appropriations Committee. Bethesda: National Cancer Institute; 1988.

Brookmeyer R, Damiano A. Statistical methods for short-term projections of AIDS incidence. *Stat Med* 1989;8:23-34.

Byrne J, Kessler LG, Devesa SS. The prevalence of cancer among adults in the United States: 1987. *Cancer* 1992;68:2154-9.

Capocaccia R, De Angelis R. Estimating the completeness of prevalence based on cancer registry data. *Stat Med* 1997;16:425-40.

Cho H, Howlader N, Mariotto AB, Cronin KA. Estimating relative survival for cancer patients from the SEER Program using expected rates based on Ederer I versus Ederer II method. Surveillance Research Program, National Cancer Institute; 2012. Technical Report #2012-01.

Clegg LX, Feuer EJ, Midthune D, Fay MP, Hankey BF. Impact of reporting delay and reporting error on cancer incidence rates and trends. *J Natl Cancer Inst* 2002;94:1537-1545.

Clegg L, Gail M, Feuer EJ. Estimating the variance of disease prevalence estimates from population-based registries. *Biometrics* 2002;58(3):684-8.

Day JC. *Population Projections of the United States by Age, Sex, Race, and Hispanic Origin: 1995 to 2050*, US Census Bureau, Current Population Reports, P25-1130, US Government Printing Office, Washington, DC, 1996. Available from: <https://www.census.gov/prod/1/pop/p25-1130/p251130.pdf>

Ederer F, Axtell LM, Cutler SJ. The relative survival rate: A statistical methodology. *J Natl Cancer Inst Monogr* 1961;6:101-121.

Ederer F, Heise H. Instructions to IBM 650 Programmers in Processing Survival Computations, Technical, End Results Evaluation Section, National Cancer Institute, 1959.

Edge SB, Byrd DR, Compton CC, Fritz AG, Greene FL, Trotti A III. *AJCC Cancer Staging Manual*, 7th ed. New York (NY): Springer; 2010.

Elandt-Johnson RC, Johnson NL. *Survival Models and Data Analysis*. New York (NY): Wiley; 1980.

Fay MP. Estimating age conditional probability of developing disease from surveillance data. *Popul Health Metr.* 2004 Jul 27;2(1):6. Available from: <https://pophealthmetrics.biomedcentral.com/articles/10.1186/1478-7954-2-6>

Fay MP, Pfeiffer R, Cronin KA, Le C, Feuer EJ. Age-conditional probabilities of developing cancer. *Stat Med.* 2003;22(11):1837-48.

Feinstein AR, Sosin DM, Wells CK. The Will Rogers phenomenon: Stage migration and new diagnostic techniques as a source of misleading statistics for survival of cancer. *New Engl J Med* 1985;312:1604-1608.

Feldman AR, Kessler L, Myers M, Naughton MD. The prevalence of cancer: Estimates based on the Connecticut Tumor Registry. *New Engl J Med* 1986; 315:1394-1397.

Feuer EJ, Wun L-M, Boring CC. Probability of developing cancer. In: Miller BA, Ries LAG, Hankey BF, Kosary CL, Edwards BK, editors. *Cancer Statistics Review: 1973-1989*. National Cancer Institute, NIH Pub. No. 92-2789, 1992. p. 1-8.

Feuer EJ, Wun L-M, Boring CC, Flanders WD, Timmel MJ, Tong T. The lifetime risk of developing breast cancer. *J Natl Cancer Inst* 1993;85:892-897.

Fritz A, Percy C, Jack A, Shanmugaratnam K, Sobin L, Parkin DM, Whelan S, editors. *International Classification of Diseases for Oncology*, 3rd ed. Geneva: World Health Organization; 2000.

Gail MH, Kessler L, Midthune D, Scoppa S. Two approaches for estimating disease prevalence from population-based registries of incidence and total mortality. *Biometrics* 1999;55:1137-44.

Greene FL, Page DL, Fleming ID, Fritz AG, Balch CM, Haller DG, Morrow M, editors. *AJCC Cancer Staging Manual*, 6th ed. New York (NY): Springer; 2002.

Hahn RA, Mulinare J, Teutsch SM. Inconsistencies in coding of race and ethnicity between birth and death in US infants. *JAMA* 1992;267:259-263.

Hakulinen T. Cancer survival corrected for heterogeneity in patient withdrawal. *Biometrics* 1982;38:933-942.

Harris JE. Reporting delays and the incidence of AIDS. *J Am Stat Assoc* 1990;85:915-924.

Howlader N, Ries LAG, Mariotto AB, Reichman ME, Ruhl J, Cronin KA. Improved estimates of cancer-specific survival rates from population-based data. *J Natl Cancer Inst* 2010;102:1-15.

Howlader N, Ries LAG, Stinchcomb DG, Edwards BK. The impact of underreported Veterans Affairs data on national cancer statistics: analysis using population-based SEER registries. *J Natl Cancer Inst* 2009;101(7):533-536.

Ingram DD, Parker JD, Schenker N, Weed JA, Hamilton B, Arias E, Madans JH. United States Census 2000 population with bridged race categories. *Vital Health Stat 2*. 2003 Sep;(135):1-55.

Keyfitz N. Sampling variance of standardized mortality rates. *Hum Biol* 1966;38:309-317.

Kim H-J, Fay MP, Feuer EJ, Midthune DN. Permutation tests for joinpoint regression with applications to cancer rates. *Stat Med* 2000;19:335-351.

Kleinbaum DG, Kupper LL, Muller KE. *Applied Regression Analysis and Other Multivariable Methods*, 2nd ed. Boston: PWS-Kent, 1988.

Mariotto A, Gigli A, Capocaccia R, Clegg L, Scoppa S, Ries LA, Tesaro GS, Rowland JS, Feuer EJ. Complete and limited duration prevalence estimates. *SEER Cancer Statistics Review, 1973-1999*. 2002;19.

Merrill RM, Feuer EJ, Capocaccia R, Mariotto A. Cancer prevalence estimates based on tumor registry data in the SEER Program. *Int J Epidemiol* 2000;29:197-207.

Midthune DN, Fay MP, Clegg LX, Feuer EJ. Modeling reporting delays and reporting corrections in cancer registry data. *J Am Stat Assoc* 2005;100(469):61-70.

Pagano M, Tu XM, De Gruttola V, MaWhinney S. Regression analysis of censored and truncated data: estimating reporting-delay distributions and AIDS incidence from surveillance data. *Biometrics* 1994;50:1203-1214.

Percy C, Ries LAG, Van Holten VD. The accuracy of liver cancer as the underlying cause of death on death certificates. *Public Health Rep* 1990;105:361-368.

Percy C, Stanek E, Gloeckler L. Accuracy of cancer death certificates and its effect on cancer mortality statistics. . *Am J Public Health* 1981;71: 3242-3250.

Percy C, Van Holten V, Muir C, editors. *International Classification of Diseases for Oncology*, 2nd ed. Geneva: World Health Organization;1990.

Ries LAG, Eisner MP, Kosary CL, Hankey BF, Miller BA, Clegg LX, Edwards BK (eds). *SEER Cancer Statistics Review, 1973-1997*. National Cancer Institute. NIH Pub. No. 00-2789. Bethesda, MD, 2000.

Robinson JG, West KK, Adlakha A. Coverage of the population in Census 2000: Results from demographic analysis. *Population Res Policy Rev* 2002;21:19-38.

Rosenberg HM, Maurer JD, Sorlie PD, Johnson NJ, MacDorman MF, Hoyert DL, Spitler JF, Scott C. Quality of death rates by race and Hispanic origin: A summary of current research. Hyattsville (MD): National Center for Health Statistics; Vital and Health Statistics, Series 2, No. 128, 1999.

Siegel R, Naishadham D, Jemal A. Cancer statistics, 2012. *CA Cancer J Clin* 2012 Jan-Feb;62(1):10-29.

Simonetti A, Gigli A, Capocaccia R, Mariotto A. Estimating complete prevalence of cancers diagnosed in childhood. *Stat Med* 2008 Mar 30;27(7):990-1007.

Snedecor GW, Cochran WG. *Statistical Methods*, 7th ed. Ames (IA): Iowa State University Press; 1980.

US Cancer Statistics Working Group. *United States Cancer Statistics: 1999-2002 Incidence and Mortality Web-based Report Version*. Atlanta: Department of Health and Human Services, Centers for Disease Control and Prevention, and National Cancer Institute; 2005. Available from: <https://www.cdc.gov/cancer/npcr/uscs/index.htm>

US Census Bureau. Current Population Reports; Series P-25 No. 985. Washington (DC): US Government Printing Office; 1986.

US Census Bureau: State and County QuickFacts. Data derived from Population Estimates, Census of Population and Housing, Small Area Income and Poverty Estimates, State and

County Housing Unit Estimates, County Business Patterns, Nonemployer Statistics, Economic Census, Survey of Business Owners, Building Permits, Consolidated Federal Funds Report. Last Revised: Thursday, 04-Nov-2010 12:46:18 EDT.

Zelen M. Theory of early detection of breast cancer in the general population. In: Heuson J-C, Mattheiem WH, Rozenzweig M, editors. *Breast Cancer: Trends in Research and Treatment*. New York (NY): Raven Press; 1976. p. 287-299.

Zhang, S, et al. Confidence intervals for ranks of age-adjusted rates across states or counties. *Statistics in Medicine*. 33(11): 1853-66.

Zou J, Huang L, Midthune D, Horner MJ, Krapcho M, Feuer EJ. Effect of reporting year on delay modeling. Statistical Research and Applications Branch, National Cancer Institute; 2009. Technical Report #2009-01. Available from: <https://surveillance.cancer.gov/reports/>.

Table 1.1

Estimated New Cancer Cases and Deaths for 2018
All Races, By Sex

Primary Site	Estimated New Cases			Estimated Deaths		
	Total	Males	Females	Total	Males	Females
All Sites	1,735,350	856,370	878,980	609,640	323,630	286,010
Oral Cavity and Pharynx	51,540	37,160	14,380	10,030	7,280	2,750
Tongue	17,110	12,490	4,620	2,510	1,750	760
Mouth	13,580	7,980	5,600	2,650	1,770	880
Pharynx	17,590	14,250	3,340	3,230	2,480	750
Other Oral Cavity	3,260	2,440	820	1,640	1,280	360
Digestive System	319,160	181,960	137,200	160,820	94,230	66,590
Esophagus	17,290	13,480	3,810	15,850	12,850	3,000
Stomach	26,240	16,520	9,720	10,800	6,510	4,290
Small Intestine	10,470	5,430	5,040	1,450	810	640
Colon ^a	97,220	49,690	47,530	50,630	27,390	23,240
Rectum	43,030	25,920	17,110			
Anus, Anal Canal, and Anorectum	8,580	2,960	5,620	1,160	480	680
Liver and Intrahepatic Bile Duct	42,220	30,610	11,610	30,200	20,540	9,660
Gallbladder and Other Biliary	12,190	5,450	6,740	3,790	1,530	2,260
Pancreas	55,440	29,200	26,240	44,330	23,020	21,310
Other Digestive	6,480	2,700	3,780	2,610	1,100	1,510
Respiratory System	253,290	136,400	116,890	158,770	87,200	71,570
Larynx	13,150	10,490	2,660	3,710	2,970	740
Lung and Bronchus	234,030	121,680	112,350	154,050	83,550	70,500
Other Respiratory	6,110	4,230	1,880	1,010	680	330
Bones and Joints	3,450	1,940	1,510	1,590	930	660
Soft Tissue	13,040	7,370	5,670	5,150	2,770	2,380
Skin (excl. basal & squamous)	99,550	60,350	39,200	13,460	9,070	4,390
Melanoma of the Skin ^b	91,270	55,150	36,120	9,320	5,990	3,330
Other non-epithelial skin	8,280	5,200	3,080	4,140	3,080	1,060
Breast ^b	268,670	2,550	266,120	41,400	480	40,920
Genital Organs	286,390	176,320	110,070	62,330	30,210	32,120
Cervix (uterus)	13,240		13,240	4,170		4,170
Endometrium (uterus)	63,230		63,230	11,350		11,350
Ovary	22,240		22,240	14,070		14,070
Vulva	6,190		6,190	1,200		1,200
Vagina and other genital organs, female	5,170		5,170	1,330		1,330
Prostate	164,690	164,690		29,430	29,430	
Testis	9,310	9,310		400	400	
Penis and other genital organs, male	2,320	2,320		380	380	
Urinary System	150,350	107,600	42,750	33,170	23,110	10,060
Urinary Bladder	81,190	62,380	18,810	17,240	12,520	4,720
Kidney and Renal Pelvis	65,340	42,680	22,660	14,970	10,010	4,960
Ureter and other urinary organs	3,820	2,540	1,280	960	580	380
Eye and Orbit	3,540	2,130	1,410	350	190	160
Brain and Other Nervous System	23,880	13,720	10,160	16,830	9,490	7,340
Endocrine System	56,430	14,350	42,080	3,080	1,490	1,590
Thyroid	53,990	13,090	40,900	2,060	960	1,100
Other Endocrine	2,440	1,260	1,180	1,020	530	490
Lymphoma	83,180	46,570	36,610	20,960	12,130	8,830
Hodgkin Lymphoma	8,500	4,840	3,660	1,050	620	430
Non-Hodgkin Lymphoma	74,680	41,730	32,950	19,910	11,510	8,400
Myeloma	30,770	16,400	14,370	12,770	6,830	5,940
Leukemia	60,300	35,030	25,270	24,370	14,270	10,100
Acute lymphocytic leukemia	5,960	3,290	2,670	1,470	830	640
Chronic lymphocytic leukemia	20,940	12,990	7,950	4,510	2,790	1,720
Acute myeloid leukemia	19,520	10,380	9,140	10,670	6,180	4,490
Chronic myeloid leukemia	8,430	4,980	3,450	1,090	620	470
Other leukemia	5,450	3,390	2,060	6,630	3,850	2,780
All Other Sites ^c	31,810	16,520	15,290	44,560	23,950	20,610

Cancer Facts & Figures - 2018, American Cancer Society (ACS), Atlanta, Georgia, 2018.
Excludes basal and squamous cell skin and *in situ* carcinomas except urinary bladder.

Estimated new cases are based on 2000-2014 incidence rates reported by the North American Association of Central Cancer Registries (NAACCR).

Estimated deaths are based on 2001-2015 US mortality data, National Center for Health Statistics, Centers for Disease Control and Prevention.

^a Estimated deaths for colon & rectum cancers are combined.

^b Carcinoma *in situ* of the breast accounts for about 63,960 new cases annually, and melanoma *in situ* accounts for about 87,290 new cases annually.

^c More deaths than cases suggests lack of specificity in recording underlying causes of death on death certificate.

Table 1.3

66-Year Trends in U.S. Cancer Death Rates^a

All Races, Males and Females

All Primary Cancer Sites Combined

Age Group	1950	1982	2015	Annual Percent Change		Total Percent Change
				1950-1982	1982-2015	1950-2015
Ages 0-4	11.1	4.4	2.0	-3.3*	-2.1*	-81.6
Ages 5-14	6.7	4.2	2.1	-1.9*	-1.7*	-68.4
Ages 15-24	8.6	5.8	3.3	-1.4*	-1.4*	-61.4
Ages 25-34	20.4	13.5	8.6	-1.4*	-1.7*	-58.0
Ages 35-44	63.6	48.2	27.0	-0.9*	-2.0*	-57.5
Ages 45-54	174.2	171.4	96.5	0.0	-1.8*	-44.6
Ages 55-64	391.3	435.5	282.0	0.4*	-1.8*	-27.9
Ages 65-74	710.0	832.7	609.0	0.6*	-1.3*	-14.2
Ages 75-84	1,167.2	1,249.3	1,095.6	0.3*	-0.6*	-6.1
Ages 85+	1,450.7	1,598.7	1,628.3	0.5*	-0.2*	12.2
All Ages	195.4	208.3	158.7	0.2*	-1.1*	-18.8

Lung and Bronchus Cancer^b

Age Group	1950	1982	2015	Annual Percent Change		Total Percent Change
				1950-1982	1982-2015	1950-2015
Ages 0-4	-	-	-	-	-	-
Ages 5-14	-	-	-	-	-	-
Ages 15-24	0.2	0.1	0.1	-2.8*	-0.3	-60.7
Ages 25-34	0.8	0.7	0.3	-1.0*	-2.9*	-61.6
Ages 35-44	4.6	8.9	2.5	1.5*	-3.3*	-44.8
Ages 45-54	20.2	52.1	19.5	2.7*	-2.9*	-3.5
Ages 55-64	48.9	143.4	76.3	3.0*	-2.7*	56.2
Ages 65-74	59.4	246.3	186.5	3.8*	-1.3*	213.8
Ages 75-84	55.4	255.0	306.4	4.5*	0.1	453.2
Ages 85+	42.3	187.4	316.9	4.7*	1.2*	649.3
All Ages	14.9	51.7	40.7	3.5*	-1.1*	172.5

Source: US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

^a Rates are per 100,000 and age-adjusted to the 2000 US Std Population (18 age groups - Census P25-1130).

^b Due to coding changes throughout the years, Lung and Bronchus includes trachea and pleura.

* The APC is significantly different from zero (p<.05).

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Trend based on less than 10 cases for at least one year within the time interval.

Table 1.4

Summary of Changes in Cancer Mortality, 1950-2015 and
5-Year Relative Survival (Percent), 1950-2014
Males and Females, By Primary Cancer Site

Primary Site	Whites			
	U.S. Mortality Percent Change 1950-2015 ^a		5-Year Relative Survival (Percent) ^b	
	Total	APC	1950-1954	2008-2014
Oral cavity and pharynx	-49.7	-1.3*	46	67.2
Esophagus	25.2	0.7*	4	21.3
Stomach	-88.7	-3.3*	12	28.1
Colon and rectum	-58.0	-1.3*	37	67.3
Colon	-52.3	-1.1*	41	67.1
Rectum	-71.0	-2.2*	40	68.1
Liver and intrahepatic bile duct	63.5	0.9*	1	16.8
Pancreas	27.8	0.1*	1	7.4
Larynx	-43.3	-0.8*	52	64.1
Lung and bronchus	173.3	1.0*	6	18.3
Males	100.2	0.3	5	15.7
Females	489.1	2.4*	9	21.1
Melanoma of the skin	147.2	1.1*	49	93.3
Breast(females)	-39.3	-0.7*	60	92.3
Cervix uteri	-81.6	-3.1*	59	71.9
Corpus and uterus, NOS	-63.9	-1.4*	72	85.9
Ovary	-20.1	-0.4*	30	44.7
Prostate	-39.2	-0.6*	43	99.8
Testis	-72.7	-2.7*	57	97.4
Urinary bladder	-28.8	-0.6*	53	80.6
Kidney and renal pelvis	30.4	0.4*	34	74.6
Brain and nervous system	56.3	0.4*	21	33.7
Thyroid	-40.6	-0.9*	80	98.0
Hodgkin lymphoma	-83.2	-3.3*	30	89.2
Non-Hodgkin lymphoma	64.0	0.6*	33	72.4
Myeloma	209.4	1.0*	6	46.0
Leukemia	-9.6	-0.3*	10	63.3
Childhood (Ages 0-14)	-74.2	-2.6*	20	84.6
All Sites	-18.8	-0.2*	35	69.7

The APC is the Annual Percent Change over the time interval.

Rates used in the calculation of the APC are age-adjusted to the 2000 U.S. standard population (18 age groups - Census P25-1130).

^a U.S. Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

Due to coding changes throughout the years: Colon excludes other digestive tract; Rectum includes anal canal; Liver & intrahepatic bile duct includes gallbladder & biliary tract, NOS; Lung & bronchus includes trachea & pleura; Ovary includes fallopian tube; Urinary bladder includes other urinary organs; Kidney & Renal pelvis includes ureter; NHL and myeloma each include a small number of leukemias; NHL includes a small number of ill-defined sites.

^b Survival estimates for 1950-54 are from NCI Survival Report 5 with the exception of All Sites, Oral cavity & pharynx, Colon & rectum, Non-Hodgkin lymphoma and Childhood cancers which come from historical Connecticut data. Survival estimates for 2008-2014 are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Rates are based on follow-up of patients into 2015.

* The APC is significantly different from zero (p<.05).

Table 1.5
Age-Adjusted SEER Incidence and U.S. Death Rates and 5-Year Relative Survival (Percent)
By Primary Cancer Site, Sex and Time Period

All Races

Site	Incidence ^a (2011-2015)			US Mortality ^b (2011-2015)			Survival ^c (%) (2008-2014)		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
All Sites	439.2	483.0	409.9	163.5	196.8	139.6	66.9	66.4	67.5
Oral Cavity & Pharynx:	11.3	17.1	6.3	2.5	3.9	1.3	64.8	64.0	66.9
Lip	0.7	1.1	0.3	0.0	0.0	0.0	88.4	88.2	89.3
Tongue	3.4	5.2	1.8	0.6	0.9	0.4	65.8	66.1	64.8
Salivary gland	1.3	1.7	1.0	0.3	0.4	0.1	71.6	64.0	82.1
Floor of mouth	0.5	0.7	0.3	0.0	0.0	0.0	52.9	51.9	55.2
Gum & other oral cavity	1.5	1.8	1.3	0.4	0.5	0.3	59.2	55.2	64.3
Nasopharynx	0.6	0.9	0.4	0.2	0.3	0.1	61.6	59.4	66.8
Tonsil	2.0	3.4	0.7	0.2	0.4	0.1	73.9	74.5	70.9
Oropharynx	0.4	0.7	0.2	0.3	0.4	0.1	45.8	47.1	40.8
Hypopharynx	0.6	1.0	0.2	0.1	0.2	0.0	32.9	32.9	32.5
Other oral cavity & pharynx	0.3	0.4	0.1	0.4	0.7	0.2	45.1	47.7	35.5
Digestive System:	81.1	98.8	66.3	41.3	53.0	31.7	43.5	41.2	46.3
Esophagus	4.2	7.2	1.7	4.0	7.2	1.5	19.2	18.9	20.0
Stomach	7.2	9.8	5.2	3.2	4.3	2.3	31.0	28.4	35.1
Small intestine	2.3	2.6	2.0	0.4	0.5	0.3	67.6	66.8	68.4
Colon & Rectum:	39.4	45.2	34.5	14.5	17.3	12.2	64.5	64.1	64.9
Colon	27.7	30.7	25.3	-	-	-	63.6	63.6	63.6
Rectum	11.7	14.6	9.2	-	-	-	66.6	65.2	68.6
Anus, anal canal & anorectum	1.8	1.5	2.1	0.3	0.2	0.3	67.4	60.8	71.4
Liver & intrahepatic bile duct	8.8	13.6	4.7	6.4	9.4	3.8	17.7	17.5	18.4
Gallbladder	1.2	0.9	1.4	0.6	0.5	0.7	18.2	18.4	18.1
Other biliary	1.9	2.3	1.5	0.4	0.5	0.4	17.5	18.8	16.1
Pancreas	12.6	14.4	11.2	10.9	12.6	9.5	8.5	8.8	8.3
Retroperitoneum	0.4	0.4	0.4	0.1	0.1	0.1	54.2	52.9	55.4
Peritoneum, omentum & mesentery	0.5	0.1	0.9	0.3	0.1	0.4	32.0	38.2	31.5
Other digestive system	0.7	0.8	0.6	0.3	0.4	0.3	8.6	7.1	10.2
Respiratory System:	58.5	70.2	49.5	44.6	56.0	35.9	21.6	20.2	23.3
Nose, nasal cavity & middle ear	0.7	0.9	0.5	0.1	0.2	0.1	58.2	59.2	56.6
Larynx	3.0	5.2	1.1	1.0	1.8	0.4	60.9	61.4	58.4
Lung & bronchus	54.6	63.8	47.8	43.4	53.8	35.4	18.6	15.5	22.0
Pleura ^d	0.0	0.0	0.0	0.1	0.1	0.0	26.2	24.7	28.3
Trachea & other respiratory organs	0.2	0.2	0.1	0.1	0.1	0.0	53.2	53.6	52.2
Bones & joints	0.9	1.1	0.8	0.4	0.5	0.3	66.9	65.1	69.2
Soft tissue (including heart)	3.4	4.1	2.9	1.3	1.5	1.2	64.5	63.8	65.3
Skin (excl. basal & squamous):	24.9	32.7	19.2	3.6	5.6	2.1	91.2	89.3	93.6
Melanoma of the skin	22.8	29.8	17.7	2.6	3.9	1.6	91.8	89.9	94.2
Other non-epithelial skin	2.1	2.9	1.5	1.0	1.7	0.5	84.3	82.5	86.7
Breast	67.5	1.2	126.0	11.6	0.3	20.9	89.6	83.0	89.7
Breast (<i>in situ</i>)	16.4	0.1	31.1	-	-	-	100.0	100.0	100.0

Note: Incidence and death rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^a SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

^b US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

^c SEER 18 areas. Based on follow-up of patients into 2015. [Expected survival rates](#) are derived from life tables by socio-economic status, geography and race developed by the SEER program.

^d Mesotheliomas of the Pleura are included in the separate group Mesothelioma for incidence but are included in the Pleura grouping for mortality.

- Statistic could not be calculated due to less than 16 cases in the time interval.

Table 1.5 - continued
 Age-Adjusted SEER Incidence and U.S. Death Rates and 5-Year Relative Survival (Percent)
 By Primary Cancer Site, Sex and Time Period

All Races

Site	Incidence ^a (2011-2015)			US Mortality ^b (2011-2015)			Survival ^c (%) (2008-2014)		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
Female Genital System:	26.2	-	49.4	8.3	-	15.1	69.3	-	69.3
Cervix uteri	3.8	-	7.4	1.2	-	2.3	66.2	-	66.2
Corpus uteri	13.3	-	25.2	1.1	-	2.0	82.5	-	82.5
Uterus, NOS	0.4	-	0.8	1.5	-	2.7	29.6	-	29.6
Ovary ^d	6.2	-	11.6	4.0	-	7.2	47.4	-	47.4
Vagina	0.4	-	0.7	0.1	-	0.2	47.0	-	47.0
Vulva	1.4	-	2.5	0.3	-	0.5	71.0	-	71.0
Other female genital system	0.7	-	1.3	0.2	-	0.3	53.6	-	53.6
Male Genital System:	54.9	119.6	-	8.1	20.0	-	97.9	97.9	-
Prostate	51.5	112.6	-	7.9	19.5	-	98.2	98.2	-
Testis	2.9	5.7	-	0.1	0.2	-	95.3	95.3	-
Penis	0.4	0.9	-	0.1	0.2	-	67.2	67.2	-
Other male genital system	0.1	0.3	-	0.0	0.0	-	84.9	84.9	-
Urinary System:	36.3	57.4	19.8	8.5	13.6	4.7	75.3	75.9	73.8
Urinary bladder	19.5	34.3	8.3	4.4	7.6	2.2	76.8	78.1	72.9
Kidney & renal pelvis	15.9	21.7	10.9	3.8	5.6	2.4	74.5	73.7	75.8
Ureter	0.5	0.8	0.4	0.1	0.2	0.1	47.1	48.3	45.4
Other urinary system	0.4	0.6	0.2	0.1	0.2	0.1	48.6	54.2	37.7
Eye & Orbit	0.9	1.0	0.7	0.1	0.1	0.1	82.9	82.4	83.6
Brain & Nervous System: ^e	6.4	7.5	5.4	4.4	5.3	3.6	33.2	32.0	34.7
Brain	6.0	7.1	5.0	-	-	-	30.1	29.4	31.1
Cranial nerves & other nervous system	0.4	0.4	0.4	-	-	-	78.9	75.6	82.0
Endocrine System:	15.2	8.1	22.1	0.8	0.8	0.7	96.3	92.1	97.7
Thyroid	14.5	7.3	21.4	0.5	0.5	0.5	98.1	95.6	98.8
Other endocrine & thymus	0.7	0.8	0.7	0.3	0.3	0.3	64.5	64.5	64.4
Lymphoma:	21.9	26.5	18.2	6.1	7.9	4.7	73.5	72.0	75.3
Hodgkin lymphoma	2.5	2.9	2.2	0.3	0.4	0.3	86.6	85.7	87.7
Non-Hodgkin lymphoma	19.4	23.6	15.9	5.7	7.4	4.5	71.4	69.9	73.3
Myeloma	6.7	8.4	5.3	3.3	4.2	2.7	50.7	50.6	50.9
Leukemia:	13.8	17.6	10.8	6.7	9.0	5.0	61.4	62.5	60.0
Lymphocytic:	6.8	9.0	5.1	1.8	2.6	1.3	79.7	80.3	78.8
Acute lymphocytic	1.7	1.9	1.6	0.4	0.5	0.4	68.1	67.8	68.5
Chronic lymphocytic	4.7	6.4	3.3	1.3	1.9	0.8	84.2	84.4	83.8
Other lymphocytic	0.4	0.6	0.2	0.1	0.2	0.1	81.3	85.0	70.2
Myeloid & Monocytic:	6.4	8.0	5.2	3.3	4.4	2.6	40.4	40.1	40.7
Acute myeloid	4.3	5.2	3.6	2.8	3.6	2.2	27.4	26.3	28.6
Chronic myeloid	1.8	2.4	1.4	0.3	0.4	0.2	67.6	66.4	69.3
Acute monocytic	0.2	0.2	0.2	0.0	0.0	0.0	23.2	21.3	25.4
Other myeloid & monocytic	0.2	0.2	0.1	0.2	0.3	0.2	34.5	36.7	31.5
Other leukemia:	0.5	0.6	0.4	1.5	2.0	1.1	33.9	33.2	34.5
Other acute leukemia	0.2	0.3	0.2	0.5	0.7	0.4	23.5	23.5	23.4
Aleukemic, subleukemic & NOS	0.3	0.3	0.2	1.0	1.4	0.8	41.4	40.6	42.1
Kaposi Sarcoma ^f	0.5	0.9	0.1	-	-	-	74.1	73.9	73.5
Mesothelioma ^f	0.9	1.6	0.4	-	-	-	9.4	7.0	16.6
Ill-defined & unspecified	7.9	9.2	6.9	11.9	15.1	9.5	18.2	21.6	14.8

Note: Incidence and death rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^a SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

^b US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

^c SEER 18 areas. Based on follow-up of patients into 2015. [Expected survival rates](#) are derived from life tables by socio-economic status, geography and race developed by the SEER program.

^d Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

^e Due to coding changes, Brain & Nervous System mortality are no longer shown separately.

^f Rate not shown for mortality. Category did not exist in mortality coding until 1999.

- Statistic could not be calculated due to less than 16 cases in the time interval.

Table 1.6
Age-Adjusted SEER Incidence and U.S. Death Rates and 5-Year Relative Survival (Percent)
By Primary Cancer Site, Sex and Time Period

Site	Whites								
	Incidence ^a (2011-2015)			US Mortality ^b (2011-2015)			Survival ^c (%) (2008-2014)		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
All Sites	448.8	488.5	423.1	163.8	196.4	140.0	67.3	66.5	68.1
Oral Cavity & Pharynx:	11.9	18.0	6.5	2.5	3.8	1.3	66.3	65.8	67.5
Lip	0.8	1.3	0.4	0.0	0.0	0.0	88.4	88.0	89.5
Tongue	3.8	5.7	2.0	0.6	1.0	0.4	67.5	68.0	66.0
Salivary gland	1.3	1.8	1.0	0.3	0.4	0.1	69.4	61.8	80.8
Floor of mouth	0.5	0.7	0.3	0.0	0.0	0.0	54.4	53.0	57.3
Gum & other oral cavity	1.5	1.8	1.3	0.4	0.5	0.3	58.9	55.2	63.5
Nasopharynx	0.4	0.6	0.2	0.1	0.2	0.1	58.5	57.1	61.9
Tonsil	2.2	3.8	0.8	0.2	0.4	0.1	75.5	75.9	73.3
Oropharynx	0.5	0.8	0.2	0.3	0.4	0.1	49.3	51.0	42.4
Hypopharynx	0.6	1.0	0.2	0.1	0.2	0.0	35.9	36.0	35.0
Other oral cavity & pharynx	0.3	0.5	0.1	0.4	0.7	0.2	49.1	51.6	39.6
Digestive System:	79.1	96.1	64.7	40.2	51.6	30.7	43.9	41.6	46.8
Esophagus	4.5	7.7	1.7	4.3	7.6	1.5	19.9	19.9	20.3
Stomach	6.5	8.9	4.5	2.8	3.7	2.0	29.6	27.0	34.2
Small intestine	2.2	2.6	1.9	0.4	0.4	0.3	68.5	67.2	69.9
Colon & Rectum:	38.8	44.4	34.0	14.1	16.8	11.9	65.0	64.7	65.4
Colon	27.3	30.1	24.9	-	-	-	64.5	64.4	64.5
Rectum	11.5	14.2	9.1	-	-	-	66.3	65.3	67.8
Anus, anal canal & anorectum	2.0	1.5	2.4	0.3	0.2	0.3	69.0	62.5	72.6
Liver & intrahepatic bile duct	8.0	12.2	4.3	6.0	8.7	3.6	17.0	16.9	17.5
Gallbladder	1.1	0.8	1.4	0.6	0.4	0.7	18.1	17.7	18.3
Other biliary	1.8	2.3	1.5	0.4	0.5	0.4	17.6	19.5	15.3
Pancreas	12.6	14.4	11.1	10.8	12.6	9.4	8.4	8.7	8.0
Retroperitoneum	0.4	0.4	0.3	0.1	0.1	0.1	53.5	52.6	54.4
Peritoneum, omentum & mesentery	0.6	0.1	1.0	0.3	0.1	0.4	31.5	36.9	31.0
Other digestive system	0.7	0.8	0.6	0.3	0.4	0.3	8.9	7.0	10.6
Respiratory System:	60.0	70.4	52.0	45.4	56.0	37.2	21.8	20.4	23.3
Nose, nasal cavity & middle ear	0.7	0.9	0.5	0.1	0.2	0.1	60.3	62.2	57.4
Larynx	3.0	5.3	1.1	1.0	1.7	0.4	62.1	62.7	59.6
Lung & bronchus	56.1	63.9	50.2	44.1	53.9	36.6	18.7	15.7	22.0
Pleura ^d	0.0	0.0	0.0	0.1	0.1	0.0	21.4	25.0	14.1
Trachea & other respiratory organs	0.2	0.3	0.1	0.1	0.1	0.0	53.5	53.7	53.0
Bones & joints	1.0	1.2	0.9	0.5	0.6	0.4	66.8	64.8	69.4
Soft tissue (including heart)	3.5	4.2	2.9	1.3	1.6	1.1	65.1	64.3	66.1
Skin (excl. basal & squamous):	29.5	38.3	23.0	4.1	6.3	2.4	90.5	88.5	93.2
Melanoma of the skin	27.3	35.2	21.5	3.0	4.5	1.9	91.2	89.2	93.8
Other non-epithelial skin	2.2	3.1	1.5	1.1	1.8	0.5	81.8	80.1	84.1
Breast	68.1	1.2	128.6	11.2	0.3	20.3	90.6	84.9	90.6
Breast (<i>in situ</i>)	16.0	0.1	30.8	-	-	-	100.0	99.9	100.0

Note: Incidence and death rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^a SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

^b US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

^c SEER 18 areas. Based on follow-up of patients into 2015. [Expected survival rates](#) are derived from life tables by socio-economic status, geography and race developed by the SEER program.

^d Mesotheliomas of the Pleura are included in the separate group Mesothelioma for incidence but are included in the Pleura grouping for mortality.

- Statistic could not be calculated due to less than 16 cases in the time interval.

Table 1.6 - continued
Age-Adjusted SEER Incidence and U.S. Death Rates and 5-Year Relative Survival (Percent)
By Primary Cancer Site, Sex and Time Period

Site	Whites								
	Incidence ^a (2011-2015)			US Mortality ^b (2011-2015)			Survival ^c (%) (2008-2014)		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
Female Genital System:	26.7	-	50.9	8.1	-	15.0	70.7	-	70.7
Cervix uteri	3.7	-	7.4	1.1	-	2.2	67.7	-	67.7
Corpus uteri	13.6	-	25.9	1.0	-	1.9	84.6	-	84.6
Uterus, NOS	0.4	-	0.8	1.3	-	2.4	32.5	-	32.5
Ovary ^d	6.4	-	12.1	4.1	-	7.5	47.4	-	47.4
Vagina	0.4	-	0.7	0.1	-	0.2	47.2	-	47.2
Vulva	1.5	-	2.7	0.3	-	0.6	70.4	-	70.4
Other female genital system	0.7	-	1.3	0.2	-	0.3	52.4	-	52.4
Male Genital System:	52.9	113.7	-	7.6	18.7	-	98.0	98.0	-
Prostate	48.9	105.7	-	7.4	18.2	-	98.3	98.3	-
Testis	3.5	6.8	-	0.1	0.3	-	95.4	95.4	-
Penis	0.4	0.9	-	0.1	0.2	-	67.0	67.0	-
Other male genital system	0.1	0.3	-	0.0	0.0	-	86.2	86.2	-
Urinary System:	38.7	61.2	20.8	8.8	14.2	4.8	75.6	76.1	74.4
Urinary bladder	21.5	37.6	8.9	4.6	8.0	2.2	77.3	78.3	74.1
Kidney & renal pelvis	16.3	22.2	11.3	4.0	5.8	2.5	74.4	73.6	75.8
Ureter	0.6	0.8	0.4	0.1	0.2	0.1	48.2	48.8	47.4
Other urinary system	0.4	0.6	0.2	0.1	0.2	0.1	49.4	53.5	39.9
Eye & Orbit	1.0	1.2	0.9	0.1	0.1	0.1	82.0	81.4	82.8
Brain & Nervous System: ^e	7.1	8.3	6.0	4.8	5.8	3.9	32.0	30.7	33.6
Brain	6.6	7.9	5.6	-	-	-	29.0	28.2	30.0
Cranial nerves & other nervous system	0.4	0.4	0.4	-	-	-	80.0	76.3	83.6
Endocrine System:	15.9	8.6	23.4	0.8	0.8	0.7	96.7	92.7	98.0
Thyroid	15.3	7.8	22.8	0.5	0.5	0.5	98.3	95.8	99.0
Other endocrine & thymus	0.7	0.8	0.6	0.3	0.3	0.3	63.5	63.9	63.1
Lymphoma:	23.0	27.7	19.1	6.3	8.2	4.9	73.7	72.3	75.5
Hodgkin lymphoma	2.7	3.0	2.4	0.3	0.4	0.3	86.9	86.2	87.7
Non-Hodgkin lymphoma	20.3	24.7	16.8	6.0	7.7	4.6	71.8	70.2	73.6
Myeloma	6.1	7.9	4.7	3.1	4.0	2.4	50.0	50.1	49.8
Leukemia:	14.6	18.6	11.4	6.9	9.3	5.2	61.8	62.7	60.6
Lymphocytic:	7.4	9.7	5.5	1.9	2.7	1.3	79.9	80.4	79.3
Acute lymphocytic	1.9	2.1	1.7	0.5	0.6	0.4	68.2	67.5	69.1
Chronic lymphocytic	5.1	6.9	3.6	1.3	2.0	0.9	84.2	84.4	83.9
Other lymphocytic	0.4	0.7	0.2	0.1	0.2	0.1	82.2	85.7	71.6
Myeloid & Monocytic:	6.7	8.3	5.4	3.5	4.6	2.7	39.5	39.1	39.9
Acute myeloid	4.4	5.4	3.7	2.9	3.8	2.2	26.9	25.9	28.0
Chronic myeloid	1.9	2.4	1.4	0.3	0.4	0.2	66.4	65.4	68.0
Acute monocytic	0.2	0.3	0.2	0.0	0.0	0.0	23.8	21.7	26.4
Other myeloid & monocytic	0.2	0.2	0.1	0.2	0.3	0.2	32.0	32.6	31.2
Other leukemia:	0.5	0.6	0.4	1.5	2.1	1.2	33.7	32.3	35.1
Other acute leukemia	0.2	0.3	0.2	0.5	0.7	0.4	23.4	23.8	22.8
Aleukemic, subleukemic & NOS	0.3	0.3	0.2	1.0	1.4	0.8	41.8	39.2	44.2
Kaposi Sarcoma ^f	0.4	0.7	0.1	-	-	-	77.2	76.3	81.7
Mesothelioma ^f	1.0	1.8	0.5	-	-	-	9.2	7.0	15.9
Ill-defined & unspecified	8.1	9.4	7.0	12.1	15.3	9.6	18.9	22.9	14.8

Note: Incidence and death rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^a SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

^b US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

^c SEER 18 areas. Based on follow-up of patients into 2015. [Expected survival rates](#) are derived from life tables by socio-economic status, geography and race developed by the SEER program.

^d Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

^e Due to coding changes, Brain & Nervous System mortality are no longer shown separately.

^f Rate not shown for mortality. Category did not exist in mortality coding until 1999.

- Statistic could not be calculated due to less than 16 cases in the time interval.

Table 1.7
Age-Adjusted SEER Incidence and U.S. Death Rates and 5-Year Relative Survival (Percent)
By Primary Cancer Site, Sex and Time Period

Site	Blacks								
	Incidence ^a (2011-2015)			US Mortality ^b (2011-2015)			Survival ^c (%) (2008-2014)		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
All Sites	453.4	535.0	397.8	189.8	239.9	159.0	61.5	63.6	59.1
Oral Cavity & Pharynx:	9.0	14.0	5.1	2.8	4.8	1.3	48.1	46.0	53.0
Lip	0.1	0.1	-	-	-	-	67.9	66.8	-
Tongue	2.1	3.3	1.1	0.5	0.9	0.3	43.1	42.9	43.7
Salivary gland	1.0	1.1	1.0	0.2	0.3	0.1	76.5	70.5	81.8
Floor of mouth	0.5	0.9	0.2	0.0	0.0	-	40.3	43.4	31.4
Gum & other oral cavity	1.3	1.7	1.0	0.3	0.5	0.2	55.8	49.5	62.5
Nasopharynx	0.7	1.0	0.4	0.2	0.4	0.1	57.0	57.4	55.1
Tonsil	1.6	2.9	0.6	0.2	0.4	0.1	55.6	57.2	48.3
Oropharynx	0.6	0.9	0.3	0.4	0.7	0.2	27.5	27.0	29.2
Hypopharynx	0.9	1.6	0.3	0.2	0.3	0.0	18.1	18.2	16.8
Other oral cavity & pharynx	0.3	0.5	0.1	0.7	1.3	0.3	26.3	29.1	16.5
Digestive System:	98.4	120.9	81.6	53.1	69.6	41.2	39.4	35.8	43.3
Esophagus	4.0	6.4	2.2	3.5	5.8	1.8	12.9	11.4	16.2
Stomach	10.1	13.6	7.7	5.7	8.3	3.9	31.1	26.4	37.0
Small intestine	3.8	4.2	3.5	0.6	0.7	0.5	64.0	64.0	63.8
Colon & Rectum:	47.6	55.5	41.9	19.4	24.4	16.1	57.8	55.7	59.8
Colon	35.4	40.2	32.0	-	-	-	56.0	55.0	56.9
Rectum	12.2	15.3	9.9	-	-	-	62.6	57.3	68.7
Anus, anal canal & anorectum	1.9	2.1	1.8	0.3	0.3	0.2	58.1	52.9	63.4
Liver & intrahepatic bile duct	10.5	17.3	5.1	8.3	13.2	4.6	14.0	13.3	16.2
Gallbladder	1.6	1.3	1.9	0.9	0.7	1.0	16.5	15.3	17.0
Other biliary	1.7	2.0	1.5	0.4	0.4	0.4	13.7	11.0	16.0
Pancreas	15.5	16.9	14.3	13.3	14.8	12.2	8.6	7.6	9.5
Retroperitoneum	0.4	0.4	0.4	0.1	0.1	0.0	55.3	58.9	52.9
Peritoneum, omentum & mesentery	0.4	0.1	0.6	0.2	0.1	0.3	27.4	33.8	26.5
Other digestive system	0.9	1.0	0.8	0.4	0.6	0.3	6.5	6.0	6.4
Respiratory System:	66.6	90.6	50.0	48.1	68.7	34.2	19.3	17.7	21.3
Nose, nasal cavity & middle ear	0.6	0.9	0.4	0.1	0.2	0.1	49.6	44.1	56.6
Larynx	4.4	8.3	1.5	1.7	3.3	0.5	52.0	52.5	50.0
Lung & bronchus	61.4	81.2	47.9	46.2	65.1	33.5	16.2	13.2	19.8
Pleura ^d	-	-	-	0.0	0.1	0.0	-	-	-
Trachea & other respiratory organs	0.1	0.2	0.1	0.1	0.1	0.0	45.7	46.8	42.8
Bones & joints	0.7	0.8	0.6	0.4	0.5	0.4	65.2	66.0	64.2
Soft tissue (including heart)	3.4	3.6	3.3	1.5	1.5	1.5	59.7	59.7	59.8
Skin (excl. basal & squamous):	2.1	2.2	2.0	0.8	1.2	0.6	81.5	77.2	84.6
Melanoma of the skin	1.1	1.1	1.0	0.4	0.5	0.3	65.2	57.1	71.0
Other non-epithelial skin	1.0	1.1	1.0	0.4	0.7	0.2	93.5	90.6	95.4
Breast	72.0	1.7	126.9	16.8	0.5	28.7	81.1	70.9	81.2
Breast (<i>in situ</i>)	18.0	0.3	32.1	-	-	-	100.0	78.3	100.0

Note: Incidence and death rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^a SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

^b US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

^c SEER 18 areas. Based on follow-up of patients into 2015. [Expected survival rates](#) are derived from life tables by socio-economic status, geography and race developed by the SEER program.

^d Mesotheliomas of the Pleura are included in the separate group Mesothelioma for incidence but are included in the Pleura grouping for mortality.

- Statistic could not be calculated due to less than 16 cases in the time interval.

Table 1.7 - continued
Age-Adjusted SEER Incidence and U.S. Death Rates and 5-Year Relative Survival (Percent)
By Primary Cancer Site, Sex and Time Period

Site	Blacks								
	Incidence ^a (2011-2015)			US Mortality ^b (2011-2015)			Survival ^c (%) (2008-2014)		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
Female Genital System:	26.4	-	46.8	11.2	-	19.2	55.7	-	55.7
Cervix uteri	4.6	-	8.4	2.1	-	3.7	55.5	-	55.5
Corpus uteri	13.5	-	23.8	1.9	-	3.3	64.0	-	64.0
Uterus, NOS	0.9	-	1.5	3.0	-	5.1	20.6	-	20.6
Ovary ^d	5.3	-	9.3	3.7	-	6.3	38.5	-	38.5
Vagina	0.6	-	1.0	0.2	-	0.3	45.0	-	45.0
Vulva	1.0	-	1.7	0.2	-	0.3	71.8	-	71.8
Other female genital system	0.5	-	0.9	0.2	-	0.3	56.2	-	56.2
Male Genital System:	77.7	181.2	-	14.8	40.3	-	95.9	95.9	-
Prostate	76.4	178.3	-	14.6	39.9	-	96.1	96.1	-
Testis	0.7	1.5	-	0.1	0.1	-	91.8	91.8	-
Penis	0.4	1.0	-	0.1	0.2	-	58.4	58.4	-
Other male genital system	0.1	0.3	-	0.0	0.1	-	63.4	63.4	-
Urinary System:	31.2	46.8	20.1	7.3	11.1	4.9	70.2	71.1	68.6
Urinary bladder	12.3	20.6	6.7	3.5	5.3	2.4	63.0	67.1	55.0
Kidney & renal pelvis	18.3	25.3	12.8	3.7	5.5	2.4	75.1	74.2	76.3
Ureter	0.2	0.3	0.2	0.0	0.0	0.0	31.9	28.1	33.4
Other urinary system	0.4	0.7	0.3	0.1	0.1	0.1	32.1	44.8	18.0
Eye & Orbit	0.2	0.3	0.3	0.0	0.0	0.0	90.3	86.3	93.2
Brain & Nervous System: ^e	4.0	4.8	3.5	2.6	3.2	2.1	39.5	38.1	40.9
Brain	3.7	4.4	3.1	-	-	-	35.7	35.0	36.6
Cranial nerves & other nervous system	0.4	0.4	0.3	-	-	-	75.2	74.2	75.7
Endocrine System:	9.8	4.7	14.3	0.9	0.8	0.9	93.8	85.5	96.0
Thyroid	8.9	3.8	13.4	0.5	0.4	0.6	97.2	92.9	97.7
Other endocrine & thymus	0.9	0.8	0.9	0.4	0.4	0.3	65.3	63.1	67.1
Lymphoma:	17.0	20.5	14.3	4.5	5.8	3.6	69.7	66.7	73.2
Hodgkin lymphoma	2.6	3.1	2.2	0.3	0.4	0.2	83.9	80.4	87.8
Non-Hodgkin lymphoma	14.4	17.5	12.1	4.2	5.4	3.4	66.2	63.3	69.6
Myeloma	13.4	15.9	11.6	6.2	7.5	5.5	52.1	51.3	52.8
Leukemia:	11.0	14.0	9.0	5.6	7.4	4.5	55.4	56.9	53.6
Lymphocytic:	4.8	6.6	3.5	1.5	2.1	1.1	70.2	72.1	67.4
Acute lymphocytic	1.0	1.0	0.9	0.3	0.4	0.3	63.6	66.0	60.2
Chronic lymphocytic	3.6	5.2	2.5	1.1	1.6	0.7	73.1	74.3	71.2
Other lymphocytic	0.2	0.4	0.1	0.1	0.1	0.1	65.9	70.7	49.2
Myeloid & Monocytic:	5.7	6.8	5.0	2.7	3.4	2.2	43.8	42.7	44.8
Acute myeloid	3.8	4.5	3.3	2.2	2.7	1.8	28.8	26.5	30.9
Chronic myeloid	1.7	2.0	1.4	0.3	0.4	0.2	70.0	67.8	72.3
Acute monocytic	0.1	0.2	0.1	0.0	0.0	-	25.2	21.1	28.4
Other myeloid & monocytic	0.1	0.1	0.1	0.2	0.2	0.1	41.5	48.8	28.2
Other leukemia:	0.5	0.6	0.5	1.5	1.9	1.2	35.2	36.4	33.3
Other acute leukemia	0.2	0.3	0.2	0.4	0.5	0.3	22.0	18.0	27.1
Aleukemic, subleukemic & NOS	0.3	0.3	0.3	1.1	1.4	0.9	40.4	45.0	35.1
Kaposi Sarcoma ^f	1.1	2.1	0.2	-	-	-	63.4	65.4	33.8
Mesothelioma ^f	0.5	0.9	0.3	-	-	-	13.3	7.0	25.8
Ill-defined & unspecified	8.9	10.1	8.0	13.1	17.0	10.5	13.4	12.5	14.1

Note: Incidence and death rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^a SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

^b US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

^c SEER 18 areas. Based on follow-up of patients into 2015. [Expected survival rates](#) are derived from life tables by socio-economic status, geography and race developed by the SEER program.

^d Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

^e Due to coding changes, Brain & Nervous System mortality are no longer shown separately.

^f Rate not shown for mortality. Category did not exist in mortality coding until 1999.

- Statistic could not be calculated due to less than 16 cases in the time interval.

Table 1.8
SEER Incidence and U.S. Mortality Trends by Primary Cancer Site and Sex
All Races, 2006-2015

Site	Incidence ^a			US Mortality ^b		
	Total APC	Males APC	Females APC	Total APC	Males APC	Females APC
All Sites	-1.2*	-2.2*	-0.2*	-1.5*	-1.8*	-1.4*
Oral Cavity & Pharynx:	0.6*	0.8*	0.1	0.2	0.5	-0.8*
Lip	0.1	-0.5	1.1	0.4	-0.8	1.6
Tongue	1.5*	1.9*	0.7	0.4	0.9*	-0.6
Salivary gland	-0.1	-0.3	-0.1	1.6*	2.1*	0.3
Floor of mouth	-3.2*	-3.5*	-2.5	-6.6*	-4.7*	-10.6*
Gum & other oral cavity	-0.1	0.1	-0.4	0.5	1.0	-0.2
Nasopharynx	-1.2	-1.1	-1.4	-1.5*	-0.9*	-3.0*
Tonsil	2.5*	2.6*	1.9*	1.9*	1.9*	1.2
Oropharynx	1.9*	2.1*	1.1	3.3*	3.5*	2.6*
Hypopharynx	-2.6*	-2.5*	-3.2*	-0.4	0.1	-2.6
Other oral cavity & pharynx	3.1	4.1*	-0.5	-2.5*	-2.4*	-3.3*
Digestive System:	-1.1*	-1.2*	-1.1*	-0.8*	-0.7*	-1.0*
Esophagus	-1.3*	-1.4*	-1.6*	-1.1*	-1.1*	-1.7*
Stomach	-0.9*	-1.2*	-0.7*	-2.1*	-2.4*	-2.0*
Small intestine	1.8*	1.1*	2.5*	0.6	0.8	0.3
Colon & Rectum:	-2.7*	-2.8*	-2.6*	-2.4*	-2.5*	-2.5*
Colon	-3.0*	-3.2*	-2.8*	-	-	-
Rectum	-1.9*	-2.0*	-1.9*	-	-	-
Anus, anal canal & anorectum	1.4*	0.5	2.1*	2.8*	3.5*	2.5*
Liver & intrahepatic bile duct	2.5*	2.3*	2.8*	2.6*	2.5*	2.5*
Gallbladder	0.2	0.7	0.0	-0.8*	-0.6	-0.8
Other biliary	0.2	0.3	-0.2	-0.7	-1.0	-0.5
Pancreas	0.5*	0.5*	0.5*	0.1	0.1	0.0
Retroperitoneum	-1.0	-1.3	-0.7	-0.2	0.2	-0.6
Peritoneum, omentum & mesentery	-4.0*	-4.6	-3.8*	-0.2	2.4	-0.5
Other digestive system	5.0*	5.2*	4.7*	2.6*	2.4*	2.5*
Respiratory System:	-2.4*	-3.0*	-1.8*	-2.6*	-3.2*	-2.0*
Nose, nasal cavity & middle ear	-0.7	-0.6	-0.8	-2.0	-1.4	-3.1*
Larynx	-2.8*	-3.0*	-2.6*	-2.3*	-2.3*	-3.1*
Lung & bronchus	-2.4*	-3.1*	-1.8*	-2.6*	-3.2*	-2.0*
Pleura	0.3	-1.7	-	1.0	0.6	1.2
Trachea & other respiratory organs	-1.4	-0.5	-2.9*	-0.6	-0.7	-0.8
Bones & joints	0.4	0.4	0.5	0.2	0.5	-0.3
Soft tissue (including heart)	0.7*	0.5*	0.9	0.4*	0.6*	0.1
Skin (excl. basal & squamous):	1.5*	1.5*	1.2*	-0.2	-0.2	-0.6*
Melanoma of the skin	1.5*	1.6*	1.3*	-1.0*	-1.1*	-1.1*
Other non-epithelial skin	0.9	1.0	0.3	2.2*	2.2*	1.5
Breast	0.0	-0.1	0.2	-1.8*	-0.5	-1.6*
Breast (<i>in situ</i>)	-0.3	0.2	-0.2	-	-	-

The APC is the Annual Percent Change over the time interval.

Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^a SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

^b US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

* The APC is significantly different from zero (p<.05).

- Statistic could not be calculated. Trend based on less than 10 cases for at least one year within the time interval.

Table 1.8 - continued
SEER Incidence and U.S. Mortality Trends by Primary Cancer Site and Sex
All Races, 2006-2015

Site	Incidence ^a			US Mortality ^b		
	Total APC	Males APC	Females APC	Total APC	Males APC	Females APC
Female Genital System:	0.0	-	0.2	-1.0*	-	-0.7*
Cervix uteri	-1.5*	-	-1.4*	-0.9*	-	-0.7*
Corpus uteri	1.1*	-	1.3*	0.4	-	0.7
Uterus, NOS	4.4*	-	4.7*	2.0*	-	2.4*
Ovary ^c	-2.0*	-	-1.8*	-2.6*	-	-2.4*
Vagina	-1.0	-	-0.8	-0.8*	-	-0.5
Vulva	0.6*	-	0.8*	0.8*	-	1.1*
Other female genital system	6.8*	-	6.8*	5.6*	-	5.8*
Male Genital System:	-5.4*	-5.7*	-	-2.3*	-3.0*	-
Prostate	-5.7*	-6.0*	-	-2.4*	-3.1*	-
Testis	0.8*	0.8*	-	0.4	0.3	-
Penis	1.1	0.9	-	0.9	0.4	-
Other male genital system	1.5	1.0	-	3.2	2.6	-
Urinary System:	-0.5*	-0.7*	-0.4	-0.4*	-0.3*	-1.1*
Urinary bladder	-1.3*	-1.5*	-1.4*	-0.1	-0.3*	-0.6*
Kidney & renal pelvis	0.5*	0.5	0.3	-0.7*	-0.4*	-1.5*
Ureter	-2.0*	-1.9*	-2.5*	0.1	0.5	-0.3
Other urinary system	3.8*	3.5*	4.2*	-0.9	-0.9	-1.5*
Eye & Orbit	0.4	0.4	0.4	1.7*	2.4*	1.0
Brain & Nervous System: ^d	-0.5*	-0.6*	-0.3	0.6*	0.6*	0.6*
Brain	-0.5*	-0.7*	-0.2	-	-	-
Cranial nerves & other nervous system	-0.6	0.7	-1.6	-	-	-
Endocrine System:	3.1*	2.9*	3.2*	-0.1	0.6*	-0.7*
Thyroid	3.3*	3.5*	3.3*	0.0	0.8	-0.6
Other endocrine & thymus	-0.8	-1.4	-0.2	-0.3	0.3	-0.9
Lymphoma:	-0.7*	-0.5*	-0.9*	-2.3*	-2.0*	-2.8*
Hodgkin lymphoma	-1.9*	-1.7*	-2.1*	-3.8*	-3.4*	-4.2*
Non-Hodgkin lymphoma	-0.5*	-0.4*	-0.7*	-2.2*	-2.0*	-2.7*
Myeloma	1.3*	1.2*	1.3*	-0.5	-0.5*	-0.6
Leukemia:	0.3	0.3	0.2	-1.3*	-1.3*	-1.4*
Lymphocytic:	-0.4	-0.4	-0.4	-1.9*	-2.0*	-2.1*
Acute lymphocytic	0.7	0.5*	0.9	-0.7*	-0.6	-1.0
Chronic lymphocytic	-0.7	-0.6	-1.0*	-2.3*	-2.4*	-2.6*
Other lymphocytic	-1.1	-1.6*	0.1	-2.0*	-2.4*	-1.3
Myeloid & Monocytic:	1.6*	1.6*	1.3*	-0.2	-0.2	-0.4
Acute myeloid	2.2*	2.4*	1.9*	-0.2	-0.2	-0.4
Chronic myeloid	1.2*	1.2*	1.0	-1.2*	-1.2	-1.3
Acute monocytic	-4.9*	-4.9*	-5.1*	-1.9	-2.3	-1.9
Other myeloid & monocytic	-1.2	-1.6	-0.7	1.1*	0.9*	1.1
Other leukemia:	-4.5*	-4.6*	-4.7*	-2.6*	-2.5*	-2.8*
Other acute leukemia	-1.4	-2.4	-0.6	-5.0*	-5.0*	-5.2*
Aleukemic, subleukemic & NOS	-6.7*	-6.2*	-7.2*	-1.2*	-1.1	-1.4*
Kaposi Sarcoma ^e	-2.1*	-2.3*	-1.6	-	-	-
Mesothelioma ^e	-1.8*	-2.2*	-1.2	-	-	-
Ill-defined & unspecified	-2.8*	-2.8*	-2.8*	-1.9*	-1.9*	-2.0*

The APC is the Annual Percent Change over the time interval.

Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^a SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

^b US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

^c Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

^d Due to coding changes, Brain & Nervous System mortality are no longer shown separately.

^e Trend not shown for mortality. Category did not exist in mortality coding until 1999.

* The APC is significantly different from zero (p<.05).

- Statistic could not be calculated. Trend based on less than 10 cases for at least one year within the time interval.

Table 1.9
SEER Incidence and U.S. Mortality Trends by Primary Cancer Site and Sex
Whites, 2006-2015

Site	Incidence ^a			US Mortality ^b		
	Total APC	Males APC	Females APC	Total APC	Males APC	Females APC
All Sites	-1.2*	-2.2*	-0.2*	-1.4*	-1.6*	-1.3*
Oral Cavity & Pharynx:	0.9*	1.1*	0.2	0.5	0.9*	-0.6
Lip	0.2	-0.4	1.4	0.3	-1.2	2.4
Tongue	1.9*	2.3*	0.8	0.8	1.3*	-0.4
Salivary gland	-0.3	-0.6	-0.2	1.8*	2.4*	0.1
Floor of mouth	-3.2*	-3.6*	-2.2	-6.3*	-4.6*	-9.5*
Gum & other oral cavity	-0.1	-0.1	-0.1	0.7	1.2	-0.1
Nasopharynx	-2.4*	-2.1*	-2.9*	-2.1*	-1.5*	-3.3*
Tonsil	2.9*	3.1*	2.2*	2.8*	2.9*	1.6
Oropharynx	3.1*	3.5*	1.7	4.0*	4.3*	3.0*
Hypopharynx	-2.6*	-2.2*	-4.2*	-0.4	0.1	-2.6
Other oral cavity & pharynx	4.1*	5.4*	-0.5	-2.2*	-2.0*	-3.3*
Digestive System:	-1.0*	-1.1*	-1.0*	-0.6*	-0.6*	-0.8*
Esophagus	-0.7*	-0.9*	-0.9	-0.5*	-0.6*	-1.2*
Stomach	-0.6*	-0.9*	-0.3	-2.0*	-2.4*	-1.7*
Small intestine	1.6*	1.0	2.4*	0.6	0.8	0.3
Colon & Rectum:	-2.6*	-2.9*	-2.4*	-2.3*	-2.4*	-2.4*
Colon	-2.9*	-3.2*	-2.7*	-	-	-
Rectum	-1.9*	-2.0*	-1.8*	-	-	-
Anus, anal canal & anorectum	1.7*	0.4	2.6*	2.9*	3.2*	2.8*
Liver & intrahepatic bile duct	3.1*	2.8*	3.8*	2.8*	2.7*	2.7*
Gallbladder	-0.2	0.6	-0.5	-1.3*	-1.3	-1.2*
Other biliary	0.2	0.2	-0.1	-0.8*	-1.2*	-0.5
Pancreas	0.6*	0.6*	0.6*	0.2*	0.2*	0.1
Retroperitoneum	-1.5*	-1.4	-1.6	-0.8	-0.6	-1.2
Peritoneum, omentum & mesentery	-4.2*	-4.9	-4.0*	-0.2	2.8*	-0.6
Other digestive system	5.2*	5.4*	4.8*	2.7*	2.5*	2.6*
Respiratory System:	-2.4*	-3.0*	-1.8*	-2.5*	-3.1*	-1.9*
Nose, nasal cavity & middle ear	-0.9	-0.7	-1.2*	-2.0	-1.3	-3.5*
Larynx	-2.7*	-2.9*	-2.5*	-2.0*	-2.0*	-2.8*
Lung & bronchus	-2.4*	-3.1*	-1.7*	-2.5*	-3.2*	-1.9*
Pleura	-0.9	-2.2	-	1.0	0.7	1.1
Trachea & other respiratory organs	-1.0	-0.1	-2.6*	-0.3	-0.3	-0.6
Bones & joints	0.6	0.3	1.1	0.3	0.7	-0.2
Soft tissue (including heart)	0.7	0.3	0.9	0.3*	0.6	0.0
Skin (excl. basal & squamous):	1.5*	1.5*	1.3*	0.0	0.1	-0.4
Melanoma of the skin	1.5*	1.6*	1.3*	-0.8	-0.9*	-0.9*
Other non-epithelial skin	0.9	0.9	0.6	2.6*	2.6*	1.6*
Breast	-0.1	0.1	0.0	-1.8*	-0.8	-1.6*
Breast (<i>in situ</i>)	-0.8	-0.4	-0.6	-	-	-

The APC is the Annual Percent Change over the time interval.

Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^a SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

^b US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

* The APC is significantly different from zero (p<.05).

- Statistic could not be calculated. Trend based on less than 10 cases for at least one year within the time interval.

Table 1.9 - continued
SEER Incidence and U.S. Mortality Trends by Primary Cancer Site and Sex
Whites, 2006-2015

Site	Incidence ^a			US Mortality ^b		
	Total APC	Males APC	Females APC	Total APC	Males APC	Females APC
Female Genital System:	-0.1	-	0.1	-1.0*	-	-0.8*
Cervix uteri	-1.3*	-	-1.2*	-0.3	-	-0.2
Corpus uteri	0.9*	-	1.1*	0.3	-	0.6
Uterus, NOS	4.7*	-	5.1*	1.8*	-	2.2*
Ovary ^c	-2.1*	-	-1.9*	-2.7*	-	-2.5*
Vagina	-1.1	-	-0.8	-0.5	-	-0.1
Vulva	0.9*	-	1.1*	0.9*	-	1.3*
Other female genital system	7.0*	-	7.1*	6.1*	-	6.4*
Male Genital System:	-5.8*	-6.1*	-	-2.1*	-2.8*	-
Prostate	-6.2*	-6.5*	-	-2.2*	-2.9*	-
Testis	0.8*	0.7*	-	0.7	0.6	-
Penis	1.2	0.9	-	1.2	0.8	-
Other male genital system	1.3	0.7	-	2.6	2.1	-
Urinary System:	-0.5*	-0.7*	-0.5	-0.2*	-0.2*	-0.9*
Urinary bladder	-1.3*	-1.5*	-1.4*	0.0	-0.2	-0.3
Kidney & renal pelvis	0.5*	0.5	0.4	-0.5*	-0.3*	-1.4*
Ureter	-1.7*	-1.8*	-2.0	0.0	0.5	-0.4
Other urinary system	3.8*	3.2*	4.7*	-0.2	-0.6	-0.2
Eye & Orbit	0.7	0.7	0.5	2.0*	2.8*	1.2
Brain & Nervous System: ^d	-0.4	-0.5*	-0.2	0.6*	0.6*	0.6*
Brain	-0.4	-0.6	-0.1	-	-	-
Cranial nerves & other nervous system	-0.5	0.4	-1.4	-	-	-
Endocrine System:	3.0*	2.9*	3.1*	-0.1	0.7*	-0.7*
Thyroid	3.2*	3.4*	3.2*	0.1	0.9*	-0.6
Other endocrine & thymus	-0.9	-1.4	-0.4	-0.3	0.4	-1.0*
Lymphoma:	-0.8*	-0.7*	-1.0*	-2.3*	-2.0*	-2.8*
Hodgkin lymphoma	-2.1*	-2.0*	-2.2*	-3.8*	-3.3*	-4.4*
Non-Hodgkin lymphoma	-0.7*	-0.5*	-0.8*	-2.2*	-1.9*	-2.7*
Myeloma	1.3*	1.2*	1.2*	-0.4	-0.4	-0.7*
Leukemia:	0.2	0.2	0.1	-1.1*	-1.1*	-1.3*
Lymphocytic:	-0.6	-0.6	-0.7	-1.7*	-1.7*	-2.1*
Acute lymphocytic	0.7	0.8*	0.5	-0.9*	-0.5	-1.5*
Chronic lymphocytic	-1.0*	-1.0*	-1.3*	-2.0*	-2.0*	-2.4*
Other lymphocytic	-1.1	-1.5*	-0.5	-1.8*	-2.0*	-1.3
Myeloid & Monocytic:	1.6*	1.6*	1.4*	-0.1	-0.2	-0.3
Acute myeloid	2.3*	2.4*	2.1*	-0.2	-0.2	-0.3
Chronic myeloid	1.2*	1.2*	1.0	-0.6	-0.6	-0.9
Acute monocytic	-5.4*	-5.3*	-5.8*	-1.9	-2.6	-1.6
Other myeloid & monocytic	-1.3	-1.5	-1.5	1.2*	1.0*	0.9
Other leukemia:	-4.3*	-4.3*	-4.6*	-2.5*	-2.4*	-2.8*
Other acute leukemia	-1.1	-2.1	-0.3	-4.9*	-4.9*	-5.0*
Aleukemic, subleukemic & NOS	-6.6*	-6.0*	-7.5*	-1.1	-1.0	-1.4*
Kaposi Sarcoma ^e	-3.9*	-4.2*	-2.1	-	-	-
Mesothelioma ^e	-1.5*	-1.9*	-0.8	-	-	-
Ill-defined & unspecified	-2.7*	-2.6*	-2.7*	-1.7*	-1.7*	-1.7*

The APC is the Annual Percent Change over the time interval.

Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^a SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

^b US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

^c Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

^d Due to coding changes, Brain & Nervous System mortality are no longer shown separately.

^e Trend not shown for mortality. Category did not exist in mortality coding until 1999.

* The APC is significantly different from zero (p<.05).

- Statistic could not be calculated. Trend based on less than 10 cases for at least one year within the time interval.

Table 1.10
SEER Incidence and U.S. Mortality Trends by Primary Cancer Site and Sex
Blacks, 2006-2015

Site	Incidence ^a			US Mortality ^b		
	Total APC	Males APC	Females APC	Total APC	Males APC	Females APC
All Sites	-1.7*	-3.1*	-0.4*	-2.1*	-2.8*	-1.6*
Oral Cavity & Pharynx:	-1.6*	-1.8*	-1.3	-2.0*	-2.3*	-1.4
Lip	-	-	-	-	-	-
Tongue	-1.5*	-1.7	-1.7	-1.7	-1.4*	-2.4
Salivary gland	0.4	0.7	0.1	1.7	1.1	2.2
Floor of mouth	-3.0	-2.3	-	-	-	-
Gum & other oral cavity	-2.3*	-1.4	-3.8*	-1.8	-1.9	-1.2
Nasopharynx	-0.7	-1.8	1.6	-1.9	-1.9	-2.3
Tonsil	-0.5	-0.6	0.3	-3.8*	-4.2*	-2.0
Oropharynx	-3.1*	-3.7*	-	0.1	-0.8	2.2
Hypopharynx	-3.8*	-5.1*	3.1	-0.4	0.9	-
Other oral cavity & pharynx	-0.2	-0.4	-	-3.9*	-4.5*	-2.5
Digestive System:	-1.9*	-1.9*	-2.0*	-1.6*	-1.7*	-1.8*
Esophagus	-4.7*	-5.3*	-4.1*	-4.7*	-5.3*	-3.9*
Stomach	-2.8*	-3.3*	-2.6*	-3.1*	-3.0*	-3.5*
Small intestine	2.3*	1.3	3.4*	0.4	0.9	-0.1
Colon & Rectum:	-3.3*	-3.3*	-3.4*	-3.1*	-3.2*	-3.3*
Colon	-3.5*	-3.6*	-3.6*	-	-	-
Rectum	-2.7*	-2.5*	-3.0*	-	-	-
Anus, anal canal & anorectum	0.8	1.1	0.1	3.2*	5.6*	0.5
Liver & intrahepatic bile duct	2.5*	2.6*	2.2*	2.2*	2.2*	2.2*
Gallbladder	2.0*	0.7	2.4	1.3	1.9	1.0
Other biliary	-0.6	-0.5	-0.6	0.6	1.4	0.3
Pancreas	-0.5	-0.4	-0.4	-0.5*	-0.3	-0.6*
Retroperitoneum	0.1	-	0.0	4.0	-	-
Peritoneum, omentum & mesentery	-0.6	-	0.4	3.5	-	4.1*
Other digestive system	3.6*	2.2	4.6*	2.4	2.5	2.5
Respiratory System:	-2.6*	-3.4*	-1.8*	-3.0*	-3.7*	-2.2*
Nose, nasal cavity & middle ear	-1.4	-1.8	-0.4	-2.3	-2.8	-1.1
Larynx	-2.9*	-3.2*	-2.0	-3.6*	-3.5*	-4.1*
Lung & bronchus	-2.6*	-3.4*	-1.8*	-3.0*	-3.7*	-2.2*
Pleura	-	-	-	-	-	-
Trachea & other respiratory organs	-4.2	-	-	-0.1	-	-
Bones & joints	-1.0	-1.1	-0.6	-0.1	-0.5	0.2
Soft tissue (including heart)	0.5	0.7	0.4	1.1	1.1	1.2*
Skin (excl. basal & squamous):	-0.7	-0.1	-1.4	-1.7*	-1.8*	-1.6
Melanoma of the skin	-0.1	0.7	-0.5	-2.3*	-1.9	-2.9
Other non-epithelial skin	-1.3	-0.8	-2.2	-1.1	-1.8	0.5
Breast	0.3	-1.0	0.4	-1.7*	1.1	-1.5*
Breast (<i>in situ</i>)	1.1*	-	1.3*	-	-	-

The APC is the Annual Percent Change over the time interval.

Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^a SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

^b US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

* The APC is significantly different from zero (p<.05).

- Statistic could not be calculated. Trend based on less than 10 cases for at least one year within the time interval.

Table 1.10 - continued
SEER Incidence and U.S. Mortality Trends by Primary Cancer Site and Sex
Blacks, 2006-2015

Site	Incidence ^a			US Mortality ^b		
	Total APC	Males APC	Females APC	Total APC	Males APC	Females APC
Female Genital System:	0.3	-	0.5	-0.3	-	0.0
Cervix uteri	-2.8*	-	-2.6*	-2.8*	-	-2.5*
Corpus uteri	2.0*	-	2.2*	0.8	-	1.0
Uterus, NOS	3.6	-	3.9	2.5*	-	2.8*
Ovary ^c	-1.7*	-	-1.4*	-1.4*	-	-1.2*
Vagina	0.4	-	0.3	-2.3	-	-2.0
Vulva	0.0	-	0.1	0.7	-	1.2
Other female genital system	5.1*	-	5.1*	3.2*	-	3.1*
Male Genital System:	-5.1*	-5.4*	-	-3.8*	-4.4*	-
Prostate	-5.2*	-5.5*	-	-3.8*	-4.5*	-
Testis	2.0	1.9	-	0.1	0.5	-
Penis	2.0	1.3	-	-1.1	-1.1	-
Other male genital system	-	-	-	-	-	-
Urinary System:	0.1	0.0	-0.1	-1.1*	-0.5	-2.0*
Urinary bladder	-0.8*	-0.8	-1.2*	-0.9	-0.4	-2.0*
Kidney & renal pelvis	0.7	0.7	0.6	-1.1*	-0.7	-1.8*
Ureter	-5.5*	-	-	-1.8	-	-
Other urinary system	2.6	1.6	-	-3.2	-	-6.2*
Eye & Orbit	2.0	-	-	-	-	-
Brain & Nervous System: ^d	-0.7	-0.5	-1.1	1.3*	1.4	1.2
Brain	-0.6	-0.9	-0.7	-	-	-
Cranial nerves & other nervous system	-1.0	3.7	-4.6	-	-	-
Endocrine System:	3.6*	1.4	4.2*	0.5	0.6	0.2
Thyroid	4.2*	3.0*	4.6*	0.7	1.0	0.3
Other endocrine & thymus	-2.1	-4.3	-0.2	0.2	0.2	0.1
Lymphoma:	-0.3	-0.2	-0.4	-1.8*	-2.0*	-1.5*
Hodgkin lymphoma	-0.9	-0.7	-1.1	-3.5*	-4.1*	-2.8*
Non-Hodgkin lymphoma	-0.2	-0.1	-0.2	-1.7*	-1.8*	-1.5*
Myeloma	0.8	0.4	0.9	-0.4	-1.0*	0.0
Leukemia:	0.9	0.5	1.3*	-1.9*	-2.3*	-1.6*
Lymphocytic:	0.6	-0.1	1.2	-3.4*	-4.5*	-2.1*
Acute lymphocytic	-0.7	-3.5*	3.2	0.0	-1.1	2.1
Chronic lymphocytic	0.9	0.8	0.5	-4.2*	-4.9*	-3.5*
Other lymphocytic	-0.3	-	-	-4.5*	-	-
Myeloid & Monocytic:	2.2*	2.2*	2.1*	-0.5	-0.5	-0.8
Acute myeloid	2.9*	3.7*	2.1	-0.2	0.0	-0.7
Chronic myeloid	1.1	-0.1	2.0	-3.7*	-4.6*	-2.9
Acute monocytic	-1.5	-	-	-	-	-
Other myeloid & monocytic	-	-	-	2.2	1.2	2.7
Other leukemia:	-6.8*	-8.7*	-5.2	-2.6*	-2.9*	-2.6*
Other acute leukemia	-1.4	-	-	-5.9*	-5.9*	-5.9*
Aleukemic, subleukemic & NOS	-10.0*	-	-8.8*	-1.3*	-1.7	-1.2
Kaposi Sarcoma ^e	-0.9	-0.9	-	-	-	-
Mesothelioma ^e	-3.6	-4.7	-	-	-	-
Ill-defined & unspecified	-3.1*	-3.1*	-3.1*	-2.9*	-3.1*	-2.8*

The APC is the Annual Percent Change over the time interval.

Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^a SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

^b US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

^c Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

^d Due to coding changes, Brain & Nervous System mortality are no longer shown separately.

^e Trend not shown for mortality. Category did not exist in mortality coding until 1999.

* The APC is significantly different from zero (p<.05).

- Statistic could not be calculated. Trend based on less than 10 cases for at least one year within the time interval.

Table 1.11

Age Distribution (%) of Incidence Cases by Site, 2011-2015

All Races, Both Sexes

Site	Age at Diagnosis								All Ages	Cases
	<20	20-34	35-44	45-54	55-64	65-74	75-84	85+		
All Sites	1.0	2.8	5.0	13.3	24.6	27.2	18.3	7.9	100.0%	2,087,262
Oral Cavity & Pharynx:	0.5	1.9	4.5	17.5	30.8	24.5	14.0	6.3	100.0%	55,070
Lip	0.1	1.4	3.9	13.9	21.2	24.0	22.2	13.2	100.0%	3,241
Tongue	0.1	1.6	3.8	17.0	33.8	26.1	12.8	4.7	100.0%	16,883
Salivary gland	1.9	6.0	7.4	12.7	19.0	23.2	18.5	11.3	100.0%	6,048
Floor of mouth	0.1	0.4	2.3	18.1	35.3	25.6	13.4	4.8	100.0%	2,438
Gum & other oral cavity	0.6	1.7	4.0	11.9	22.9	25.0	21.3	12.5	100.0%	7,179
Nasopharynx	2.8	6.3	12.4	22.8	26.6	18.4	8.5	2.3	100.0%	2,904
Tonsil	0.0	0.4	4.3	25.4	40.3	21.3	6.7	1.5	100.0%	10,093
Oropharynx	0.2	0.3	2.8	18.2	34.0	28.3	12.2	4.0	100.0%	2,198
Hypopharynx	0.0	0.2	1.1	14.5	33.1	30.4	15.8	4.9	100.0%	2,814
Other oral cavity & pharynx	0.3	0.9	2.7	16.8	31.0	26.3	14.9	7.1	100.0%	1,272
Digestive System:	0.2	1.2	3.5	13.0	24.5	25.9	20.9	10.7	100.0%	386,637
Esophagus	0.0	0.4	1.9	10.4	27.1	30.6	21.2	8.5	100.0%	20,146
Stomach	0.1	1.7	4.3	11.9	21.5	26.3	22.7	11.4	100.0%	33,965
Small intestine	0.1	1.7	4.8	14.7	24.9	27.4	19.0	7.3	100.0%	10,894
Colon & Rectum:	0.2	1.6	4.4	15.0	22.6	24.3	20.6	11.4	100.0%	186,532
Colon	0.2	1.5	3.8	12.6	20.6	25.0	23.0	13.3	100.0%	130,439
Rectum	0.1	1.9	5.8	20.6	27.0	22.7	15.0	7.1	100.0%	56,093
Colon & Rectum (Male)	0.1	1.5	4.3	15.7	24.9	25.9	19.1	8.4	100.0%	97,424
Colon & Rectum (Female)	0.2	1.7	4.4	14.3	20.0	22.7	22.2	14.7	100.0%	89,108
Anus, anal canal & anorectum	0.0	1.2	5.3	21.7	30.2	22.4	13.2	6.0	100.0%	8,965
Liver & intrahepatic bile duct	0.8	0.7	1.8	12.1	37.1	25.8	15.9	5.9	100.0%	44,172
Gallbladder	0.0	0.5	2.4	9.3	20.5	27.8	25.7	13.7	100.0%	5,539
Other biliary	0.1	0.6	2.5	8.2	20.7	27.3	26.6	14.0	100.0%	8,891
Pancreas	0.1	0.5	1.9	8.6	22.4	28.4	24.7	13.4	100.0%	59,942
Retroperitoneum	7.5	4.5	5.1	16.4	23.0	23.8	14.8	4.9	100.0%	1,767
Peritoneum, omentum & mesentery	0.3	1.2	1.9	10.9	24.9	32.7	21.0	7.2	100.0%	2,543
Other digestive system	0.2	0.8	2.7	9.3	21.6	26.0	25.5	14.0	100.0%	3,281
Respiratory System:	0.1	0.3	1.2	7.8	22.2	33.0	26.1	9.2	100.0%	275,950
Nose, nasal cavity & middle ear	1.7	3.4	7.4	15.0	23.5	22.6	18.1	8.5	100.0%	3,204
Larynx	0.0	0.4	2.1	13.5	31.4	30.0	17.1	5.6	100.0%	14,536
Lung & bronchus	0.0	0.2	1.0	7.4	21.7	33.4	26.8	9.4	100.0%	257,301
Lung & bronchus (Male)	0.0	0.2	1.0	7.0	22.7	34.0	26.4	8.6	100.0%	133,651
Lung & bronchus (Female)	0.0	0.3	1.1	7.7	20.6	32.7	27.3	10.3	100.0%	123,650
Pleura	1.4	2.2	4.3	5.1	15.9	21.7	31.9	17.4	100.0%	138
Trachea & other respiratory organs	17.1	20.0	8.0	10.0	14.5	14.8	10.5	5.1	100.0%	771
Bones & joints	27.3	15.4	9.4	12.1	12.8	11.7	7.6	3.7	100.0%	4,207
Soft tissue (including heart)	8.3	9.0	8.5	14.2	19.5	18.3	14.6	7.5	100.0%	15,839
Skin (excl. basal & squamous):	0.5	5.4	7.4	14.6	21.8	23.4	17.7	9.1	100.0%	116,628
Melanoma of the skin	0.4	5.5	7.6	15.1	22.4	23.5	17.1	8.3	100.0%	107,109
Other non-epithelial skin	1.3	4.9	5.2	8.8	14.8	22.1	25.3	17.8	100.0%	9,519
Breast (Female)	0.0	1.9	8.6	20.4	25.9	24.1	13.6	5.5	100.0%	317,468
Breast (Female -in situ)	0.0	0.7	9.5	26.4	27.9	23.7	9.8	1.9	100.0%	77,965

Source: SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG). Percents may not sum to 100 due to rounding.

Table 1.11 - continued

Age Distribution (%) of Incidence Cases by Site, 2011-2015

All Races, Both Sexes

Site	Age at Diagnosis								All Ages	Cases
	<20	20-34	35-44	45-54	55-64	65-74	75-84	85+		
Female Genital System:	0.4	4.0	8.1	17.5	29.1	23.3	12.3	5.4	100.0%	125,897
Cervix uteri	0.1	14.0	23.6	23.4	19.1	11.4	5.9	2.5	100.0%	16,935
Corpus uteri	0.0	1.7	5.4	16.3	35.0	26.8	11.2	3.7	100.0%	66,066
Uterus, NOS	0.2	1.9	5.5	16.1	24.6	22.7	16.3	12.7	100.0%	2,147
Ovary ^a	1.3	4.0	6.8	18.1	24.6	22.1	15.3	7.7	100.0%	29,281
Vagina	1.0	1.6	4.7	13.0	23.1	24.4	19.6	12.6	100.0%	1,868
Vulva	0.1	1.7	5.2	14.5	20.6	22.0	21.1	14.8	100.0%	6,374
Other female genital system	0.5	4.5	3.9	13.4	25.5	27.2	17.4	7.6	100.0%	3,226
Male Genital System:	0.3	2.3	1.6	9.2	31.3	37.0	14.5	3.8	100.0%	272,609
Prostate	0.0	0.0	0.5	9.0	32.7	38.8	15.1	3.9	100.0%	257,650
Testis	5.7	50.5	23.1	12.9	5.5	1.7	0.5	0.2	100.0%	12,481
Penis	0.0	1.7	4.9	12.2	21.0	26.3	23.3	10.7	100.0%	1,873
Other male genital system	1.5	3.3	4.3	12.6	22.8	27.4	20.2	7.9	100.0%	605
Urinary System:	0.5	1.1	3.4	10.2	21.9	28.7	23.2	10.9	100.0%	171,139
Urinary bladder	0.1	0.4	1.3	6.1	18.3	29.8	28.8	15.1	100.0%	91,295
Kidney & renal pelvis	1.1	1.9	6.0	15.5	26.6	27.3	16.0	5.6	100.0%	75,678
Ureter	0.0	0.1	0.4	3.6	15.0	30.7	34.2	15.9	100.0%	2,433
Other urinary system	0.1	0.2	1.2	5.9	17.5	27.7	28.9	18.5	100.0%	1,733
Eye & Orbit	11.8	3.8	5.9	13.3	21.3	22.8	15.1	6.0	100.0%	4,027
Brain & Nervous System:	12.5	8.9	7.6	13.7	20.3	19.2	12.9	4.9	100.0%	29,330
Brain	11.6	8.8	7.4	13.6	20.7	19.6	13.2	5.0	100.0%	27,575
Cranial nerves & other nervous system	27.2	10.3	10.7	14.7	13.8	12.9	6.8	3.6	100.0%	1,755
Endocrine System:	2.8	14.9	18.1	22.3	20.9	13.9	5.6	1.4	100.0%	69,278
Thyroid	1.9	15.4	18.6	22.8	21.0	13.7	5.4	1.3	100.0%	65,899
Other endocrine & thymus	20.9	7.0	8.0	12.7	20.2	18.0	10.0	3.2	100.0%	3,379
Lymphoma:	2.9	6.7	6.2	12.1	20.2	23.8	19.4	8.7	100.0%	101,968
Hodgkin lymphoma	12.2	31.2	13.9	13.0	11.8	9.3	6.3	2.3	100.0%	11,324
Non-Hodgkin lymphoma	1.7	3.6	5.3	12.0	21.3	25.6	21.1	9.5	100.0%	90,644
Myeloma	0.0	0.5	2.8	10.9	23.2	29.8	23.7	9.0	100.0%	31,483
Leukemia:	8.8	4.6	4.7	9.8	17.9	23.2	20.5	10.6	100.0%	63,616
Lymphocytic:	13.2	2.8	3.2	9.2	19.2	23.6	19.1	9.8	100.0%	31,743
Acute lymphocytic	55.4	10.4	6.2	7.2	8.5	6.6	4.2	1.5	100.0%	7,535
Chronic lymphocytic	0.0	0.3	1.6	9.1	22.4	29.6	24.5	12.6	100.0%	22,377
Other lymphocytic	1.1	1.9	10.0	18.5	22.9	21.1	15.1	9.3	100.0%	1,831
Myeloid & Monocytic:	4.1	6.5	6.3	10.6	17.1	23.3	21.7	10.3	100.0%	29,539
Acute myeloid	4.9	6.0	5.3	9.5	16.9	24.3	22.6	10.5	100.0%	19,570
Chronic myeloid	2.1	7.7	8.6	13.5	17.9	21.4	19.3	9.5	100.0%	8,371
Acute monocytic	7.6	4.7	7.0	10.0	16.3	21.6	22.6	10.2	100.0%	898
Other myeloid & monocytic	3.6	7.4	5.9	8.7	14.4	21.4	22.7	15.9	100.0%	700
Other leukemia:	6.1	4.9	4.4	7.5	11.7	16.2	24.2	24.9	100.0%	2,334
Other acute leukemia	8.0	5.7	4.5	6.9	11.5	16.7	25.1	21.6	100.0%	1,031
Aleukemic, subleukemic & NOS	4.7	4.3	4.3	8.0	11.9	15.9	23.4	27.6	100.0%	1,303
Kaposi Sarcoma	0.3	24.8	19.7	21.7	10.7	8.1	8.8	6.1	100.0%	2,202
Mesothelioma	0.0	1.1	1.6	5.8	14.8	28.8	31.7	16.1	100.0%	4,233
Ill-defined & unspecified	0.4	0.9	2.2	8.0	18.9	23.8	25.7	20.0	100.0%	37,177

Source: SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG). Percents may not sum to 100 due to rounding.

^a Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

Table 1.12
 Median Age of Cancer Patients at Diagnosis^a, 2011-2015
 By Primary Cancer Site, Race and Sex

Site	All Races			Whites			Blacks		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
All Sites	66.0	66.0	65.0	66.0	67.0	65.0	63.0	63.0	62.0
Oral Cavity & Pharynx:	63.0	62.0	65.0	63.0	63.0	66.0	60.0	60.0	59.0
Lip	68.0	68.0	70.0	69.0	68.0	70.0	60.0	61.0	-
Tongue	63.0	62.0	64.0	63.0	62.0	65.0	60.0	60.0	59.0
Salivary gland	66.0	68.0	62.0	67.0	70.0	64.0	57.0	58.0	56.0
Floor of mouth	63.0	62.0	65.0	64.0	63.0	65.0	60.0	61.0	59.0
Gum & other oral cavity	68.0	66.0	71.0	69.0	67.0	73.0	62.0	61.0	63.0
Nasopharynx	57.0	57.0	56.0	60.0	60.0	59.0	55.0	56.0	53.0
Tonsil	59.0	59.0	62.0	59.0	59.0	62.0	59.0	59.0	59.0
Oropharynx	63.0	63.0	64.0	63.0	63.0	65.0	61.0	61.0	60.0
Hypopharynx	65.0	65.0	65.0	65.0	65.0	67.0	61.0	61.0	61.0
Other oral cavity & pharynx	64.0	63.0	67.0	64.0	63.0	68.0	61.0	60.0	62.0
Digestive System:	67.0	66.0	69.0	68.0	67.0	70.0	64.0	63.0	65.0
Esophagus	68.0	67.0	71.0	68.0	67.0	72.0	64.0	63.0	65.0
Stomach	68.0	68.0	69.0	69.0	68.0	70.0	66.0	65.0	68.0
Small intestine	66.0	65.0	66.0	66.0	66.0	67.0	64.0	63.0	64.0
Colon & Rectum:	67.0	66.0	69.0	68.0	66.0	70.0	64.0	63.0	64.0
Colon	69.0	67.0	71.0	70.0	68.0	72.0	65.0	64.0	65.0
Rectum	63.0	62.0	63.0	63.0	63.0	64.0	60.0	60.0	60.0
Anus, anal canal & anorectum	62.0	59.0	63.0	62.0	61.0	63.0	56.0	52.0	60.0
Liver & intrahepatic bile duct	64.0	63.0	68.0	64.0	63.0	68.0	62.0	62.0	63.0
Gallbladder	71.0	71.0	71.0	72.0	71.0	72.0	66.0	69.0	65.0
Other biliary	71.0	70.0	72.0	71.0	70.0	73.0	68.0	68.0	68.0
Pancreas	70.0	69.0	72.0	71.0	69.0	73.0	66.0	65.0	69.0
Retroperitoneum	62.0	63.0	61.0	63.0	63.0	62.0	58.0	58.0	59.0
Peritoneum, omentum & mesentery	68.0	65.5	68.0	68.0	66.0	68.0	64.0	59.5	64.0
Other digestive system	70.0	70.0	71.0	71.0	70.0	72.0	66.0	65.0	66.0
Respiratory System:	70.0	70.0	71.0	71.0	70.0	71.0	66.0	66.0	67.0
Nose, nasal cavity & middle ear	64.0	64.0	65.0	65.0	64.0	67.0	57.0	56.5	61.5
Larynx	65.0	65.0	64.0	66.0	66.0	65.0	63.0	63.0	62.0
Lung & bronchus	70.0	70.0	71.0	71.0	71.0	71.0	67.0	66.0	67.0
Pleura	74.0	77.0	72.0	77.0	79.0	73.0	-	-	-
Trachea & other respiratory organs	49.0	44.0	58.0	50.0	45.0	59.5	52.0	52.0	56.5
Bones & joints	42.0	42.0	43.0	45.0	44.0	46.0	33.0	30.5	35.0
Soft tissue (including heart)	60.0	61.0	59.0	61.0	62.0	60.0	53.0	52.0	54.0
Skin (excl. basal & squamous):	65.0	67.0	60.0	65.0	67.0	61.0	58.0	58.0	59.0
Melanoma of the skin	64.0	66.0	60.0	64.0	66.0	60.0	65.0	65.0	65.5
Other non-epithelial skin	71.0	72.0	70.0	73.0	74.0	72.0	53.0	50.0	54.0
Breast	62.0	68.0	62.0	63.0	68.0	63.0	59.0	64.0	59.0
Breast (<i>in situ</i>)	59.0	61.0	59.0	60.0	61.0	60.0	60.0	61.5	60.0

^a SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).
 - Statistic could not be calculated. Less than 16 cases were diagnosed during the time interval.

Table 1.12 - continued
 Median Age of Cancer Patients at Diagnosis^a, 2011-2015
 By Primary Cancer Site, Race and Sex

Site	All Races			Whites			Blacks		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
Female Genital System:	61.0	-	61.0	62.0	-	62.0	61.0	-	61.0
Cervix uteri	50.0	-	50.0	49.0	-	49.0	52.0	-	52.0
Corpus uteri	62.0	-	62.0	63.0	-	63.0	63.0	-	63.0
Uterus, NOS	65.0	-	65.0	66.0	-	66.0	64.0	-	64.0
Ovary ^b	63.0	-	63.0	64.0	-	64.0	61.0	-	61.0
Vagina	67.0	-	67.0	67.0	-	67.0	65.0	-	65.0
Vulva	68.0	-	68.0	69.0	-	69.0	59.0	-	59.0
Other female genital system	65.0	-	65.0	66.0	-	66.0	60.0	-	60.0
Male Genital System:	66.0	66.0	-	66.0	66.0	-	63.0	63.0	-
Prostate	66.0	66.0	-	66.0	66.0	-	64.0	64.0	-
Testis	33.0	33.0	-	33.0	33.0	-	36.0	36.0	-
Penis	68.0	68.0	-	69.0	69.0	-	65.0	65.0	-
Other male genital system	67.0	67.0	-	67.0	67.0	-	57.0	57.0	-
Urinary System:	69.0	69.0	69.0	69.0	70.0	69.0	64.0	64.0	65.0
Urinary bladder	72.0	72.0	73.0	73.0	73.0	73.0	70.0	69.0	72.0
Kidney & renal pelvis	64.0	64.0	65.0	65.0	64.0	65.0	61.0	61.0	63.0
Ureter	75.0	74.0	75.0	75.0	74.0	76.0	70.0	69.5	72.0
Other urinary system	74.0	74.0	73.0	74.0	75.0	74.0	67.0	67.0	65.0
Eye & Orbit	62.0	62.0	61.0	62.0	63.0	62.0	34.0	42.0	11.0
Brain & Nervous System:	58.0	58.0	59.0	59.0	59.0	60.0	52.0	52.0	52.0
Brain	59.0	58.0	59.0	60.0	59.0	60.0	52.0	53.0	52.0
Cranial nerves & other nervous system	46.0	44.0	48.0	46.0	44.0	48.0	47.0	45.0	51.0
Endocrine System:	51.0	55.0	50.0	51.0	55.0	50.0	52.0	55.0	51.0
Thyroid	51.0	55.0	50.0	51.0	55.0	50.0	52.0	56.0	51.0
Other endocrine & thymus	55.0	54.0	57.0	56.0	54.0	58.0	51.0	49.0	54.0
Lymphoma:	65.0	64.0	66.0	66.0	65.0	67.0	57.0	55.0	59.0
Hodgkin lymphoma	39.0	41.0	37.0	40.0	42.0	37.0	38.0	39.0	37.0
Non-Hodgkin lymphoma	67.0	66.0	68.0	68.0	67.0	69.0	59.0	58.0	61.0
Myeloma	69.0	68.0	69.0	70.0	69.0	70.0	66.0	65.0	66.0
Leukemia:	66.0	66.0	67.0	67.0	67.0	68.0	62.0	62.0	62.0
Lymphocytic:	65.0	65.0	66.0	66.0	66.0	67.0	63.0	63.0	64.0
Acute lymphocytic	15.0	16.0	14.0	16.0	17.0	14.0	16.0	15.0	19.0
Chronic lymphocytic	70.0	69.0	72.0	70.0	69.0	72.0	67.0	66.5	68.0
Other lymphocytic	63.0	61.0	65.0	63.0	62.0	67.0	61.5	59.0	66.0
Myeloid & Monocytic:	67.0	67.0	67.0	68.0	68.0	68.0	60.0	60.0	60.0
Acute myeloid	68.0	68.0	67.0	69.0	69.0	68.0	62.0	62.0	61.0
Chronic myeloid	65.0	65.0	65.0	66.0	66.0	66.0	57.0	56.0	58.0
Acute monocytic	67.0	68.0	65.5	67.0	67.0	66.0	64.0	64.5	62.0
Other myeloid & monocytic	70.0	69.0	70.0	71.0	70.5	71.0	62.5	60.0	63.0
Other leukemia:	74.0	71.0	77.0	76.0	73.0	78.0	64.0	63.0	64.5
Other acute leukemia	73.0	69.0	77.0	74.0	69.0	77.0	66.0	66.0	66.0
Aleukemic, subleukemic & NOS	75.0	73.0	77.0	77.0	76.0	80.0	63.0	62.0	64.0
Kaposi Sarcoma	47.0	45.0	77.0	51.0	49.0	80.0	35.0	34.0	43.0
Mesothelioma	74.0	75.0	72.0	74.0	75.0	72.0	68.5	70.0	66.0
Ill-defined & unspecified	73.0	71.0	75.0	73.0	71.0	76.0	68.0	66.0	70.0

^a SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

^b Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

- Statistic could not be calculated. Less than 16 cases were diagnosed during the time interval.

Table 1.13

Age Distribution (%) of Deaths by Site, 2011-2015

All Races, Both Sexes

Site	Age at Death								All Ages	Deaths
	<20	20-34	35-44	45-54	55-64	65-74	75-84	85+		
All Sites	0.3	0.8	1.9	7.9	19.5	26.5	26.3	16.9	100.0%	2,931,769
Oral Cavity & Pharynx:	0.1	0.8	2.1	12.1	26.9	25.7	19.8	12.6	100.0%	45,589
Lip	0.3	0.3	1.1	7.1	15.8	20.9	26.6	28.0	100.0%	368
Tongue	0.0	1.1	2.6	12.2	27.8	26.2	18.8	11.2	100.0%	11,406
Salivary gland	0.1	1.1	2.5	7.9	17.1	22.9	25.5	23.0	100.0%	4,521
Floor of mouth	0.0	0.3	1.0	15.8	30.3	23.8	19.0	10.0	100.0%	400
Gum & other oral cavity	0.0	0.5	1.5	8.6	20.5	23.4	23.5	22.0	100.0%	6,563
Nasopharynx	0.6	2.9	5.5	17.1	28.2	22.9	15.8	7.0	100.0%	3,320
Tonsil	0.0	0.2	2.2	16.9	35.3	26.9	13.5	5.0	100.0%	4,455
Oropharynx	0.0	0.4	1.5	13.2	30.5	26.6	18.9	8.9	100.0%	4,803
Hypopharynx	0.0	0.4	1.2	13.3	28.8	28.5	18.8	9.1	100.0%	1,709
Other oral cavity & pharynx	0.0	0.1	1.1	11.3	28.7	28.3	20.4	10.0	100.0%	8,044
Digestive System:	0.0	0.5	2.0	8.8	21.7	25.7	24.7	16.6	100.0%	746,822
Esophagus	0.0	0.3	1.6	9.2	25.3	29.5	23.0	11.1	100.0%	73,930
Stomach	0.0	1.3	3.7	10.1	18.7	23.6	25.3	17.3	100.0%	56,128
Small intestine	0.0	0.7	2.6	8.4	19.0	26.8	25.8	16.6	100.0%	6,595
Colon & Rectum:	0.0	0.7	2.6	9.4	18.6	22.8	24.8	21.0	100.0%	259,159
Colon & Rectum (Male)	0.0	0.8	2.7	10.3	21.2	25.2	24.2	15.5	100.0%	135,542
Colon & Rectum (Female)	0.0	0.7	2.4	8.5	15.8	20.2	25.4	27.0	100.0%	123,617
Anus, anal canal & anorectum	0.0	0.7	4.5	16.1	26.7	23.6	17.4	11.0	100.0%	4,588
Liver & intrahepatic bile duct	0.2	0.5	1.4	9.6	31.2	25.9	20.9	10.3	100.0%	119,070
Gallbladder	0.0	0.2	1.4	6.9	17.8	27.7	28.5	17.5	100.0%	10,765
Other biliary	0.0	0.3	1.3	6.0	16.9	25.1	28.6	21.8	100.0%	7,765
Pancreas	0.0	0.2	1.1	7.0	20.1	28.2	27.1	16.3	100.0%	197,170
Retroperitoneum	0.5	1.1	2.6	9.4	21.3	26.2	24.7	14.2	100.0%	1,101
Peritoneum, omentum & mesentery	0.0	0.4	1.8	6.5	18.5	30.3	28.6	13.9	100.0%	4,525
Other digestive system	0.0	0.5	1.5	7.0	17.6	24.8	27.4	21.2	100.0%	6,026
Respiratory System:	0.0	0.1	0.7	6.7	20.3	31.6	28.2	12.3	100.0%	803,178
Nose, nasal cavity & middle ear	0.4	2.5	3.6	12.3	20.0	25.2	22.0	14.0	100.0%	2,303
Larynx	0.0	0.1	0.9	9.6	27.1	30.1	22.0	10.2	100.0%	18,643
Lung & bronchus	0.0	0.1	0.7	6.6	20.2	31.7	28.4	12.3	100.0%	779,796
Lung & bronchus (Male)	0.0	0.1	0.7	6.5	21.4	32.7	27.8	10.8	100.0%	427,587
Lung & bronchus (Female)	0.0	0.1	0.8	6.8	18.7	30.5	29.1	14.1	100.0%	352,209
Pleura	0.3	0.1	1.0	2.9	13.5	28.6	35.0	18.6	100.0%	1,260
Trachea & other respiratory organs	1.6	3.8	3.0	10.4	21.4	24.9	21.3	13.6	100.0%	1,176
Bones & joints	12.3	13.8	5.5	9.3	13.5	16.2	16.4	13.1	100.0%	7,399
Soft tissue (including heart)	3.5	5.9	6.0	12.4	19.8	21.4	19.2	11.8	100.0%	23,080
Skin (excl. basal & squamous):	0.1	1.5	3.6	9.4	18.5	22.6	24.7	19.6	100.0%	63,570
Melanoma of the skin	0.1	1.9	4.5	10.6	19.7	23.1	24.1	15.9	100.0%	45,982
Other non-epithelial skin	0.0	0.4	1.1	6.2	15.3	21.3	26.2	29.4	100.0%	17,588
Breast (Female)	0.0	0.9	4.7	13.3	22.0	22.4	19.7	16.9	100.0%	205,675

Source: US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.
Percents may not sum to 100 due to rounding.

Table 1.13 - continued

Age Distribution (%) of Deaths by Site, 2011-2015

All Races, Both Sexes

Site	Age at Death								All Ages	Deaths
	<20	20-34	35-44	45-54	55-64	65-74	75-84	85+		
Female Genital System:	0.0	1.2	3.7	10.8	22.0	26.3	22.2	13.8	100.0%	148,985
Cervix uteri	0.0	5.2	13.4	22.7	24.1	17.2	11.2	6.3	100.0%	20,673
Corpus uteri	0.0	0.4	1.7	6.8	24.3	31.5	22.8	12.5	100.0%	19,939
Uterus, NOS	0.0	0.5	1.9	8.1	23.0	29.3	22.3	14.9	100.0%	26,761
Ovary	0.1	0.7	2.3	9.9	21.2	26.8	24.6	14.4	100.0%	71,141
Vagina	0.1	0.6	2.5	7.3	15.5	21.6	26.2	26.2	100.0%	2,162
Vulva	0.0	0.6	1.6	7.4	13.4	19.0	28.0	30.0	100.0%	5,271
Other female genital system	0.1	0.9	2.2	8.5	20.7	29.4	24.2	14.1	100.0%	3,038
Male Genital System:	0.0	0.5	0.4	1.9	9.4	21.5	33.5	32.9	100.0%	143,694
Prostate	0.0	0.0	0.1	1.5	9.2	21.7	34.0	33.5	100.0%	140,086
Testis	2.0	34.5	18.0	19.0	11.7	6.7	4.9	3.2	100.0%	1,934
Penis	0.0	1.3	3.8	10.3	18.7	25.9	24.6	15.4	100.0%	1,392
Other male genital system	0.0	1.1	2.1	8.2	11.0	23.8	33.3	20.6	100.0%	282
Urinary System:	0.1	0.3	1.0	5.6	15.9	24.4	28.9	23.7	100.0%	151,566
Urinary bladder	0.0	0.1	0.5	3.4	11.6	21.9	32.2	30.4	100.0%	78,045
Kidney & renal pelvis	0.3	0.6	1.6	8.3	21.1	27.3	24.8	16.0	100.0%	69,348
Ureter	0.1	0.1	0.3	2.1	10.2	22.8	36.8	27.8	100.0%	1,970
Other urinary system	0.0	0.2	1.2	4.5	12.6	23.3	32.5	25.7	100.0%	2,203
Eye & Orbit	2.0	2.0	3.0	9.7	22.2	24.5	22.0	14.6	100.0%	1,536
Brain & Nervous System:	3.6	3.4	5.3	12.8	24.0	25.2	18.0	7.7	100.0%	77,375
Endocrine System:	5.7	2.2	3.4	9.2	18.3	23.4	22.7	15.2	100.0%	13,906
Thyroid	0.2	0.8	1.7	7.5	18.0	25.6	27.1	19.1	100.0%	9,013
Other endocrine & thymus	15.9	4.9	6.4	12.3	18.8	19.3	14.6	7.8	100.0%	4,893
Lymphoma:	0.4	1.7	2.2	5.8	14.1	23.4	30.7	21.7	100.0%	106,944
Hodgkin lymphoma	1.0	10.7	7.5	11.0	15.1	20.8	22.3	11.7	100.0%	5,585
Non-Hodgkin lymphoma	0.3	1.2	1.9	5.5	14.1	23.6	31.2	22.2	100.0%	101,359
Myeloma	0.0	0.1	0.9	5.1	15.6	27.4	32.3	18.7	100.0%	59,376
Leukemia:	2.2	2.6	2.4	5.5	12.6	23.1	30.0	21.6	100.0%	116,463
Lymphocytic:	3.3	3.4	2.1	4.5	10.7	19.4	28.6	27.9	100.0%	31,743
Acute lymphocytic	14.5	14.1	8.1	11.3	14.8	16.3	13.2	7.7	100.0%	7,276
Chronic lymphocytic	0.0	0.1	0.3	2.3	9.4	20.4	33.4	34.1	100.0%	22,478
Other lymphocytic	0.5	1.3	1.1	4.3	11.0	20.3	30.7	31.0	100.0%	1,989
Myeloid & Monocytic:	1.6	2.5	2.8	6.5	14.6	26.1	29.9	15.9	100.0%	58,357
Acute myeloid	1.8	2.6	2.8	6.7	15.2	27.1	29.5	14.2	100.0%	48,632
Chronic myeloid	0.6	3.1	3.7	7.1	11.3	18.2	29.8	26.1	100.0%	5,300
Acute monocytic	2.4	0.4	1.5	5.0	13.1	23.3	31.9	22.4	100.0%	464
Other myeloid & monocytic	0.7	1.1	1.6	4.1	11.6	24.6	34.9	21.4	100.0%	3,961
Other leukemia:	1.9	2.0	1.9	4.7	10.6	20.6	31.8	26.6	100.0%	26,363
Other acute leukemia	1.2	2.2	2.0	4.6	10.3	21.1	33.1	25.6	100.0%	8,794
Aleukemic, subleukemic & NOS	2.2	2.0	1.8	4.7	10.7	20.3	31.1	27.1	100.0%	17,569
Ill-defined & unspecified	0.2	0.7	1.6	7.0	18.0	24.7	27.0	20.8	100.0%	214,371

Source: US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.
 Percents may not sum to 100 due to rounding.

Table 1.14
 Median Age of Cancer Patients at Death^a, 2011-2015
 By Primary Cancer Site, Race and Sex

Site	All Races			Whites			Blacks		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
All Sites	72.0	72.0	73.0	73.0	72.0	73.0	67.0	67.0	68.0
Oral Cavity & Pharynx:	67.0	66.0	73.0	68.0	66.0	74.0	63.0	63.0	64.0
Lip	76.0	73.0	82.0	77.0	74.0	83.0	-	-	-
Tongue	67.0	65.0	71.0	67.0	65.5	72.0	63.0	63.0	64.0
Salivary gland	74.0	74.0	74.5	75.0	75.0	76.0	64.0	64.0	66.5
Floor of mouth	66.0	64.0	72.0	66.0	64.0	72.5	64.0	64.0	-
Gum & other oral cavity	73.0	68.0	80.0	74.0	69.0	81.0	64.0	63.0	68.0
Nasopharynx	63.0	62.0	65.0	65.0	64.0	69.0	59.0	59.0	61.0
Tonsil	63.0	63.0	66.0	64.0	63.0	68.0	62.0	62.0	62.0
Oropharynx	66.0	65.0	71.0	67.0	65.0	72.0	62.0	62.0	63.0
Hypopharynx	66.0	66.0	68.0	67.0	67.0	69.0	62.0	63.0	58.0
Other oral cavity & pharynx	67.0	67.0	71.0	68.0	67.0	72.0	64.0	64.0	66.0
Digestive System:	71.0	69.0	74.0	72.0	70.0	75.0	67.0	65.0	70.0
Esophagus	69.0	68.0	73.0	69.0	68.0	74.0	65.0	64.0	66.0
Stomach	71.0	70.0	74.0	72.0	71.0	75.0	69.0	67.0	72.0
Small intestine	72.0	71.0	73.0	73.0	72.0	74.0	66.0	66.0	66.0
Colon & Rectum	73.0	70.0	76.0	74.0	71.0	77.0	67.0	66.0	69.0
Anus, anal canal & anorectum	65.0	63.0	66.0	66.0	64.0	67.0	60.0	57.0	63.0
Liver & intrahepatic bile duct	67.0	65.0	72.0	68.0	66.0	73.0	63.0	62.0	66.0
Gallbladder	73.0	72.0	74.0	74.0	73.0	75.0	70.0	71.0	69.0
Other biliary	75.0	73.0	76.0	75.0	74.0	77.0	70.0	68.0	71.0
Pancreas	72.0	70.0	75.0	73.0	71.0	75.0	69.0	66.0	71.0
Retroperitoneum	71.0	70.0	71.0	71.0	70.0	74.0	69.0	69.5	66.0
Peritoneum, omentum & mesentery	72.0	69.0	73.0	72.0	69.0	73.0	70.0	64.0	71.0
Other digestive system	74.0	71.0	77.0	75.0	72.0	78.0	70.0	68.0	73.0
Respiratory System:	72.0	71.0	72.0	72.0	71.0	73.0	68.0	67.0	69.0
Nose, nasal cavity & middle ear	69.0	66.0	73.0	69.0	67.0	74.0	66.0	63.0	69.0
Larynx	68.0	68.0	69.0	69.0	69.0	70.0	66.0	66.0	65.0
Lung & bronchus	72.0	71.0	72.0	72.0	72.0	73.0	68.0	67.0	69.0
Pleura	76.0	76.0	75.0	76.0	76.0	76.0	72.0	72.0	72.0
Trachea & other respiratory organs	68.0	66.0	73.0	70.0	67.0	73.0	63.0	63.0	63.0
Bones & joints	61.0	59.0	65.0	63.0	61.0	66.0	54.0	52.0	57.0
Soft tissue (including heart)	66.0	66.0	65.0	67.0	67.0	66.0	59.0	57.0	60.0
Skin (excl. basal & squamous):	72.0	72.0	73.0	72.0	72.0	73.0	65.0	63.0	67.0
Melanoma of the skin	70.0	70.0	70.0	70.0	70.0	70.0	67.0	65.0	70.0
Other non-epithelial skin	77.0	75.0	82.0	78.0	76.0	82.0	63.0	62.0	64.0
Breast	68.0	70.0	68.0	70.0	71.0	70.0	62.0	66.0	62.0

^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.
 - Statistic could not be calculated. Less than 16 deaths occurred during the time interval.

Table 1.14 - continued
 Median Age of Cancer Patients at Death^a, 2011-2015
 By Primary Cancer Site, Race and Sex

Site	All Races			Whites			Blacks		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
Female Genital System:	69.0	-	69.0	70.0	-	70.0	66.0	-	66.0
Cervix uteri	58.0	-	58.0	58.0	-	58.0	58.5	-	58.5
Corpus uteri	70.0	-	70.0	70.0	-	70.0	68.0	-	68.0
Uterus, NOS	70.0	-	70.0	71.0	-	71.0	67.0	-	67.0
Ovary	70.0	-	70.0	71.0	-	71.0	68.0	-	68.0
Vagina	75.5	-	75.5	77.0	-	77.0	71.0	-	71.0
Vulva	78.0	-	78.0	79.0	-	79.0	66.0	-	66.0
Other female genital system	70.0	-	70.0	71.0	-	71.0	67.0	-	67.0
Male Genital System:	80.0	80.0	-	81.0	81.0	-	76.0	76.0	-
Prostate	80.0	80.0	-	81.0	81.0	-	76.0	76.0	-
Testis	42.0	42.0	-	42.0	42.0	-	43.5	43.5	-
Penis	71.0	71.0	-	72.0	72.0	-	67.5	67.5	-
Other male genital system	76.0	76.0	-	78.0	78.0	-	64.5	64.5	-
Urinary System:	75.0	75.0	78.0	76.0	75.0	78.0	70.0	68.0	73.0
Urinary bladder	79.0	78.0	81.0	79.0	79.0	81.0	74.0	73.0	77.0
Kidney & renal pelvis	71.0	70.0	74.0	72.0	70.0	75.0	67.0	65.0	70.5
Ureter	79.0	78.0	80.0	79.0	78.0	81.0	75.0	72.0	75.0
Other urinary system	77.0	77.0	78.0	78.0	77.0	80.0	70.0	72.0	68.5
Eye & Orbit	69.0	67.0	71.0	69.0	68.0	71.0	61.0	62.5	59.0
Brain & Nervous System	65.0	64.0	66.0	65.0	64.0	66.0	60.0	59.0	62.0
Endocrine System:	69.0	67.0	72.0	70.0	67.0	72.0	64.0	62.0	66.5
Thyroid	73.0	71.0	75.0	74.0	71.0	76.0	69.0	66.5	71.0
Other endocrine & thymus	60.0	59.0	62.0	61.0	60.0	63.0	57.0	55.0	58.0
Lymphoma:	75.0	74.0	78.0	76.0	74.0	78.0	66.0	64.0	69.0
Hodgkin lymphoma	67.0	65.0	69.0	68.0	67.0	71.0	53.0	53.0	56.0
Non-Hodgkin lymphoma	76.0	74.0	78.0	76.0	75.0	79.0	67.0	65.0	69.0
Myeloma	75.0	74.0	76.0	76.0	75.0	77.0	71.0	70.0	73.0
Leukemia:	75.0	74.0	76.0	76.0	75.0	77.0	68.0	67.0	70.0
Lymphocytic:	77.0	75.0	80.0	78.0	76.0	81.0	70.0	68.0	74.0
Acute lymphocytic	56.0	53.0	59.0	57.0	55.0	60.0	48.5	44.0	53.0
Chronic lymphocytic	80.0	78.0	83.0	81.0	79.0	84.0	74.0	72.0	78.0
Other lymphocytic	79.0	77.0	81.0	79.0	78.0	82.0	73.0	67.0	77.0
Myeloid & Monocytic:	73.0	73.0	74.0	74.0	73.0	74.0	66.0	66.0	67.0
Acute myeloid	72.0	72.0	73.0	73.0	73.0	73.0	66.0	66.0	66.5
Chronic myeloid	77.0	75.0	79.0	78.0	76.0	80.0	63.0	59.0	69.0
Acute monocytic	76.0	75.0	77.0	77.0	76.0	78.0	69.0	68.0	-
Other myeloid & monocytic	76.0	76.0	78.0	77.0	76.0	79.0	70.0	68.0	71.5
Other leukemia:	77.0	76.0	79.0	78.0	77.0	80.0	70.0	68.0	73.0
Other acute leukemia	77.0	76.0	79.0	78.0	77.0	79.0	71.0	69.0	74.0
Aleukemic, subleukemic & NOS	77.0	76.0	79.0	78.0	77.0	80.0	70.0	68.0	73.0
Ill-defined & unspecified	74.0	72.0	76.0	74.0	73.0	76.0	68.0	66.0	70.0

^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.
 - Statistic could not be calculated. Less than 16 deaths occurred during the time interval.

Table 1.15

Lifetime Risk (Percent) of Being Diagnosed with Cancer by Site and Race/Ethnicity
Both Sexes, 18 SEER Areas, 2013-2015

Site	All Races		Whites		Blacks	
	Percent	(95% C.I.)	Percent	(95% C.I.)	Percent	(95% C.I.)
All Sites	38.38	(38.30, 38.46)	38.61	(38.52, 38.70)	35.18	(34.94, 35.42)
Invasive and In Situ	40.80	(40.72, 40.89)	41.02	(40.93, 41.11)	36.47	(36.23, 36.71)
Oral Cavity and Pharynx	1.15	(1.14, 1.17)	1.21	(1.20, 1.23)	0.79	(0.76, 0.82)
Esophagus	0.48	(0.47, 0.49)	0.51	(0.50, 0.52)	0.38	(0.36, 0.41)
Stomach	0.83	(0.82, 0.84)	0.73	(0.72, 0.74)	1.01	(0.97, 1.06)
Colon and Rectum	4.23	(4.21, 4.26)	4.16	(4.13, 4.19)	4.32	(4.24, 4.41)
Invasive and In Situ	4.35	(4.33, 4.38)	4.27	(4.24, 4.30)	4.47	(4.38, 4.56)
Liver and Intrahepatic Bile Duct	1.01	(1.00, 1.03)	0.92	(0.90, 0.93)	1.06	(1.03, 1.10)
Pancreas	1.57	(1.56, 1.59)	1.57	(1.55, 1.58)	1.63	(1.58, 1.69)
Larynx	0.32	(0.31, 0.32)	0.32	(0.31, 0.33)	0.42	(0.40, 0.45)
Invasive and In Situ	0.34	(0.33, 0.35)	0.34	(0.33, 0.35)	0.44	(0.41, 0.47)
Lung and Bronchus	6.24	(6.21, 6.28)	6.38	(6.34, 6.42)	5.94	(5.84, 6.04)
Melanoma of the Skin	2.27	(2.25, 2.29)	2.67	(2.65, 2.70)	0.11	(0.09, 0.12)
Invasive and In Situ	3.97	(3.95, 4.00)	4.54	(4.51, 4.57)	0.15	(0.13, 0.17)
Breast	6.39	(6.36, 6.42)	6.44	(6.40, 6.48)	6.11	(6.02, 6.21)
Invasive and In Situ	7.55	(7.51, 7.58)	7.55	(7.51, 7.59)	7.30	(7.20, 7.40)
Urinary Bladder (Invasive and In Situ)	2.33	(2.31, 2.35)	2.52	(2.50, 2.55)	1.26	(1.21, 1.31)
Kidney and Renal Pelvis	1.65	(1.64, 1.67)	1.70	(1.69, 1.72)	1.61	(1.56, 1.66)
Brain and Other Nervous System	0.62	(0.61, 0.63)	0.68	(0.67, 0.69)	0.34	(0.32, 0.36)
Thyroid	1.22	(1.21, 1.23)	1.28	(1.27, 1.30)	0.74	(0.71, 0.77)
Hodgkin Lymphoma	0.20	(0.20, 0.21)	0.22	(0.21, 0.22)	0.19	(0.18, 0.21)
Non-Hodgkin Lymphoma	2.11	(2.09, 2.13)	2.21	(2.18, 2.23)	1.29	(1.25, 1.34)
Myeloma	0.77	(0.76, 0.78)	0.71	(0.69, 0.72)	1.30	(1.26, 1.35)
Leukemia	1.51	(1.49, 1.52)	1.59	(1.57, 1.61)	1.03	(0.99, 1.07)
Acute Lymphocytic Leukemia	0.13	(0.13, 0.14)	0.15	(0.14, 0.15)	0.07	(0.06, 0.08)
Chronic Lymphocytic Leukemia	0.57	(0.56, 0.58)	0.61	(0.60, 0.62)	0.36	(0.34, 0.39)
Acute Myeloid Leukemia	0.48	(0.47, 0.49)	0.50	(0.49, 0.51)	0.35	(0.33, 0.38)
Chronic Myeloid Leukemia	0.20	(0.19, 0.21)	0.20	(0.20, 0.21)	0.16	(0.14, 0.17)
Kaposi Sarcoma	0.12	(0.11, 0.12)	0.13	(0.13, 0.14)	0.05	(0.04, 0.06)
Mesothelioma	0.04	(0.04, 0.05)	0.04	(0.03, 0.04)	0.07	(0.06, 0.08)

Devcan Version 6.7.6, April 2018, National Cancer Institute (<https://surveillance.cancer.gov/devcan/>).

Source: SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, and Georgia excluding ATL/RG).

Note: Invasive cancer only unless specified otherwise.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table 1.15 - continued

Lifetime Risk (Percent) of Being Diagnosed with Cancer by Site and Race/Ethnicity
Both Sexes, 18 SEER Areas, 2013-2015

Site	Asian/Pacific Islanders	American Indian/ Alaska Natives ^a	Hispanics ^b
	Percent (95% C.I.)	Percent (95% C.I.)	Percent (95% C.I.)
All Sites	33.77 (33.46, 34.09)	28.31 (27.26, 29.44)	35.07 (34.80, 35.35)
Invasive and In Situ	35.14 (34.82, 35.46)	29.26 (28.21, 30.41)	36.28 (36.00, 36.56)
Oral Cavity and Pharynx	0.90 (0.86, 0.95)	0.80 (0.66, 1.06)	0.74 (0.70, 0.78)
Esophagus	0.29 (0.26, 0.33)	0.41 (0.30, 0.64)	0.35 (0.33, 0.39)
Stomach	1.64 (1.56, 1.72)	1.19 (0.96, 1.52)	1.35 (1.29, 1.41)
Colon and Rectum	4.41 (4.29, 4.53)	4.61 (4.18, 5.14)	4.16 (4.06, 4.26)
Invasive and In Situ	4.53 (4.41, 4.65)	4.69 (4.26, 5.22)	4.28 (4.18, 4.38)
Liver and Intrahepatic Bile Duct	1.84 (1.77, 1.91)	1.49 (1.30, 1.79)	1.71 (1.65, 1.77)
Pancreas	1.63 (1.56, 1.71)	1.14 (0.94, 1.45)	1.65 (1.59, 1.72)
Larynx	0.13 (0.11, 0.15)	0.20 (0.13, 0.42)	0.24 (0.22, 0.27)
Invasive and In Situ	0.14 (0.13, 0.17)	0.23 (0.15, 0.45)	0.26 (0.24, 0.29)
Lung and Bronchus	5.45 (5.31, 5.59)	4.15 (3.76, 4.64)	3.87 (3.77, 3.97)
Melanoma of the Skin	0.19 (0.16, 0.22)	0.49 (0.38, 0.73)	0.58 (0.55, 0.63)
Invasive and In Situ	0.25 (0.22, 0.28)	0.79 (0.64, 1.06)	0.92 (0.88, 0.98)
Breast	5.79 (5.68, 5.90)	3.99 (3.64, 4.43)	5.14 (5.04, 5.23)
Invasive and In Situ	7.16 (7.04, 7.28)	4.68 (4.30, 5.14)	6.05 (5.95, 6.15)
Urinary Bladder (Invasive and In Situ)	1.46 (1.39, 1.54)	1.08 (0.86, 1.42)	1.53 (1.47, 1.60)
Kidney and Renal Pelvis	1.13 (1.07, 1.19)	1.63 (1.42, 1.94)	1.84 (1.78, 1.90)
Brain and Other Nervous System	0.41 (0.38, 0.44)	0.31 (0.22, 0.53)	0.54 (0.51, 0.58)
Thyroid	1.28 (1.24, 1.33)	0.83 (0.70, 1.08)	1.14 (1.11, 1.18)
Hodgkin Lymphoma	0.11 (0.10, 0.13)	0.16 (0.09, 0.38)	0.21 (0.20, 0.23)
Non-Hodgkin Lymphoma	1.84 (1.77, 1.92)	1.26 (1.01, 1.62)	2.15 (2.08, 2.22)
Myeloma	0.56 (0.53, 0.61)	0.70 (0.54, 0.96)	0.78 (0.74, 0.82)
Leukemia	0.98 (0.93, 1.04)	0.78 (0.63, 1.04)	1.19 (1.14, 1.24)
Acute Lymphocytic Leukemia	0.11 (0.10, 0.13)	0.13 (0.09, 0.34)	0.20 (0.19, 0.22)
Chronic Lymphocytic Leukemia	0.14 (0.13, 0.17)	0.16 (0.09, 0.37)	0.29 (0.26, 0.32)
Acute Myeloid Leukemia	0.47 (0.43, 0.51)	0.25 (0.16, 0.47)	0.42 (0.39, 0.45)
Chronic Myeloid Leukemia	0.16 (0.14, 0.18)	0.14 (0.09, 0.34)	0.16 (0.14, 0.18)
Kaposi Sarcoma	0.06 (0.04, 0.08)	0.05 (0.01, 0.26)	0.13 (0.11, 0.15)
Mesothelioma	0.02 (0.02, 0.04)	0.04 (0.01, 0.26)	0.07 (0.06, 0.08)

Devcan Version 6.7.6, April 2018, National Cancer Institute (<https://surveillance.cancer.gov/devcan/>).

Source: SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, and Georgia excluding ATL/RG).

Note: Invasive cancer only unless specified otherwise.

^a Underlying incidence data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry. A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table 1.16

Lifetime Risk (Percent) of Being Diagnosed with Cancer by Site and Race/Ethnicity
Males, 18 SEER Areas, 2013-2015

Site	All Races		Whites		Blacks	
	Percent	(95% C.I.)	Percent	(95% C.I.)	Percent	(95% C.I.)
All Sites	39.26	(39.14, 39.38)	39.08	(38.95, 39.22)	36.64	(36.28, 37.00)
Invasive and In Situ	40.91	(40.79, 41.04)	40.75	(40.61, 40.89)	36.94	(36.59, 37.31)
Oral Cavity and Pharynx	1.64	(1.61, 1.66)	1.73	(1.70, 1.75)	1.12	(1.06, 1.18)
Esophagus	0.75	(0.74, 0.77)	0.80	(0.79, 0.82)	0.55	(0.51, 0.59)
Stomach	1.04	(1.02, 1.06)	0.93	(0.91, 0.95)	1.20	(1.14, 1.27)
Colon and Rectum	4.42	(4.38, 4.46)	4.34	(4.29, 4.38)	4.42	(4.30, 4.55)
Invasive and In Situ	4.55	(4.51, 4.59)	4.46	(4.41, 4.50)	4.58	(4.46, 4.71)
Liver and Intrahepatic Bile Duct	1.42	(1.40, 1.45)	1.28	(1.26, 1.30)	1.60	(1.54, 1.67)
Pancreas	1.60	(1.58, 1.63)	1.60	(1.58, 1.63)	1.57	(1.49, 1.65)
Larynx	0.52	(0.51, 0.54)	0.52	(0.51, 0.54)	0.72	(0.67, 0.77)
Invasive and In Situ	0.56	(0.54, 0.57)	0.56	(0.54, 0.57)	0.75	(0.70, 0.80)
Lung and Bronchus	6.68	(6.62, 6.73)	6.67	(6.62, 6.73)	6.89	(6.73, 7.06)
Melanoma of the Skin	2.82	(2.79, 2.85)	3.28	(3.24, 3.32)	0.11	(0.09, 0.14)
Invasive and In Situ	4.87	(4.83, 4.92)	5.52	(5.47, 5.57)	0.16	(0.13, 0.19)
Breast	0.12	(0.12, 0.13)	0.12	(0.12, 0.13)	0.15	(0.13, 0.18)
Invasive and In Situ	0.14	(0.13, 0.14)	0.14	(0.13, 0.14)	0.17	(0.14, 0.20)
Prostate	11.18	(11.12, 11.24)	10.46	(10.40, 10.53)	14.77	(14.55, 14.99)
Testis	0.40	(0.39, 0.41)	0.48	(0.46, 0.49)	0.10	(0.09, 0.12)
Urinary Bladder (Invasive and In Situ)	3.69	(3.65, 3.73)	3.99	(3.95, 4.03)	1.79	(1.70, 1.88)
Kidney and Renal Pelvis	2.11	(2.09, 2.14)	2.18	(2.15, 2.21)	2.00	(1.93, 2.09)
Brain and Other Nervous System	0.69	(0.67, 0.70)	0.76	(0.74, 0.78)	0.36	(0.33, 0.40)
Thyroid	0.64	(0.63, 0.65)	0.68	(0.67, 0.70)	0.31	(0.28, 0.34)
Hodgkin Lymphoma	0.23	(0.22, 0.24)	0.24	(0.23, 0.25)	0.23	(0.21, 0.25)
Non-Hodgkin Lymphoma	2.37	(2.34, 2.40)	2.47	(2.44, 2.51)	1.43	(1.36, 1.50)
Myeloma	0.89	(0.87, 0.91)	0.83	(0.81, 0.85)	1.38	(1.31, 1.45)
Leukemia	1.78	(1.75, 1.80)	1.87	(1.84, 1.90)	1.16	(1.10, 1.23)
Acute Lymphocytic Leukemia	0.14	(0.14, 0.15)	0.16	(0.15, 0.17)	0.07	(0.06, 0.09)
Chronic Lymphocytic Leukemia	0.71	(0.69, 0.72)	0.75	(0.73, 0.77)	0.45	(0.42, 0.50)
Acute Myeloid Leukemia	0.55	(0.53, 0.56)	0.57	(0.55, 0.58)	0.40	(0.36, 0.44)
Chronic Myeloid Leukemia	0.24	(0.23, 0.25)	0.25	(0.24, 0.26)	0.17	(0.15, 0.20)
Kaposi Sarcoma	0.19	(0.18, 0.20)	0.21	(0.20, 0.22)	0.08	(0.06, 0.11)
Mesothelioma	0.07	(0.07, 0.08)	0.06	(0.06, 0.07)	0.13	(0.12, 0.15)

Devcan Version 6.7.6, April 2018, National Cancer Institute (<https://surveillance.cancer.gov/devcan/>).

Source: SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, and Georgia excluding ATL/RG).

Note: Invasive cancer only unless specified otherwise.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table 1.16 - continued

Lifetime Risk (Percent) of Being Diagnosed with Cancer by Site and Race/Ethnicity
Males, 18 SEER Areas, 2013-2015

Site	Asian/Pacific Islanders	American Indian/ Alaska Natives ^a	Hispanics ^b
	Percent (95% C.I.)	Percent (95% C.I.)	Percent (95% C.I.)
All Sites	33.93 (33.46, 34.42)	27.36 (25.87, 29.09)	36.09 (35.66, 36.52)
Invasive and In Situ	34.23 (33.76, 34.72)	27.78 (26.28, 29.52)	36.55 (36.12, 36.99)
Oral Cavity and Pharynx	1.23 (1.15, 1.31)	0.90 (0.72, 1.45)	0.99 (0.93, 1.07)
Esophagus	0.47 (0.41, 0.54)	0.61 (0.42, 1.19)	0.55 (0.50, 0.61)
Stomach	1.97 (1.84, 2.10)	1.34 (1.06, 1.95)	1.56 (1.47, 1.66)
Colon and Rectum	4.80 (4.63, 4.98)	4.30 (3.69, 5.19)	4.49 (4.35, 4.65)
Invasive and In Situ	4.96 (4.78, 5.14)	4.40 (3.79, 5.29)	4.63 (4.48, 4.79)
Liver and Intrahepatic Bile Duct	2.57 (2.45, 2.71)	1.90 (1.60, 2.51)	2.32 (2.22, 2.43)
Pancreas	1.66 (1.55, 1.78)	1.33 (1.01, 1.97)	1.60 (1.50, 1.70)
Larynx	0.25 (0.21, 0.30)	0.34 (0.21, 0.89)	0.45 (0.40, 0.50)
Invasive and In Situ	0.27 (0.23, 0.32)	0.39 (0.24, 0.95)	0.48 (0.43, 0.54)
Lung and Bronchus	6.59 (6.37, 6.82)	4.80 (4.21, 5.66)	4.36 (4.20, 4.53)
Melanoma of the Skin	0.20 (0.16, 0.25)	0.50 (0.33, 1.06)	0.61 (0.55, 0.68)
Invasive and In Situ	0.27 (0.23, 0.33)	0.90 (0.65, 1.50)	0.95 (0.88, 1.04)
Breast	0.08 (0.06, 0.12)	0.02 (0.00, 0.59)	0.09 (0.07, 0.14)
Invasive and In Situ	0.09 (0.07, 0.13)	0.02 (0.00, 0.59)	0.10 (0.08, 0.14)
Prostate	7.32 (7.12, 7.54)	5.53 (4.87, 6.46)	10.39 (10.18, 10.62)
Testis	0.17 (0.15, 0.20)	0.41 (0.32, 0.93)	0.39 (0.37, 0.42)
Urinary Bladder (Invasive and In Situ)	2.38 (2.24, 2.54)	1.46 (1.13, 2.11)	2.47 (2.34, 2.60)
Kidney and Renal Pelvis	1.48 (1.39, 1.59)	1.97 (1.63, 2.61)	2.30 (2.21, 2.41)
Brain and Other Nervous System	0.45 (0.41, 0.51)	0.37 (0.22, 0.92)	0.59 (0.54, 0.65)
Thyroid	0.67 (0.62, 0.73)	0.34 (0.22, 0.89)	0.50 (0.46, 0.55)
Hodgkin Lymphoma	0.13 (0.11, 0.17)	0.11 (0.04, 0.67)	0.23 (0.21, 0.27)
Non-Hodgkin Lymphoma	2.13 (2.02, 2.26)	1.20 (0.82, 1.92)	2.29 (2.19, 2.41)
Myeloma	0.69 (0.62, 0.77)	0.69 (0.47, 1.28)	0.89 (0.83, 0.97)
Leukemia	1.16 (1.08, 1.26)	0.91 (0.70, 1.48)	1.38 (1.30, 1.48)
Acute Lymphocytic Leukemia	0.11 (0.09, 0.14)	0.15 (0.10, 0.69)	0.22 (0.20, 0.25)
Chronic Lymphocytic Leukemia	0.18 (0.15, 0.23)	0.19 (0.09, 0.74)	0.34 (0.30, 0.40)
Acute Myeloid Leukemia	0.54 (0.48, 0.61)	0.29 (0.17, 0.84)	0.48 (0.43, 0.54)
Chronic Myeloid Leukemia	0.20 (0.17, 0.25)	0.20 (0.12, 0.74)	0.21 (0.18, 0.25)
Kaposi Sarcoma	0.09 (0.06, 0.13)	0.10 (0.03, 0.66)	0.22 (0.18, 0.26)
Mesothelioma	0.05 (0.03, 0.08)	0.04 (0.01, 0.61)	0.11 (0.09, 0.15)

Devcan Version 6.7.6, April 2018, National Cancer Institute (<https://surveillance.cancer.gov/devcan/>).

Source: SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, and Georgia excluding ATL/RG).

Note: Invasive cancer only unless specified otherwise.

^a Underlying incidence data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry. A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table 1.17

Lifetime Risk (Percent) of Being Diagnosed with Cancer by Site and Race/Ethnicity
Females, 18 SEER Areas, 2013-2015

Site	All Races		Whites		Blacks	
	Percent	(95% C.I.)	Percent	(95% C.I.)	Percent	(95% C.I.)
All Sites	37.73	(37.62, 37.84)	38.38	(38.25, 38.50)	33.96	(33.64, 34.29)
Invasive and In Situ	40.93	(40.82, 41.05)	41.56	(41.43, 41.69)	36.16	(35.83, 36.49)
Oral Cavity and Pharynx	0.69	(0.67, 0.71)	0.71	(0.69, 0.73)	0.50	(0.46, 0.54)
Esophagus	0.22	(0.21, 0.23)	0.23	(0.22, 0.24)	0.24	(0.21, 0.27)
Stomach	0.64	(0.62, 0.66)	0.54	(0.53, 0.56)	0.84	(0.79, 0.90)
Colon and Rectum	4.06	(4.02, 4.10)	4.00	(3.96, 4.04)	4.24	(4.12, 4.36)
Invasive and In Situ	4.16	(4.13, 4.20)	4.10	(4.05, 4.14)	4.37	(4.25, 4.49)
Liver and Intrahepatic Bile Duct	0.62	(0.60, 0.63)	0.56	(0.54, 0.57)	0.58	(0.54, 0.62)
Pancreas	1.55	(1.53, 1.57)	1.53	(1.50, 1.56)	1.69	(1.62, 1.77)
Larynx	0.12	(0.12, 0.13)	0.13	(0.12, 0.13)	0.16	(0.14, 0.18)
Invasive and In Situ	0.13	(0.13, 0.14)	0.14	(0.13, 0.14)	0.17	(0.15, 0.19)
Lung and Bronchus	5.87	(5.83, 5.92)	6.14	(6.09, 6.19)	5.15	(5.02, 5.28)
Melanoma of the Skin	1.78	(1.76, 1.81)	2.12	(2.09, 2.15)	0.11	(0.09, 0.13)
Invasive and In Situ	3.17	(3.14, 3.20)	3.65	(3.61, 3.68)	0.15	(0.13, 0.17)
Breast	12.44	(12.38, 12.50)	12.69	(12.62, 12.76)	11.48	(11.30, 11.65)
Invasive and In Situ	14.72	(14.65, 14.79)	14.91	(14.83, 14.98)	13.73	(13.54, 13.92)
Cervix Uteri	0.62	(0.60, 0.63)	0.60	(0.59, 0.62)	0.71	(0.67, 0.76)
Corpus and Uterus, NOS	2.86	(2.84, 2.89)	2.93	(2.90, 2.96)	2.67	(2.59, 2.76)
Invasive and In Situ	2.88	(2.85, 2.91)	2.95	(2.91, 2.98)	2.70	(2.62, 2.79)
Ovary ^a	1.27	(1.25, 1.29)	1.33	(1.30, 1.35)	0.93	(0.88, 0.99)
Urinary Bladder (Invasive and In Situ)	1.10	(1.08, 1.12)	1.16	(1.14, 1.18)	0.81	(0.76, 0.87)
Kidney and Renal Pelvis	1.22	(1.20, 1.23)	1.24	(1.22, 1.27)	1.26	(1.20, 1.32)
Brain and Other Nervous System	0.55	(0.54, 0.56)	0.61	(0.59, 0.62)	0.32	(0.29, 0.35)
Thyroid	1.81	(1.78, 1.83)	1.90	(1.88, 1.93)	1.14	(1.10, 1.20)
Hodgkin Lymphoma	0.18	(0.17, 0.19)	0.19	(0.18, 0.20)	0.16	(0.15, 0.18)
Non-Hodgkin Lymphoma	1.87	(1.84, 1.89)	1.96	(1.93, 1.99)	1.18	(1.13, 1.24)
Myeloma	0.66	(0.65, 0.68)	0.59	(0.57, 0.60)	1.24	(1.18, 1.31)
Leukemia	1.26	(1.24, 1.28)	1.32	(1.30, 1.34)	0.92	(0.86, 0.97)
Acute Lymphocytic Leukemia	0.12	(0.12, 0.13)	0.13	(0.13, 0.14)	0.07	(0.06, 0.08)
Chronic Lymphocytic Leukemia	0.44	(0.43, 0.46)	0.48	(0.46, 0.49)	0.28	(0.25, 0.32)
Acute Myeloid Leukemia	0.42	(0.41, 0.44)	0.44	(0.43, 0.45)	0.32	(0.29, 0.36)
Chronic Myeloid Leukemia	0.16	(0.15, 0.17)	0.16	(0.16, 0.17)	0.14	(0.12, 0.17)
Kaposi Sarcoma	0.05	(0.05, 0.06)	0.06	(0.05, 0.06)	0.03	(0.02, 0.04)
Mesothelioma	0.01	(0.01, 0.01)	0.01	(0.01, 0.01)	0.01	(0.01, 0.02)

Devcan Version 6.7.6, April 2018, National Cancer Institute (<https://surveillance.cancer.gov/devcan/>).

Source: SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, and Georgia excluding ATL/RG).

Note: Invasive cancer only unless specified otherwise.

^a Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table 1.17 - continued

Lifetime Risk (Percent) of Being Diagnosed with Cancer by Site and Race/Ethnicity
Females, 18 SEER Areas, 2013-2015

Site	Asian/Pacific Islanders	American Indian/ Alaska Natives ^a	Hispanics ^b
	Percent (95% C.I.)	Percent (95% C.I.)	Percent (95% C.I.)
All Sites	33.81 (33.40, 34.23)	29.38 (27.91, 31.01)	34.63 (34.26, 35.00)
Invasive and In Situ	36.11 (35.69, 36.54)	30.88 (29.38, 32.53)	36.56 (36.19, 36.94)
Oral Cavity and Pharynx	0.62 (0.57, 0.68)	0.70 (0.50, 1.12)	0.51 (0.46, 0.57)
Esophagus	0.15 (0.12, 0.19)	0.21 (0.11, 0.59)	0.18 (0.15, 0.22)
Stomach	1.37 (1.27, 1.48)	1.02 (0.71, 1.55)	1.17 (1.10, 1.26)
Colon and Rectum	4.07 (3.92, 4.24)	4.93 (4.31, 5.72)	3.87 (3.74, 4.01)
Invasive and In Situ	4.17 (4.01, 4.33)	4.99 (4.37, 5.78)	3.97 (3.84, 4.11)
Liver and Intrahepatic Bile Duct	1.22 (1.13, 1.31)	1.09 (0.84, 1.55)	1.15 (1.08, 1.22)
Pancreas	1.61 (1.51, 1.72)	0.98 (0.72, 1.45)	1.71 (1.62, 1.81)
Larynx	0.03 (0.02, 0.06)	0.08 (0.03, 0.44)	0.06 (0.05, 0.09)
Invasive and In Situ	0.04 (0.03, 0.07)	0.08 (0.03, 0.44)	0.07 (0.06, 0.10)
Lung and Bronchus	4.52 (4.35, 4.69)	3.57 (3.06, 4.25)	3.48 (3.35, 3.61)
Melanoma of the Skin	0.18 (0.15, 0.22)	0.49 (0.35, 0.88)	0.58 (0.53, 0.63)
Invasive and In Situ	0.23 (0.19, 0.27)	0.70 (0.52, 1.10)	0.92 (0.86, 0.99)
Breast	10.70 (10.50, 10.91)	7.82 (7.15, 8.65)	9.85 (9.68, 10.03)
Invasive and In Situ	13.25 (13.03, 13.47)	9.18 (8.47, 10.06)	11.63 (11.45, 11.82)
Cervix Uteri	0.61 (0.57, 0.67)	0.75 (0.57, 1.15)	0.81 (0.77, 0.86)
Corpus and Uterus, NOS	2.33 (2.25, 2.42)	2.08 (1.77, 2.57)	2.57 (2.49, 2.67)
Invasive and In Situ	2.34 (2.25, 2.43)	2.08 (1.77, 2.57)	2.59 (2.50, 2.68)
Ovary ^c	1.13 (1.06, 1.20)	0.94 (0.72, 1.38)	1.21 (1.14, 1.28)
Urinary Bladder (Invasive and In Situ)	0.71 (0.65, 0.79)	0.73 (0.46, 1.23)	0.75 (0.69, 0.82)
Kidney and Renal Pelvis	0.82 (0.76, 0.90)	1.31 (1.07, 1.76)	1.43 (1.37, 1.51)
Brain and Other Nervous System	0.37 (0.33, 0.42)	0.26 (0.16, 0.62)	0.50 (0.46, 0.55)
Thyroid	1.83 (1.76, 1.91)	1.32 (1.09, 1.76)	1.79 (1.73, 1.85)
Hodgkin Lymphoma	0.09 (0.08, 0.12)	0.21 (0.10, 0.59)	0.19 (0.17, 0.22)
Non-Hodgkin Lymphoma	1.60 (1.51, 1.70)	1.34 (1.00, 1.88)	2.03 (1.94, 2.13)
Myeloma	0.47 (0.42, 0.52)	0.71 (0.50, 1.13)	0.68 (0.63, 0.74)
Leukemia	0.84 (0.77, 0.91)	0.64 (0.45, 1.05)	1.03 (0.97, 1.10)
Acute Lymphocytic Leukemia	0.11 (0.09, 0.14)	0.12 (0.06, 0.48)	0.19 (0.17, 0.21)
Chronic Lymphocytic Leukemia	0.11 (0.09, 0.14)	0.13 (0.06, 0.50)	0.24 (0.21, 0.28)
Acute Myeloid Leukemia	0.41 (0.37, 0.47)	0.20 (0.10, 0.58)	0.37 (0.34, 0.42)
Chronic Myeloid Leukemia	0.12 (0.09, 0.16)	0.08 (0.03, 0.44)	0.11 (0.09, 0.14)
Kaposi Sarcoma	0.04 (0.02, 0.07)	0.00 (0.00, 0.38)	0.05 (0.04, 0.07)
Mesothelioma	0.00 (0.00, 0.03)	0.04 (0.01, 0.42)	0.03 (0.02, 0.05)

Devcan Version 6.7.6, April 2018, National Cancer Institute (<https://surveillance.cancer.gov/devcan/>).

Source: SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, and Georgia excluding ATL/RG).

Note: Invasive cancer only unless specified otherwise.

^a Underlying incidence data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

^c Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table 1.18

Lifetime Risk (Percent) of Dying from Cancer by Site and Race/Ethnicity
Both Sexes, Total U.S., 2013-2015

Site	All Races	Whites	Blacks
	Percent (95% C.I.)	Percent (95% C.I.)	Percent (95% C.I.)
All Sites	19.95 (19.92, 19.97)	20.07 (20.04, 20.10)	20.09 (20.01, 20.18)
Oral Cavity and Pharynx	0.29 (0.29, 0.30)	0.29 (0.29, 0.30)	0.27 (0.26, 0.28)
Esophagus	0.47 (0.47, 0.48)	0.50 (0.49, 0.50)	0.34 (0.33, 0.35)
Stomach	0.38 (0.37, 0.38)	0.33 (0.33, 0.34)	0.61 (0.59, 0.63)
Colon and Rectum	1.79 (1.78, 1.80)	1.75 (1.74, 1.76)	2.07 (2.03, 2.10)
Liver and Intrahepatic Bile Duct	0.76 (0.76, 0.77)	0.72 (0.71, 0.72)	0.86 (0.84, 0.88)
Pancreas	1.36 (1.35, 1.37)	1.36 (1.35, 1.36)	1.46 (1.44, 1.49)
Larynx	0.12 (0.11, 0.12)	0.11 (0.11, 0.12)	0.17 (0.16, 0.18)
Lung and Bronchus	5.15 (5.13, 5.16)	5.26 (5.25, 5.28)	4.76 (4.71, 4.80)
Melanoma of the Skin	0.30 (0.30, 0.31)	0.35 (0.34, 0.35)	0.04 (0.04, 0.05)
Breast	1.35 (1.35, 1.36)	1.32 (1.32, 1.33)	1.70 (1.67, 1.73)
Urinary Bladder	0.62 (0.61, 0.62)	0.65 (0.64, 0.65)	0.43 (0.41, 0.44)
Kidney and Renal Pelvis	0.47 (0.46, 0.47)	0.48 (0.48, 0.49)	0.40 (0.38, 0.41)
Brain and Other Nervous System	0.47 (0.47, 0.47)	0.51 (0.51, 0.52)	0.25 (0.24, 0.26)
Thyroid	0.06 (0.06, 0.07)	0.06 (0.06, 0.07)	0.06 (0.05, 0.06)
Hodgkin Lymphoma	0.03 (0.03, 0.04)	0.04 (0.03, 0.04)	0.03 (0.02, 0.03)
Non-Hodgkin Lymphoma	0.73 (0.72, 0.73)	0.76 (0.76, 0.77)	0.43 (0.42, 0.45)
Myeloma	0.43 (0.42, 0.43)	0.40 (0.39, 0.40)	0.70 (0.68, 0.72)
Leukemia	0.83 (0.83, 0.84)	0.87 (0.86, 0.88)	0.59 (0.57, 0.60)
Acute Lymphocytic Leukemia	0.04 (0.04, 0.04)	0.04 (0.04, 0.05)	0.03 (0.02, 0.03)
Chronic Lymphocytic Leukemia	0.18 (0.17, 0.18)	0.19 (0.18, 0.19)	0.12 (0.11, 0.13)
Acute Myeloid Leukemia	0.33 (0.33, 0.34)	0.35 (0.34, 0.35)	0.22 (0.21, 0.23)
Chronic Myeloid Leukemia	0.04 (0.04, 0.04)	0.04 (0.04, 0.04)	0.03 (0.03, 0.03)

Devcan Version 6.7.6, April 2018, National Cancer Institute (<https://surveillance.cancer.gov/devcan/>).
Source: NCHS public use data file for the total US.
A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table 1.18 - continued

Lifetime Risk (Percent) of Dying from Cancer by Site and Race/Ethnicity
Both Sexes, Total U.S., 2013-2015

Site	Asian/Pacific Islanders	American Indian/ Alaska Natives ^a	Hispanics ^b
	Percent (95% C.I.)	Percent (95% C.I.)	Percent (95% C.I.)
All Sites	18.07 (17.86, 18.28)	16.49 (16.05, 16.94)	17.25 (17.13, 17.38)
Oral Cavity and Pharynx	0.32 (0.29, 0.35)	0.22 (0.17, 0.30)	0.23 (0.21, 0.25)
Esophagus	0.27 (0.24, 0.30)	0.38 (0.32, 0.48)	0.30 (0.28, 0.31)
Stomach	0.98 (0.92, 1.03)	0.53 (0.45, 0.64)	0.74 (0.71, 0.77)
Colon and Rectum	1.85 (1.77, 1.92)	1.85 (1.69, 2.03)	1.77 (1.72, 1.81)
Liver and Intrahepatic Bile Duct	1.52 (1.46, 1.58)	1.12 (1.02, 1.26)	1.29 (1.26, 1.32)
Pancreas	1.44 (1.38, 1.51)	1.00 (0.89, 1.14)	1.30 (1.26, 1.33)
Larynx	0.06 (0.05, 0.07)	0.08 (0.06, 0.15)	0.10 (0.09, 0.12)
Lung and Bronchus	4.17 (4.07, 4.28)	3.89 (3.68, 4.13)	2.87 (2.81, 2.92)
Melanoma of the Skin	0.06 (0.05, 0.07)	0.08 (0.04, 0.15)	0.11 (0.10, 0.12)
Breast	0.99 (0.94, 1.05)	0.90 (0.78, 1.06)	1.12 (1.09, 1.16)
Urinary Bladder	0.43 (0.39, 0.48)	0.29 (0.23, 0.39)	0.45 (0.42, 0.47)
Kidney and Renal Pelvis	0.32 (0.29, 0.36)	0.63 (0.55, 0.74)	0.52 (0.50, 0.54)
Brain and Other Nervous System	0.31 (0.28, 0.34)	0.24 (0.19, 0.32)	0.37 (0.35, 0.39)
Thyroid	0.11 (0.09, 0.13)	0.07 (0.04, 0.14)	0.09 (0.08, 0.11)
Hodgkin Lymphoma	0.02 (0.02, 0.04)	0.03 (0.01, 0.09)	0.05 (0.05, 0.06)
Non-Hodgkin Lymphoma	0.73 (0.68, 0.77)	0.50 (0.42, 0.62)	0.73 (0.70, 0.76)
Myeloma	0.27 (0.25, 0.30)	0.38 (0.31, 0.48)	0.41 (0.39, 0.43)
Leukemia	0.60 (0.56, 0.64)	0.47 (0.39, 0.57)	0.67 (0.65, 0.70)
Acute Lymphocytic Leukemia	0.04 (0.03, 0.05)	0.06 (0.04, 0.11)	0.07 (0.06, 0.07)
Chronic Lymphocytic Leukemia	0.04 (0.03, 0.05)	0.06 (0.02, 0.13)	0.09 (0.08, 0.10)
Acute Myeloid Leukemia	0.32 (0.29, 0.35)	0.19 (0.15, 0.27)	0.26 (0.25, 0.28)
Chronic Myeloid Leukemia	0.03 (0.02, 0.04)	0.01 (0.01, 0.07)	0.04 (0.03, 0.05)

Devcan Version 6.7.6, April 2018, National Cancer Institute (<https://surveillance.cancer.gov/devcan/>).

Source: NCHS public use data file for the total US.

^a Underlying mortality data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian Pacific Islanders, and American Indians/Alaska Natives. A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table 1.19

Lifetime Risk (Percent) of Dying from Cancer by Site and Race/Ethnicity
Males, Total U.S., 2013-2015

Site	All Races		Whites		Blacks	
	Percent	(95% C.I.)	Percent	(95% C.I.)	Percent	(95% C.I.)
All Sites	21.66	(21.62, 21.70)	21.75	(21.70, 21.79)	22.02	(21.88, 22.16)
Oral Cavity and Pharynx	0.41	(0.40, 0.42)	0.41	(0.40, 0.42)	0.41	(0.39, 0.43)
Esophagus	0.76	(0.75, 0.77)	0.81	(0.80, 0.82)	0.50	(0.48, 0.52)
Stomach	0.46	(0.45, 0.46)	0.40	(0.40, 0.41)	0.76	(0.73, 0.79)
Colon and Rectum	1.87	(1.86, 1.89)	1.83	(1.82, 1.85)	2.20	(2.15, 2.25)
Liver and Intrahepatic Bile Duct	1.01	(1.00, 1.02)	0.94	(0.93, 0.95)	1.21	(1.18, 1.24)
Pancreas	1.38	(1.37, 1.39)	1.39	(1.38, 1.40)	1.38	(1.34, 1.41)
Larynx	0.19	(0.19, 0.20)	0.19	(0.18, 0.19)	0.29	(0.28, 0.31)
Lung and Bronchus	5.74	(5.71, 5.76)	5.78	(5.76, 5.81)	5.78	(5.71, 5.86)
Melanoma of the Skin	0.41	(0.41, 0.42)	0.47	(0.47, 0.48)	0.04	(0.04, 0.05)
Breast	0.03	(0.03, 0.03)	0.03	(0.03, 0.03)	0.05	(0.04, 0.06)
Prostate	2.42	(2.41, 2.44)	2.27	(2.26, 2.29)	3.97	(3.89, 4.05)
Testis	0.02	(0.02, 0.02)	0.02	(0.02, 0.02)	0.01	(0.01, 0.01)
Urinary Bladder	0.93	(0.92, 0.94)	0.99	(0.98, 1.00)	0.55	(0.52, 0.57)
Kidney and Renal Pelvis	0.61	(0.61, 0.62)	0.63	(0.63, 0.64)	0.51	(0.49, 0.54)
Brain and Other Nervous System	0.52	(0.52, 0.53)	0.57	(0.56, 0.58)	0.27	(0.26, 0.29)
Thyroid	0.06	(0.05, 0.06)	0.06	(0.06, 0.06)	0.03	(0.03, 0.04)
Hodgkin Lymphoma	0.04	(0.04, 0.04)	0.04	(0.04, 0.04)	0.03	(0.02, 0.03)
Non-Hodgkin Lymphoma	0.82	(0.81, 0.83)	0.86	(0.85, 0.87)	0.47	(0.45, 0.50)
Myeloma	0.48	(0.47, 0.48)	0.46	(0.45, 0.46)	0.70	(0.68, 0.73)
Leukemia	0.99	(0.98, 1.00)	1.04	(1.03, 1.05)	0.67	(0.64, 0.69)
Acute Lymphocytic Leukemia	0.05	(0.05, 0.05)	0.05	(0.05, 0.05)	0.03	(0.02, 0.03)
Chronic Lymphocytic Leukemia	0.22	(0.21, 0.22)	0.23	(0.23, 0.24)	0.15	(0.13, 0.16)
Acute Myeloid Leukemia	0.39	(0.39, 0.40)	0.41	(0.40, 0.42)	0.25	(0.23, 0.26)
Chronic Myeloid Leukemia	0.05	(0.04, 0.05)	0.05	(0.05, 0.05)	0.03	(0.03, 0.04)

Devcan Version 6.7.6, April 2018, National Cancer Institute (<https://surveillance.cancer.gov/devcan/>).
Source: NCHS public use data file for the total US.
A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table 1.19 - continued

Lifetime Risk (Percent) of Dying from Cancer by Site and Race/Ethnicity
Males, Total U.S., 2013-2015

Site	Asian/Pacific Islanders	American Indian/ Alaska Natives ^a	Hispanics ^b
	Percent (95% C.I.)	Percent (95% C.I.)	Percent (95% C.I.)
All Sites	20.40 (20.07, 20.74)	17.48 (16.83, 18.18)	19.49 (19.28, 19.70)
Oral Cavity and Pharynx	0.43 (0.39, 0.48)	0.32 (0.24, 0.50)	0.32 (0.29, 0.35)
Esophagus	0.43 (0.39, 0.49)	0.63 (0.52, 0.84)	0.48 (0.45, 0.51)
Stomach	1.14 (1.06, 1.24)	0.62 (0.51, 0.82)	0.88 (0.83, 0.93)
Colon and Rectum	1.98 (1.87, 2.10)	2.03 (1.80, 2.34)	2.00 (1.93, 2.07)
Liver and Intrahepatic Bile Duct	2.02 (1.92, 2.12)	1.37 (1.22, 1.61)	1.64 (1.59, 1.70)
Pancreas	1.42 (1.33, 1.52)	0.97 (0.82, 1.20)	1.27 (1.21, 1.33)
Larynx	0.11 (0.09, 0.15)	0.15 (0.09, 0.32)	0.20 (0.18, 0.23)
Lung and Bronchus	5.12 (4.95, 5.30)	4.18 (3.86, 4.58)	3.61 (3.52, 3.71)
Melanoma of the Skin	0.05 (0.04, 0.08)	0.08 (0.04, 0.24)	0.13 (0.11, 0.15)
Breast	0.02 (0.01, 0.05)	0.00 (0.00, 0.17)	0.02 (0.02, 0.04)
Prostate	2.13 (1.99, 2.28)	2.05 (1.78, 2.40)	2.77 (2.67, 2.88)
Testis	0.00 (0.00, 0.03)	0.03 (0.02, 0.20)	0.02 (0.02, 0.04)
Urinary Bladder	0.66 (0.58, 0.75)	0.36 (0.24, 0.57)	0.68 (0.63, 0.74)
Kidney and Renal Pelvis	0.45 (0.40, 0.52)	0.83 (0.70, 1.06)	0.66 (0.63, 0.71)
Brain and Other Nervous System	0.32 (0.28, 0.37)	0.24 (0.17, 0.41)	0.40 (0.37, 0.43)
Thyroid	0.08 (0.06, 0.11)	0.06 (0.03, 0.23)	0.07 (0.06, 0.09)
Hodgkin Lymphoma	0.03 (0.02, 0.05)	0.03 (0.01, 0.20)	0.06 (0.05, 0.08)
Non-Hodgkin Lymphoma	0.86 (0.79, 0.94)	0.57 (0.41, 0.81)	0.84 (0.79, 0.89)
Myeloma	0.30 (0.27, 0.35)	0.40 (0.27, 0.62)	0.46 (0.43, 0.50)
Leukemia	0.78 (0.71, 0.86)	0.51 (0.39, 0.73)	0.81 (0.77, 0.86)
Acute Lymphocytic Leukemia	0.04 (0.03, 0.07)	0.05 (0.03, 0.21)	0.08 (0.07, 0.09)
Chronic Lymphocytic Leukemia	0.06 (0.04, 0.08)	0.09 (0.02, 0.30)	0.12 (0.10, 0.15)
Acute Myeloid Leukemia	0.42 (0.37, 0.47)	0.20 (0.14, 0.37)	0.31 (0.29, 0.34)
Chronic Myeloid Leukemia	0.03 (0.01, 0.06)	0.02 (0.01, 0.19)	0.05 (0.04, 0.07)

Devcan Version 6.7.6, April 2018, National Cancer Institute (<https://surveillance.cancer.gov/devcan/>).

Source: NCHS public use data file for the total US.

^a Underlying mortality data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian Pacific Islanders, and American Indians/Alaska Natives. A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table 1.20

Lifetime Risk (Percent) of Dying from Cancer by Site and Race/Ethnicity
Females, Total U.S., 2013-2015

Site	All Races	Whites	Blacks
	Percent (95% C.I.)	Percent (95% C.I.)	Percent (95% C.I.)
All Sites	18.53 (18.50, 18.57)	18.67 (18.63, 18.71)	18.73 (18.61, 18.85)
Oral Cavity and Pharynx	0.18 (0.18, 0.18)	0.18 (0.18, 0.19)	0.15 (0.14, 0.16)
Esophagus	0.20 (0.19, 0.20)	0.20 (0.20, 0.21)	0.20 (0.19, 0.21)
Stomach	0.30 (0.30, 0.31)	0.27 (0.26, 0.27)	0.48 (0.46, 0.51)
Colon and Rectum	1.71 (1.70, 1.72)	1.68 (1.66, 1.69)	1.96 (1.92, 2.01)
Liver and Intrahepatic Bile Duct	0.52 (0.52, 0.53)	0.50 (0.49, 0.51)	0.55 (0.53, 0.57)
Pancreas	1.34 (1.33, 1.35)	1.32 (1.31, 1.33)	1.54 (1.50, 1.58)
Larynx	0.05 (0.04, 0.05)	0.05 (0.04, 0.05)	0.06 (0.05, 0.07)
Lung and Bronchus	4.62 (4.60, 4.64)	4.80 (4.78, 4.82)	3.91 (3.85, 3.97)
Melanoma of the Skin	0.20 (0.20, 0.21)	0.23 (0.23, 0.24)	0.04 (0.03, 0.05)
Breast	2.59 (2.58, 2.61)	2.56 (2.54, 2.57)	3.12 (3.07, 3.17)
Cervix Uteri	0.22 (0.22, 0.23)	0.21 (0.20, 0.21)	0.36 (0.34, 0.37)
Corpus and Uterus, NOS	0.61 (0.61, 0.62)	0.57 (0.56, 0.58)	0.99 (0.96, 1.02)
Ovary	0.90 (0.89, 0.91)	0.93 (0.92, 0.94)	0.72 (0.69, 0.74)
Urinary Bladder	0.34 (0.34, 0.35)	0.35 (0.34, 0.35)	0.34 (0.32, 0.36)
Kidney and Renal Pelvis	0.33 (0.33, 0.34)	0.34 (0.34, 0.35)	0.30 (0.28, 0.31)
Brain and Other Nervous System	0.42 (0.41, 0.42)	0.45 (0.45, 0.46)	0.23 (0.22, 0.25)
Thyroid	0.07 (0.07, 0.07)	0.07 (0.06, 0.07)	0.08 (0.07, 0.09)
Hodgkin Lymphoma	0.03 (0.03, 0.03)	0.03 (0.03, 0.03)	0.02 (0.02, 0.03)
Non-Hodgkin Lymphoma	0.64 (0.64, 0.65)	0.67 (0.66, 0.68)	0.40 (0.39, 0.42)
Myeloma	0.38 (0.38, 0.39)	0.35 (0.34, 0.35)	0.71 (0.68, 0.73)
Leukemia	0.69 (0.68, 0.70)	0.72 (0.71, 0.73)	0.52 (0.50, 0.55)
Acute Lymphocytic Leukemia	0.04 (0.03, 0.04)	0.04 (0.04, 0.04)	0.03 (0.02, 0.03)
Chronic Lymphocytic Leukemia	0.14 (0.14, 0.15)	0.15 (0.14, 0.15)	0.10 (0.09, 0.11)
Acute Myeloid Leukemia	0.28 (0.27, 0.28)	0.29 (0.29, 0.30)	0.20 (0.19, 0.21)
Chronic Myeloid Leukemia	0.03 (0.03, 0.04)	0.03 (0.03, 0.04)	0.03 (0.02, 0.03)

Devcan Version 6.7.6, April 2018, National Cancer Institute (<https://surveillance.cancer.gov/devcan/>).
Source: NCHS public use data file for the total US.
A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table 1.20 - continued

Lifetime Risk (Percent) of Dying from Cancer by Site and Race/Ethnicity
Females, Total U.S., 2013-2015

Site	Asian/Pacific Islanders	American Indian/ Alaska Natives ^a	Hispanics ^b
	Percent (95% C.I.)	Percent (95% C.I.)	Percent (95% C.I.)
All Sites	16.30 (16.03, 16.57)	15.62 (15.01, 16.26)	15.61 (15.45, 15.78)
Oral Cavity and Pharynx	0.23 (0.19, 0.27)	0.12 (0.07, 0.24)	0.15 (0.13, 0.18)
Esophagus	0.13 (0.11, 0.16)	0.15 (0.09, 0.26)	0.14 (0.12, 0.16)
Stomach	0.84 (0.77, 0.92)	0.45 (0.34, 0.61)	0.62 (0.59, 0.66)
Colon and Rectum	1.74 (1.64, 1.84)	1.68 (1.46, 1.94)	1.57 (1.51, 1.63)
Liver and Intrahepatic Bile Duct	1.11 (1.04, 1.18)	0.88 (0.74, 1.06)	0.98 (0.94, 1.02)
Pancreas	1.47 (1.39, 1.56)	1.03 (0.87, 1.25)	1.33 (1.28, 1.38)
Larynx	0.02 (0.01, 0.03)	0.02 (0.01, 0.12)	0.02 (0.01, 0.03)
Lung and Bronchus	3.42 (3.29, 3.55)	3.64 (3.36, 3.97)	2.27 (2.20, 2.34)
Melanoma of the Skin	0.06 (0.04, 0.09)	0.07 (0.02, 0.19)	0.09 (0.08, 0.11)
Breast	1.79 (1.71, 1.89)	1.75 (1.52, 2.02)	2.08 (2.02, 2.15)
Cervix Uteri	0.24 (0.21, 0.27)	0.22 (0.16, 0.33)	0.30 (0.28, 0.32)
Corpus and Uterus, NOS	0.43 (0.40, 0.48)	0.38 (0.28, 0.53)	0.56 (0.53, 0.60)
Ovary	0.68 (0.62, 0.74)	0.73 (0.61, 0.90)	0.77 (0.74, 0.81)
Urinary Bladder	0.26 (0.22, 0.32)	0.24 (0.16, 0.38)	0.27 (0.24, 0.30)
Kidney and Renal Pelvis	0.22 (0.19, 0.26)	0.44 (0.35, 0.59)	0.40 (0.37, 0.43)
Brain and Other Nervous System	0.30 (0.26, 0.34)	0.24 (0.18, 0.36)	0.35 (0.32, 0.37)
Thyroid	0.13 (0.11, 0.16)	0.08 (0.03, 0.20)	0.11 (0.10, 0.13)
Hodgkin Lymphoma	0.02 (0.01, 0.04)	0.03 (0.01, 0.13)	0.05 (0.04, 0.06)
Non-Hodgkin Lymphoma	0.62 (0.57, 0.68)	0.47 (0.36, 0.63)	0.64 (0.61, 0.68)
Myeloma	0.24 (0.21, 0.28)	0.38 (0.29, 0.53)	0.37 (0.35, 0.40)
Leukemia	0.46 (0.42, 0.51)	0.43 (0.32, 0.59)	0.57 (0.54, 0.60)
Acute Lymphocytic Leukemia	0.03 (0.02, 0.06)	0.06 (0.03, 0.15)	0.06 (0.05, 0.07)
Chronic Lymphocytic Leukemia	0.03 (0.02, 0.05)	0.04 (0.01, 0.15)	0.07 (0.06, 0.09)
Acute Myeloid Leukemia	0.24 (0.21, 0.28)	0.19 (0.13, 0.31)	0.22 (0.20, 0.24)
Chronic Myeloid Leukemia	0.02 (0.01, 0.05)	0.00 (0.00, 0.10)	0.03 (0.02, 0.04)

Devcan Version 6.7.6, April 2018, National Cancer Institute (<https://surveillance.cancer.gov/devcan/>).

Source: NCHS public use data file for the total US.

^a Underlying mortality data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian Pacific Islanders, and American Indians/Alaska Natives. A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table 1.21
U.S. and SEER Death Rates by Primary Cancer Site and Race/Ethnicity, 2011-2015

Site		Total United States ^a							SEER 18 Areas ^{ab}						
		Total	White	Black	AI/AN ^c	API ^d	Hisp ^e	W-NHisp ^e	Total	White	Black	AI/AN ^c	API ^d	Hisp ^e	W-NHisp ^e
All Sites	Both Sexes	163.5	163.8	189.8	150.4	101.3	114.6	167.8	157.7	160.4	189.6	132.8	106.8	115.1	166.3
	Male	196.7	196.4	239.9	181.4	120.4	140.0	200.7	188.7	191.0	237.7	162.0	128.1	138.8	197.5
	Female	139.5	140.0	159.0	127.6	87.7	96.7	143.7	135.6	138.3	159.8	111.5	91.7	98.5	143.7
Oral Cavity & Pharynx	Both Sexes	2.5	2.5	2.8	2.2	2.0	1.5	2.6	2.5	2.5	2.9	1.8	2.1	1.5	2.6
	Male	3.9	3.8	4.8	3.7	3.0	2.4	3.9	3.8	3.8	4.6	2.9	3.3	2.4	4.0
	Female	1.3	1.3	1.3	1.0	1.1	0.8	1.4	1.3	1.3	1.6	-	1.2	0.8	1.4
Esophagus	Both Sexes	4.0	4.3	3.5	3.5	1.6	2.1	4.5	3.7	3.9	3.4	2.8	1.6	2.2	4.2
	Male	7.2	7.6	5.8	5.9	2.8	3.9	8.0	6.5	7.0	5.7	4.4	2.9	4.0	7.5
	Female	1.5	1.5	1.8	1.6	0.7	0.8	1.5	1.3	1.4	1.7	1.6	0.6	0.8	1.5
Stomach	Both Sexes	3.2	2.8	5.7	5.2	5.3	5.1	2.4	3.6	3.1	5.8	6.6	5.4	5.6	2.6
	Male	4.3	3.7	8.3	7.3	6.8	6.7	3.4	4.8	4.2	8.3	9.5	6.9	7.0	3.6
	Female	2.3	2.0	3.9	3.5	4.2	4.0	1.7	2.7	2.3	4.0	4.2	4.4	4.5	1.8
Colon & Rectum	Both Sexes	14.5	14.1	19.4	16.5	10.1	11.4	14.3	14.1	13.9	19.7	15.9	10.8	11.2	14.2
	Male	17.3	16.8	24.4	20.2	12.0	14.6	16.9	16.8	16.4	24.9	20.0	13.1	14.5	16.6
	Female	12.2	11.9	16.0	13.6	8.6	9.0	12.1	12.0	11.8	16.2	12.8	9.0	8.7	12.2
Liver & Intrahepatic Bile Duct	Both Sexes	6.4	6.0	8.3	10.6	9.5	9.1	5.6	6.9	6.4	8.3	8.5	9.8	9.6	5.7
	Male	9.4	8.7	13.2	14.8	14.0	13.0	8.2	10.1	9.3	13.1	11.2	14.5	13.7	8.4
	Female	3.8	3.6	4.6	7.0	6.0	5.9	3.4	4.1	3.8	4.6	6.1	6.0	6.1	3.4
Pancreas	Both Sexes	10.9	10.8	13.3	8.8	7.7	8.5	11.0	10.8	10.9	13.1	8.3	8.4	8.8	11.1
	Male	12.6	12.6	14.8	9.6	8.3	9.5	12.8	12.4	12.5	14.5	9.5	9.2	9.6	12.9
	Female	9.5	9.4	12.2	8.0	7.3	7.7	9.5	9.5	9.5	11.9	7.3	7.7	8.2	9.6
Larynx	Both Sexes	1.0	1.0	1.7	0.7	0.4	0.7	1.0	0.9	0.9	1.5	0.6	0.3	0.6	0.9
	Male	1.8	1.7	3.3	1.4	0.7	1.5	1.7	1.6	1.5	3.0	-	0.7	1.2	1.6
	Female	0.4	0.4	0.5	-	0.1	0.1	0.4	0.3	0.3	0.5	-	0.1	0.1	0.4
Lung & Bronchus	Both Sexes	43.4	44.1	46.2	36.7	23.3	18.8	46.6	39.0	40.2	45.1	26.2	24.3	17.6	43.7
	Male	53.8	53.9	65.1	45.0	31.0	26.4	56.3	48.0	48.2	63.2	35.4	33.0	23.7	51.8
	Female	35.4	36.6	33.5	30.6	17.7	13.3	39.0	32.3	34.1	33.0	19.3	18.0	13.1	37.5
Melanoma of the Skin	Both Sexes	2.6	3.0	0.4	0.8	0.3	0.7	3.3	2.5	3.0	0.4	0.8	0.4	0.7	3.4
	Male	3.9	4.5	0.5	1.1	0.4	1.0	4.8	3.7	4.5	0.4	-	0.4	0.9	5.1
	Female	1.6	1.9	0.3	0.5	0.3	0.6	2.1	1.5	1.8	0.3	-	0.3	0.5	2.1
Breast	Female	20.9	20.3	28.6	14.3	11.3	14.2	20.8	20.8	29.6	13.2	12.4	14.2	21.7	

^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b The SEER 18 areas are San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG.

^c Rates for American Indian/Alaska Native (AI/AN) are based on the CHSDA (Contract Health Service Delivery Area) counties.

^d Asian/Pacific Islander.

^e Hispanic (Hisp) and White Non-Hispanic (W-NHisp) are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives.

- Statistic could not be calculated due to less than 16 cases in the time interval.

Table 1.21 - continued
 U.S. and SEER Death Rates by Primary Cancer Site and Race/Ethnicity, 2011-2015

Site		Total United States ^a							SEER 18 Areas ^{ab}						
		Total	White	Black	AI/AN ^c	API ^d	Hisp ^e	W-NHisp ^e	Total	White	Black	AI/AN ^c	API ^d	Hisp ^e	W-NHisp ^e
Cervix	Female	2.3	2.2	3.7	2.6	1.8	2.6	2.1	2.2	2.1	3.3	2.2	1.9	2.5	2.0
Corpus & Uterus, NOS	Female	4.6	4.3	8.3	3.6	2.9	3.8	4.3	4.6	4.4	8.2	3.9	3.2	3.7	4.4
Ovary	Female	7.2	7.5	6.3	6.3	4.3	5.3	7.6	7.2	7.7	6.4	6.4	4.4	5.6	7.9
Prostate	Male	19.5	18.2	39.9	19.7	8.7	16.1	18.2	19.8	19.1	40.4	17.3	9.6	16.6	19.2
Testis	Male	0.2	0.3	0.1	0.4	0.1	0.3	0.3	0.3	0.3	0.2	-	0.1	0.3	0.3
Urinary Bladder	Both Sexes	4.4	4.6	3.5	2.3	1.7	2.3	4.8	4.3	4.6	3.8	2.3	1.7	2.3	4.9
	Male	7.6	8.0	5.3	3.6	2.9	3.9	8.3	7.4	8.0	5.7	3.4	3.0	3.7	8.6
	Female	2.2	2.2	2.4	1.4	0.9	1.3	2.3	2.1	2.2	2.6	1.5	0.9	1.3	2.3
Kidney & Renal Pelvis	Both Sexes	3.8	4.0	3.7	6.0	1.8	3.5	4.0	3.7	3.9	3.7	5.4	2.0	3.5	3.9
	Male	5.6	5.8	5.5	8.4	2.6	5.0	5.8	5.4	5.7	5.5	7.7	3.1	5.1	5.7
	Female	2.4	2.5	2.4	4.1	1.1	2.3	2.5	2.3	2.4	2.3	3.7	1.3	2.3	2.4
Brain & Nervous System	Both Sexes	4.4	4.8	2.6	2.5	2.1	2.9	5.0	4.3	4.8	2.7	2.3	2.2	3.0	5.1
	Male	5.3	5.8	3.2	3.0	2.5	3.4	6.1	5.3	5.9	3.4	2.9	2.6	3.6	6.3
	Female	3.5	3.9	2.1	2.0	1.8	2.5	4.0	3.5	3.9	2.2	1.7	1.9	2.5	4.1
Thyroid	Both Sexes	0.5	0.5	0.5	0.5	0.6	0.6	0.5	0.5	0.5	0.5	-	0.7	0.7	0.5
	Male	0.5	0.5	0.4	-	0.5	0.5	0.5	0.5	0.5	0.4	-	0.6	0.5	0.5
	Female	0.5	0.5	0.6	0.5	0.7	0.7	0.4	0.5	0.5	0.5	-	0.8	0.7	0.5
Hodgkin Lymphoma	Both Sexes	0.3	0.3	0.3	-	0.1	0.4	0.3	0.3	0.4	0.3	-	0.2	0.4	0.3
	Male	0.4	0.4	0.4	-	0.2	0.5	0.4	0.4	0.4	0.4	-	0.2	0.5	0.4
	Female	0.3	0.3	0.2	-	0.1	0.3	0.3	0.3	0.3	0.2	-	0.1	0.3	0.3
Non-Hodgkin Lymphoma	Both Sexes	5.7	6.0	4.2	4.4	3.9	4.8	6.0	5.7	6.0	4.2	4.2	4.1	5.0	6.1
	Male	7.4	7.7	5.4	5.6	5.0	6.1	7.8	7.4	7.8	5.4	5.3	5.1	6.3	7.9
	Female	4.5	4.6	3.4	3.4	3.2	3.9	4.7	4.4	4.7	3.3	3.3	3.3	4.0	4.7
Myeloma	Both Sexes	3.3	3.1	6.2	3.0	1.6	2.8	3.1	3.3	3.1	6.4	3.0	1.7	2.8	3.2
	Male	4.2	4.0	7.5	3.4	2.0	3.4	4.0	4.2	4.1	7.5	3.1	2.1	3.5	4.2
	Female	2.7	2.4	5.5	2.7	1.3	2.3	2.4	2.6	2.4	5.6	3.0	1.4	2.3	2.4
Leukemia	Both Sexes	6.7	6.9	5.6	4.3	3.7	4.8	7.0	6.4	6.8	5.6	3.6	4.0	4.8	6.9
	Male	9.0	9.3	7.4	5.5	4.9	6.0	9.5	8.5	9.0	7.2	5.1	5.2	6.0	9.3
	Female	5.0	5.2	4.5	3.3	2.9	3.9	5.2	4.9	5.2	4.5	2.4	3.1	3.9	5.2

^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b The SEER 18 areas are San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG.

^c Rates for American Indian/Alaska Native (AI/AN) are based on the CHSDA (Contract Health Service Delivery Area) counties.

^d Asian/Pacific Islander.

^e Hispanic (Hisp) and White Non-Hispanic (W-NHisp) are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives.

- Statistic could not be calculated due to less than 16 cases in the time interval.

Table 1.22
U.S. Prevalence Counts, Invasive Cancers Only, January 1, 2015^a
Using Different Tumor Inclusion Criteria^b

Site	Sex	5-Year Limited Duration			40-year Limited Duration	
		1st Invasive Tumor Ever ^c	1st Per Site in Previous 40 Years ^d	1st Per Site in Previous 5 Years ^e	1st Invasive Tumor Ever ^c	1st Per Site in Previous 40 Years ^d
All Sites	Both Sexes	4,722,844	4,824,176	5,364,403	14,564,667	14,803,473
	Male	2,336,741	2,376,359	2,637,853	6,886,073	6,969,613
	Female	2,386,103	2,447,817	2,726,550	7,678,594	7,833,860
Oral Cavity & Pharynx	Both Sexes	119,650	140,282	146,087	315,830	351,683
	Male	85,201	99,042	102,445	213,376	235,817
	Female	34,449	41,240	43,642	102,454	115,866
Esophagus	Both Sexes	22,718	28,397	28,552	39,725	47,178
	Male	17,684	22,100	22,194	30,875	36,584
	Female	5,034	6,297	6,358	8,850	10,594
Stomach	Both Sexes	40,050	48,431	49,134	83,535	96,852
	Male	23,823	29,145	29,511	47,851	55,915
	Female	16,227	19,286	19,623	35,684	40,937
Colon & Rectum	Both Sexes	388,393	447,554	457,029	1,184,519	1,317,472
	Male	200,410	230,711	235,400	592,981	656,924
	Female	187,983	216,843	221,629	591,538	660,548
Liver & Intrahepatic Bile Duct	Both Sexes	43,476	50,595	50,790	62,989	71,436
	Male	32,091	36,509	36,632	45,370	50,472
	Female	11,385	14,086	14,158	17,619	20,964
Pancreas	Both Sexes	41,541	51,213	51,287	56,595	68,335
	Male	21,365	26,612	26,669	28,337	34,642
	Female	20,176	24,601	24,618	28,258	33,693
Larynx	Both Sexes	31,888	37,523	38,098	87,909	98,750
	Male	25,677	30,180	30,662	70,763	79,299
	Female	6,211	7,343	7,436	17,146	19,451
Lung & Bronchus	Both Sexes	241,884	314,788	327,383	427,053	534,600
	Male	107,976	143,225	148,287	186,018	235,174
	Female	133,908	171,563	179,096	241,035	299,426
Melanoma of the Skin	Both Sexes	311,301	357,482	378,631	1,093,856	1,188,494
	Male	169,064	198,951	212,770	546,187	602,574
	Female	142,237	158,531	165,861	547,669	585,920
Breast	Female	919,418	1,000,641	1,067,300	3,186,386	3,383,211
Cervix	Female	39,276	41,471	41,584	222,360	228,508
Corpus & Uterus, NOS	Female	199,512	224,764	224,949	648,910	711,315
Ovary ^f	Female	61,387	70,296	70,355	188,419	211,175

^a U.S. 2015 cancer prevalence counts are based on 2015 cancer prevalence proportions from the SEER 9 registries and 1/1/2015 U.S. population estimates based on the average of 2014 and 2015 population estimates from the U.S. Bureau of the Census.

^b Prevalence estimates are ambiguous for those with multiple cancers, unless the tumor inclusion criteria are understood. Depending on the application, different inclusion criteria may be appropriate. This table provides three different methods of tumor inclusion:

^c (c) First invasive tumor ever

^d (d) First invasive tumor for each cancer site diagnosed during the previous 40 years (1975-2014)

^e (e) First invasive tumor for each cancer site diagnosed during the previous 5 years (2010-2014)

For definitions (d) and (e) all sites is treated as a separate cancer "site".

Consider a woman who had three invasive cancers: Melanoma in 1981; Breast cancer in 2010; Melanoma in 2011.

In method (c) the melanoma is the woman's first cancer, and is counted for the melanoma and all sites 40-year limited duration prevalence. For 5-year limited duration prevalence, the woman is not counted at all since her first cancer occurred more than 5 years prior to 1/1/2015.

In method (d) the 1981 melanoma is counted for the melanoma and all sites 40-year limited duration prevalence. The 2010 breast cancer is counted for the breast 5-year and 40-year limited duration prevalence.

In method (e) the 2010 breast cancer is counted for the breast cancer and all sites 5-year limited duration prevalence. The 2011 melanoma is counted for 5-year limited duration prevalence for melanoma.

^f Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

Table 1.22 - continued
 U.S. Prevalence Counts, Invasive Cancers Only, January 1, 2015^a
 Using Different Tumor Inclusion Criteria^b

Site	Sex	5-Year Limited Duration			40-year Limited Duration	
		1st Invasive Tumor Ever ^c	1st Per Site in Previous 40 Years ^d	1st Per Site in Previous 5 Years ^e	1st Invasive Tumor Ever ^c	1st Per Site in Previous 40 Years ^d
Prostate	Male	904,841	981,660	981,708	2,930,034	3,119,586
Testis	Male	43,676	44,722	45,419	240,638	244,108
Urinary Bladder	Both Sexes	209,289	264,140	270,460	595,422	699,906
	Male	159,522	202,342	207,410	444,468	523,805
	Female	49,767	61,798	63,050	150,954	176,101
Kidney & Renal Pelvis	Both Sexes	164,860	202,123	205,347	421,973	494,409
	Male	103,366	127,961	130,183	254,134	300,381
	Female	61,494	74,162	75,164	167,839	194,028
Brain & Nervous System	Both Sexes	46,283	49,632	50,142	148,219	153,481
	Male	25,950	27,706	28,005	79,027	81,695
	Female	20,333	21,926	22,137	69,192	71,786
Thyroid	Both Sexes	206,116	231,486	232,345	673,927	726,654
	Male	46,591	55,337	55,526	145,745	161,902
	Female	159,525	176,149	176,819	528,182	564,752
Hodgkin Lymphoma	Both Sexes	37,408	39,654	39,732	190,947	196,508
	Male	20,709	22,013	22,057	99,255	102,171
	Female	16,699	17,641	17,675	91,692	94,337
Non-Hodgkin Lymphoma	Both Sexes	212,016	250,895	256,180	603,480	678,222
	Male	114,846	137,105	139,836	319,682	360,297
	Female	97,170	113,790	116,344	283,798	317,925
Myeloma	Both Sexes	64,947	76,977	77,405	107,880	124,483
	Male	35,484	43,060	43,330	58,762	68,975
	Female	29,463	33,917	34,075	49,118	55,508
Leukemia	Both Sexes	132,739	156,843	157,533	359,582	399,967
	Male	77,006	91,685	92,061	204,125	228,132
	Female	55,733	65,158	65,472	155,457	171,835
Acute Lymphocytic Leukemia	Both Sexes	17,562	18,283	18,283	80,053	81,139
	Male	9,392	9,771	9,771	43,576	44,099
	Female	8,170	8,512	8,512	36,477	37,040
Childhood (Ages 0-19)	Both Sexes	67,139	67,259	67,807	360,164	360,164
	Male	34,528	34,605	34,868	184,580	184,580
	Female	32,611	32,654	32,939	175,584	175,584
Kaposi Sarcoma	Both Sexes	7,206	7,837	7,837	28,593	30,123
	Male	6,671	7,226	7,226	27,086	28,436
	Female	535	611	611	1,507	1,687
Mesothelioma	Both Sexes	3,059	4,067	4,079	4,861	5,982
	Male	2,126	2,837	2,837	2,904	3,671
	Female	933	1,230	1,242	1,957	2,311

^a U.S. 2015 cancer prevalence counts are based on 2015 cancer prevalence proportions from the SEER 9 registries and 1/1/2015 U.S. population estimates based on the average of 2014 and 2015 population estimates from the U.S. Bureau of the Census.

^b Prevalence estimates are ambiguous for those with multiple cancers, unless the tumor inclusion criteria are understood. Depending on the application, different inclusion criteria may be appropriate. This table provides three different methods of tumor inclusion:

^c (c) First invasive tumor ever

^d (d) First invasive tumor for each cancer site diagnosed during the previous 40 years (1975-2014)

^e (e) First invasive tumor for each cancer site diagnosed during the previous 5 years (2010-2014)

For definitions (d) and (e) all sites is treated as a separate cancer "site".

Consider a woman who had three invasive cancers: Melanoma in 1981; Breast cancer in 2010; Melanoma in 2011.

In method (c) the melanoma is the woman's first cancer, and is counted for the melanoma and all sites 40-year limited duration prevalence. For 5-year limited duration prevalence, the woman is not counted at all since her first cancer occurred more than 5 years prior to 1/1/2015.

In method (d) the 1981 melanoma is counted for the melanoma and all sites 40-year limited duration prevalence. The 2010 breast cancer is counted for the breast 5-year and 40-year limited duration prevalence.

In method (e) the 2010 breast cancer is counted for the breast cancer and all sites 5-year limited duration prevalence. The 2011 melanoma is counted for 5-year limited duration prevalence for melanoma.

Table 1.23
U.S. Complete Prevalence Counts, Invasive Cancers Only, January 1, 2015^a
By Age at Prevalence

Site/Sex	Age at Prevalence								
	All Ages ^c	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70+
All Sites									
Males	7,032,494	18,799	43,681	92,992	172,866	345,981	942,023	1,888,544	3,527,607
Females	8,079,604	17,434	38,006	96,195	248,166	638,524	1,448,944	2,081,139	3,511,196
Oral Cavity & Pharynx									
Males	240,247	60	336	1,691	3,835	13,583	51,730	81,471	87,541
Females	119,471	96	496	1,605	4,119	8,518	21,670	31,757	51,208
Esophagus									
Males	36,637	0	0	28	231	975	4,943	12,273	18,186
Females	10,647	0	0	11	55	198	1,426	3,121	5,836
Stomach									
Males	56,530	0	23	150	711	2,701	8,186	15,329	29,430
Females	41,385	0	61	172	792	2,373	6,357	9,414	22,216
Colon & Rectum									
Males	661,891	33	250	1,423	7,046	26,892	94,246	161,215	370,786
Females	670,194	0	371	1,716	7,187	25,306	82,985	135,011	417,617
Liver & Intrahep									
Males	50,546	578	624	681	635	1,774	10,577	22,897	12,780
Females	21,444	441	607	449	509	1,106	3,718	7,157	7,456
Pancreas									
Males	34,774	23	46	131	515	1,660	6,226	11,193	14,981
Females	33,841	11	46	330	783	1,891	6,174	9,420	15,186
Larynx									
Males	80,116	0	23	79	241	1,631	10,408	23,663	44,070
Females	19,640	0	0	17	152	770	3,601	6,083	9,016
Lung & Bronchus									
Males	238,904	45	110	427	1,280	5,040	25,457	66,067	140,479
Females	302,131	44	81	398	1,613	6,862	36,368	77,674	179,091
Melanoma of the Skin									
Males	613,293	68	636	5,049	18,308	45,387	105,775	168,413	269,658
Females	608,730	102	784	10,401	36,985	74,100	131,803	152,396	202,159

^a U.S. 2015 cancer prevalence counts are based on 2015 cancer prevalence proportions from the SEER 9 registries (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta) and 1/1/2015 U.S. population estimates based on the average of 2014 and 2015 population estimates from the U.S. Bureau of the Census. Prevalence was calculated using the first invasive tumor for each cancer site diagnosed during the previous 40 years (1975-2014).

^b Cases diagnosed more than 40 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000).

^c Due to rounding, the sum of the age specific estimates may not equal the all ages estimate.

Table 1.23 - continued
 U.S. Complete Prevalence Counts, Invasive Cancers Only, January 1, 2015^a
 By Age at Prevalence

Site/Sex	Age at Prevalence								
	All Ages ^c	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70+
Breast									
Males	19,095	0	0	11	153	624	2,094	5,126	11,088
Females	3,418,124	11	12	3,313	39,205	214,191	604,877	942,597	1,613,918
Cervix									
Females	257,524	0	81	2,063	14,486	40,266	60,508	63,055	77,065
Corpus & Uterus, NOS									
Females	727,200	11	23	844	6,718	27,300	101,936	215,845	374,522
Ovary ^d									
Females	224,940	113	948	3,985	7,585	18,327	46,034	62,323	85,625
Prostate									
Males	3,120,176	57	46	80	339	15,748	231,637	864,374	2,007,896
Urinary Bladder									
Males	529,318	62	105	595	2,505	9,096	43,303	120,318	353,336
Females	179,126	68	52	251	1,033	3,695	14,768	37,488	121,771
Kidney & Renal Pelvis									
Males	306,570	1,572	2,258	3,120	6,122	20,682	54,352	87,265	131,198
Females	198,810	1,676	2,482	3,197	5,496	13,754	33,887	50,099	88,220
Hodgkin Lymphoma									
Males	108,267	186	2,304	10,001	16,774	22,863	25,896	19,002	11,241
Females	100,538	45	1,912	9,703	16,856	22,201	23,482	15,606	10,735
Non-Hodgkin Lymphoma									
Males	364,491	828	4,278	8,341	14,678	30,274	62,974	91,757	151,361
Females	321,551	626	1,720	4,732	10,140	21,355	48,928	80,035	154,015
Myeloma									
Males	69,065	0	0	88	694	3,299	11,384	20,714	32,886
Females	55,668	0	0	67	464	2,560	8,464	16,640	27,472
Leukemia									
Males	230,623	6,548	13,735	14,815	13,747	17,054	29,724	50,174	84,826
Females	175,192	6,088	10,463	13,168	11,932	13,097	21,058	32,759	66,628
Acute Lymphocytic Leuk									
Males	45,907	5,461	11,360	11,466	7,880	5,297	2,407	1,257	780
Females	38,863	5,145	8,877	9,392	6,746	4,468	2,136	1,295	803

^a U.S. 2015 cancer prevalence counts are based on 2015 cancer prevalence proportions from the SEER 9 registries (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta) and 1/1/2015 U.S. population estimates based on the average of 2014 and 2015 population estimates from the U.S. Bureau of the Census. Prevalence was calculated using the first invasive tumor for each cancer site diagnosed during the previous 40 years (1975-2014).

^b Cases diagnosed more than 40 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000).

^c Due to rounding, the sum of the age specific estimates may not equal the all ages estimate.

Table 1.24
Age-Adjusted SEER Incidence Rates and Trends for the Top 15 Cancer Sites^a by Race/Ethnicity
Both Sexes

	All Races		White		Black	
	Rate ^b 2011-2015	APC ^c 2006-2015	Rate ^b 2011-2015	APC ^c 2006-2015	Rate ^b 2011-2015	APC ^c 2006-2015
All Sites	439.2	-1.2*	448.8	-1.2*	453.4	-1.7*
Breast	67.5	0.0	68.1	-0.1	76.4	-5.2*
Lung and Bronchus	54.6	-2.4*	56.1	-2.4*	72.0	0.3
Prostate ^f	51.5	-5.7*	48.9	-6.2*	61.4	-2.6*
Colon and Rectum	39.4	-2.7*	38.8	-2.6*	47.6	-3.3*
Melanoma of the Skin	22.8	1.5*	27.3	1.5*	18.3	0.7
Urinary Bladder	19.5	-1.3*	21.5	-1.3*	15.5	-0.5
Non-Hodgkin Lymphoma	19.4	-0.5*	20.3	-0.7*	14.4	-0.2
Kidney and Renal Pelvis	15.9	0.5*	16.3	0.5*	14.3	2.1*
Thyroid	14.5	3.3*	15.3	3.2*	13.4	0.8
Leukemia	13.8	0.3	14.6	0.2	12.3	-0.8*
Corpus and Uterus, NOS ^f	13.8	1.2*	14.0	1.0*	11.0	0.9
Pancreas	12.6	0.5*	12.6	0.6*	10.5	2.5*
Oral Cavity and Pharynx	11.3	0.6*	11.9	0.9*	10.1	-2.8*
Liver & IBD ^g	8.8	2.5*	8.0	3.1*	9.0	-1.6*
Stomach	7.2	-0.9*	7.1	-0.4	8.9	4.2*
	Asian/Pacific Islander		American Indian/Alaska Native ^d		Hispanic ^e	
	Rate ^b 2011-2015	APC ^c 2006-2015	Rate ^b 2011-2015	APC ^c 2006-2015	Rate ^b 2011-2015	APC ^c 2006-2015
All Sites	297.9	-1.2*	315.1	-0.9*	336.3	-1.1*
Breast	55.3	1.1*	44.4	0.5	50.0	0.5
Lung and Bronchus	35.6	-1.6*	41.8	0.4	40.9	-6.1*
Colon and Rectum	33.0	-2.9*	37.3	-2.7*	33.5	-2.2*
Prostate ^f	25.9	-7.0*	24.3	-6.9*	27.8	-2.5*
Thyroid	14.0	2.6*	17.2	-3.1*	17.5	-0.2
Non-Hodgkin Lymphoma	13.6	0.0	13.5	0.9	15.9	1.0*
Liver & IBD ^g	13.1	-2.0*	11.8	-1.9	13.3	1.7*
Corpus and Uterus, NOS ^f	11.4	1.9*	10.4	-0.2	12.5	4.0*
Stomach	10.3	-3.0*	10.2	0.2	11.9	3.1*
Pancreas	10.0	0.2	9.3	-1.9	11.2	0.2
Urinary Bladder	8.6	-1.5*	9.3	4.4*	10.8	-1.6*
Kidney and Renal Pelvis	8.5	0.7	9.0	1.2	10.6	-0.2
Leukemia	7.8	0.1	8.8	0.2	10.2	-1.6*
Oral Cavity and Pharynx	7.8	0.0	8.3	1.1	6.5	-0.2
Ovary ^{fh}	5.3	-0.4	5.8	4.5	6.2	1.2

Source: SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

^a Top 15 cancer sites selected based on 2011-2015 age-adjusted rates for the race/ethnic group.

^b Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^c The APC is the Annual Percent Change over the time interval.

Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^d Rates for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^e Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives.

Incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

^f The rates for sex-specific cancer sites are calculated using the population for both sexes combined.

^g IBD = Intrahepatic Bile Duct. ONS = Other Nervous System.

^h Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

* The APC is significantly different from zero (p<.05).

- Statistic not shown. Rate based on less than 16 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.

Table 1.25
Age-Adjusted SEER Incidence Rates and Trends for the Top 15 Cancer Sites^a by Race/Ethnicity

			Males					
All Races			White			Black		
	Rate ^b	APC ^c		Rate ^b	APC ^c		Rate ^b	APC ^c
	2011-2015	2006-2015		2011-2015	2006-2015		2011-2015	2006-2015
All Sites	483.0	-2.2*	All Sites	488.5	-2.2*	All Sites	535.0	-3.1*
Prostate	112.6	-6.0*	Prostate	105.7	-6.5*	Prostate	178.3	-5.5*
Lung and Bronchus	63.8	-3.1*	Lung and Bronchus	63.9	-3.1*	Lung and Bronchus	81.2	-3.4*
Colon and Rectum	45.2	-2.8*	Colon and Rectum	44.4	-2.9*	Colon and Rectum	55.5	-3.3*
Urinary Bladder	34.3	-1.5*	Urinary Bladder	37.6	-1.5*	Kidney and Renal Pelvis	25.3	0.7
Melanoma of the Skin	29.8	1.6*	Melanoma of the Skin	35.2	1.6*	Urinary Bladder	20.6	-0.8
Non-Hodgkin Lymphoma	23.6	-0.4*	Non-Hodgkin Lymphoma	24.7	-0.5*	Non-Hodgkin Lymphoma	17.5	-0.1
Kidney and Renal Pelvis	21.7	0.5	Kidney and Renal Pelvis	22.2	0.5	Liver & IBD ^f	17.3	2.6*
Leukemia	17.6	0.3	Leukemia	18.6	0.2	Pancreas	16.9	-0.4
Oral Cavity and Pharynx	17.1	0.8*	Oral Cavity and Pharynx	18.0	1.1*	Myeloma	15.9	0.4
Pancreas	14.4	0.5*	Pancreas	14.4	0.6*	Leukemia	14.0	0.5
Liver & IBD ^f	13.6	2.3*	Liver & IBD ^f	12.2	2.8*	Oral Cavity and Pharynx	14.0	-1.8*
Stomach	9.8	-1.2*	Stomach	8.9	-0.9*	Stomach	13.6	-3.3*
Myeloma	8.4	1.2*	Brain and ONS ^f	8.3	-0.5*	Larynx	8.3	-3.2*
Brain and ONS ^f	7.5	-0.6*	Myeloma	7.9	1.2*	Esophagus	6.4	-5.3*
Thyroid	7.3	3.5*	Thyroid	7.8	3.4*	Brain and ONS ^f	4.8	-0.5
Asian/Pacific Islander			American Indian/Alaska Native^d			Hispanic^e		
	Rate ^b	APC ^c		Rate ^b	APC ^c		Rate ^b	APC ^c
	2011-2015	2006-2015		2011-2015	2006-2015		2011-2015	2006-2015
All Sites	303.5	-2.6*	All Sites	325.6	-1.7*	All Sites	362.9	-2.6*
Prostate	59.1	-7.2*	Prostate	54.8	-7.1*	Prostate	91.8	-6.5*
Lung and Bronchus	45.9	-2.3*	Lung and Bronchus	45.4	-2.4*	Colon and Rectum	39.7	-2.7*
Colon and Rectum	39.3	-2.7*	Colon and Rectum	45.1	0.2	Lung and Bronchus	34.1	-3.5*
Liver & IBD ^f	20.2	-1.8*	Kidney and Renal Pelvis	22.6	-3.3*	Kidney and Renal Pelvis	20.7	0.5
Non-Hodgkin Lymphoma	16.7	0.1	Liver & IBD ^f	18.5	0.3	Non-Hodgkin Lymphoma	20.2	-0.5
Urinary Bladder	15.0	-1.9*	Urinary Bladder	14.7	-1.5	Liver & IBD ^f	19.8	1.3*
Stomach	13.6	-3.0*	Stomach	13.7	-0.1	Urinary Bladder	18.9	-1.9*
Kidney and Renal Pelvis	12.2	1.1	Non-Hodgkin Lymphoma	12.9	-2.5	Leukemia	12.8	-0.3
Oral Cavity and Pharynx	11.1	-0.1	Oral Cavity and Pharynx	12.3	0.9	Stomach	12.8	-2.7*
Pancreas	11.0	0.2	Pancreas	11.3	-	Pancreas	12.0	-0.2
Leukemia	9.7	0.1	Leukemia	10.6	2.7	Oral Cavity and Pharynx	9.6	-0.3
Thyroid	7.0	4.4*	Myeloma	6.2	-	Myeloma	7.6	1.2
Myeloma	4.9	0.7	Testis	5.6	5.2	Brain and ONS ^f	5.7	-0.6
Brain and ONS ^f	4.3	-0.5	Esophagus	5.1	-	Testis	5.3	2.6*
Esophagus	3.3	-2.3	Melanoma of the Skin	5.0	-	Thyroid	5.3	3.5*

Source: SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

^a Top 15 cancer sites selected based on 2011-2015 age-adjusted rates for the race/ethnic group.

^b Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^c The APC is the Annual Percent Change over the time interval.

Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^d Rates for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^e Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives.

Incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

^f IBD = Intrahepatic Bile Duct. ONS = Other Nervous System.

* The APC is significantly different from zero (p<.05).

- Statistic not shown. Rate based on less than 16 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.

Table 1.26
Age-Adjusted SEER Incidence Rates and Trends for the Top 15 Cancer Sites^a by Race/Ethnicity

Females

All Races	Rate ^b		APC ^c	All Sites	Rate ^b		APC ^c	All Sites	Rate ^b		APC ^c
	2011-2015	2006-2015			2011-2015	2006-2015			2011-2015	2006-2015	
All Sites	409.9	-0.2*		All Sites	423.1	-0.2*		All Sites	397.8	-0.4*	
Breast	126.0	0.2		Breast	128.6	0.0		Breast	126.9	0.4	
Lung and Bronchus	47.8	-1.8*		Lung and Bronchus	50.2	-1.7*		Lung and Bronchus	47.9	-1.8*	
Colon and Rectum	34.5	-2.6*		Colon and Rectum	34.0	-2.4*		Colon and Rectum	41.9	-3.4*	
Corpus and Uterus, NOS	26.0	1.4*		Corpus and Uterus, NOS	26.6	1.2*		Corpus and Uterus, NOS	25.4	2.3*	
Thyroid	21.4	3.3*		Thyroid	22.8	3.2*		Pancreas	14.3	-0.4	
Melanoma of the Skin	17.7	1.3*		Melanoma of the Skin	21.5	1.3*		Thyroid	13.4	4.6*	
Non-Hodgkin Lymphoma	15.9	-0.7*		Non-Hodgkin Lymphoma	16.8	-0.8*		Kidney and Renal Pelvis	12.8	0.6	
Ovary ^g	11.6	-1.8*		Ovary ^g	12.1	-1.9*		Non-Hodgkin Lymphoma	12.1	-0.2	
Pancreas	11.2	0.5*		Leukemia	11.4	0.1		Myeloma	11.6	0.9	
Kidney and Renal Pelvis	10.9	0.3		Kidney and Renal Pelvis	11.3	0.4		Ovary ^g	9.3	-1.4*	
Leukemia	10.8	0.2		Pancreas	11.1	0.6*		Leukemia	9.0	1.3*	
Urinary Bladder	8.3	-1.4*		Urinary Bladder	8.9	-1.4*		Cervix Uteri	8.4	-2.6*	
Cervix Uteri	7.4	-1.4*		Cervix Uteri	7.4	-1.2*		Stomach	7.7	-2.6*	
Oral Cavity and Pharynx	6.3	0.1		Oral Cavity and Pharynx	6.5	0.2		Urinary Bladder	6.7	-1.2*	
Brain and ONS ^f	5.4	-0.3		Brain and ONS ^f	6.0	-0.2		Oral Cavity and Pharynx	5.1	-1.3	
Asian/Pacific Islander				American Indian/Alaska Native^d				Hispanic^e			
	Rate ^b	APC ^c		Rate ^b	APC ^c		Rate ^b	APC ^c			
	2011-2015	2006-2015		2011-2015	2006-2015		2011-2015	2006-2015			
All Sites	298.3	-0.1		All Sites	311.4	-0.2		All Sites	322.9	0.3*	
Breast	100.6	1.2*		Breast	82.6	0.6		Breast	93.7	0.6*	
Lung and Bronchus	28.0	-0.8		Colon and Rectum	39.2	0.7		Colon and Rectum	28.6	-1.7*	
Colon and Rectum	28.0	-3.2*		Lung and Bronchus	31.2	-3.2		Lung and Bronchus	23.2	-1.5*	
Corpus and Uterus, NOS	20.8	1.9*		Corpus and Uterus, NOS	19.6	-0.1		Corpus and Uterus, NOS	22.5	3.3*	
Thyroid	20.2	2.2*		Thyroid	14.3	4.9*		Thyroid	19.6	4.3*	
Non-Hodgkin Lymphoma	11.1	-0.1		Kidney and Renal Pelvis	13.0	-2.5		Non-Hodgkin Lymphoma	15.3	0.2	
Ovary ^g	9.6	-0.3		Non-Hodgkin Lymphoma	10.8	-1.6		Kidney and Renal Pelvis	11.9	1.6*	
Pancreas	9.2	0.2		Ovary ^g	9.0	-5.0*		Pancreas	10.5	0.6	
Stomach	7.9	-3.1*		Liver & IBD ^f	8.9	2.0		Ovary ^g	10.4	-1.2	
Liver & IBD ^f	7.4	-2.2*		Cervix Uteri	8.1	1.7		Cervix Uteri	8.9	-3.6*	
Leukemia	6.4	-0.1		Pancreas	7.8	-5.8*		Leukemia	8.8	-0.3	
Cervix Uteri	6.1	-2.2*		Stomach	7.3	0.3		Stomach	8.3	-0.2	
Kidney and Renal Pelvis	5.6	0.0		Leukemia	6.5	-1.0		Liver & IBD ^f	7.7	2.6*	
Oral Cavity and Pharynx	5.1	0.2		Oral Cavity and Pharynx	6.1	-		Myeloma	5.0	1.0	
Urinary Bladder	3.9	-0.1		Myeloma	5.5	-		Urinary Bladder	4.8	-1.4*	

Source: SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

^a Top 15 cancer sites selected based on 2011-2015 age-adjusted rates for the race/ethnic group.

^b Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^c The APC is the Annual Percent Change over the time interval.

Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^d Rates for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^e Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives.

Incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

^f IBD = Intrahepatic Bile Duct. ONS = Other Nervous System.

^g Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

* The APC is significantly different from zero (p<.05).

- Statistic not shown. Rate based on less than 16 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.

Table 1.27
Age-Adjusted U.S. Death Rates and Trends for the Top 15 Cancer Sites^a by Race/Ethnicity
Both Sexes

	All Races		White		Black	
	Rate ^b 2011-2015	APC ^c 2006-2015	Rate ^b 2011-2015	APC ^c 2006-2015	Rate ^b 2011-2015	APC ^c 2006-2015
All Sites	163.5	-1.5*	163.8	-1.4*	189.8	-2.1*
Lung and Bronchus	43.4	-2.6*	44.1	-2.5*	46.2	-3.0*
Colon and Rectum	14.5	-2.4*	14.1	-2.3*	19.4	-3.1*
Breast	11.6	-1.8*	11.2	-1.8*	16.8	-1.7*
Pancreas	10.9	0.1	10.8	0.2*	14.6	-3.8*
Prostate ^f	7.9	-2.4*	7.4	-2.1*	13.3	-0.5*
Leukemia	6.7	-1.3*	6.9	-1.1*	8.3	2.2*
Liver & IBD ^g	6.4	2.6*	6.0	-2.2*	6.2	-0.4
Non-Hodgkin Lymphoma	5.7	-2.3*	6.0	2.8*	5.7	-3.1*
Urinary Bladder	4.4	-0.1	4.8	0.6*	5.6	-1.9*
Brain and ONS ^g	4.4	0.6*	4.6	0.0	4.9	1.8*
Esophagus	4.0	-1.1*	4.3	-0.5*	4.2	-1.7*
Ovary ^f	4.0	-2.6*	4.1	-2.7*	3.7	-1.4*
Kidney and Renal Pelvis	3.8	-0.7*	4.0	-0.5*	3.7	-1.1*
Myeloma	3.3	-0.5	3.1	-0.4	3.5	-0.9
Stomach	3.2	-2.1*	3.0	-0.8	3.5	-4.8*
	Asian/Pacific Islander		American Indian/Alaska Native ^d		Hispanic ^e	
	Rate ^b 2011-2015	APC ^c 2006-2015	Rate ^b 2011-2015	APC ^c 2006-2015	Rate ^b 2011-2015	APC ^c 2006-2015
All Sites	101.3	-1.3*	150.4	-1.1*	114.6	-1.2*
Lung and Bronchus	23.3	-1.9*	36.7	-1.7*	18.8	-2.4*
Colon and Rectum	10.1	-2.0*	16.5	-0.1	11.4	-2.1*
Liver & IBD ^g	9.5	-0.5	10.6	2.7*	9.1	1.3*
Pancreas	7.7	0.1	8.8	-1.3	8.5	-0.1
Breast	6.4	-0.1	8.0	-0.3	7.9	-0.9*
Stomach	5.3	-4.0*	8.0	-1.9*	6.5	-2.7*
Non-Hodgkin Lymphoma	3.9	-1.0*	6.0	-1.6	5.1	-2.0*
Leukemia	3.7	-1.1	5.2	-2.2	4.8	-1.9*
Prostate ^f	3.5	-2.5*	4.4	-1.1	4.8	-0.5
Ovary ^f	2.4	-2.0*	4.3	-1.1	3.5	-0.3
Brain and ONS ^g	2.1	2.8*	3.5	-1.8	3.0	-2.0*
Oral Cavity and Pharynx	2.0	0.3	3.5	0.3	2.9	0.9*
Kidney and Renal Pelvis	1.8	-1.7*	3.0	1.5	2.8	-0.7
Urinary Bladder	1.7	0.5	2.5	-0.1	2.3	-0.4
Corpus and Uterus, NOS ^f	1.6	2.0*	2.3	-0.1	2.1	-1.9*

Source: US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

^a Top 15 cancer sites selected based on 2011-2015 age-adjusted rates for the race/ethnic group.

^b Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^c The APC is the Annual Percent Change over the time interval.

Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^d Rates for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^e Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives.

^f The rates for sex-specific cancer sites are calculated using the population for both sexes combined.

^g IBD = Intrahepatic Bile Duct. ONS = Other Nervous System.

* The APC is significantly different from zero ($p < .05$).

- Statistic not shown. Rate based on less than 16 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.

Table 1.28
Age-Adjusted U.S. Death Rates and Trends for the Top 15 Cancer Sites^a by Race/Ethnicity

Males

All Races	Rate ^b		APC ^c	All Sites	Rate ^b		APC ^c	All Sites	Rate ^b		APC ^c
	2011-2015	2006-2015			2011-2015	2006-2015			2011-2015	2006-2015	
All Sites	196.7	-1.8*		All Sites	196.4	-1.6*		All Sites	239.9	-2.8*	
Lung and Bronchus	53.8	-3.3*		Lung and Bronchus	53.9	-3.2*		Lung and Bronchus	65.1	-3.7*	
Prostate	19.5	-3.1*		Prostate	18.2	-2.9*		Prostate	39.9	-4.5*	
Colon and Rectum	17.3	-2.5*		Colon and Rectum	16.8	-2.4*		Colon and Rectum	24.4	-3.2*	
Pancreas	12.6	0.1		Pancreas	12.6	0.2*		Pancreas	14.8	-0.3	
Liver & IBD ^f	9.4	2.5*		Leukemia	9.3	-1.1*		Liver & IBD ^f	13.2	2.2*	
Leukemia	9.0	-1.3*		Liver & IBD ^f	8.7	2.7*		Stomach	8.3	-3.0*	
Urinary Bladder	7.6	-0.3*		Urinary Bladder	8.0	-0.2		Myeloma	7.5	-1.0*	
Non-Hodgkin Lymphoma	7.4	-2.0*		Non-Hodgkin Lymphoma	7.7	-1.9*		Leukemia	7.4	-2.3*	
Esophagus	7.2	-1.1*		Esophagus	7.6	-0.6*		Esophagus	5.8	-5.3*	
Kidney and Renal Pelvis	5.6	-0.4*		Kidney and Renal Pelvis	5.8	-0.3*		Kidney and Renal Pelvis	5.5	-0.7	
Brain and ONS ^f	5.3	0.5*		Brain and ONS ^f	5.8	0.6*		Non-Hodgkin Lymphoma	5.4	-1.8*	
Stomach	4.3	-2.4*		Melanoma of the Skin	4.5	-0.9*		Urinary Bladder	5.3	-0.4	
Myeloma	4.2	-0.5*		Myeloma	4.0	-0.4		Oral Cavity and Pharynx	4.8	-2.3*	
Melanoma of the Skin	3.9	-1.1*		Oral Cavity and Pharynx	3.8	0.9*		Larynx	3.3	-3.5*	
Oral Cavity and Pharynx	3.9	0.5		Stomach	3.7	-2.4*		Brain and ONS ^f	3.2	1.4	
Asian/Pacific Islander				American Indian/Alaska Native ^d				Hispanic ^e			
	Rate ^b	APC ^c		Rate ^b	APC ^c		Rate ^b	APC ^c			
	2011-2015	2006-2015		2011-2015	2006-2015		2011-2015	2006-2015			
All Sites	120.4	-1.7*		All Sites	181.4	-0.8		All Sites	140.0	-1.6*	
Lung and Bronchus	31.0	-2.7*		Lung and Bronchus	45.0	-1.9		Lung and Bronchus	26.4	-3.3*	
Liver & IBD ^f	14.0	-0.6*		Colon and Rectum	20.2	1.8		Prostate	16.1	-3.2*	
Colon and Rectum	12.0	-1.8*		Prostate	19.7	-0.9		Colon and Rectum	14.6	-1.7*	
Prostate	8.7	-2.7*		Liver & IBD ^f	14.8	2.5*		Liver & IBD ^f	13.0	1.0*	
Pancreas	8.3	-0.3		Pancreas	9.6	-0.5		Pancreas	9.5	-0.4	
Stomach	6.8	-4.4*		Kidney and Renal Pelvis	8.4	-1.4		Stomach	6.7	-2.3*	
Non-Hodgkin Lymphoma	5.0	-1.0		Stomach	7.3	-3.0		Non-Hodgkin Lymphoma	6.1	-1.4*	
Leukemia	4.9	-0.5		Esophagus	5.9	1.5		Leukemia	6.0	-0.5	
Oral Cavity and Pharynx	3.0	0.7		Non-Hodgkin Lymphoma	5.6	0.5		Kidney and Renal Pelvis	5.0	-0.6	
Urinary Bladder	2.9	0.6		Leukemia	5.5	-2.1		Urinary Bladder	3.9	-0.6	
Esophagus	2.8	-2.1*		Oral Cavity and Pharynx	3.7	0.1		Esophagus	3.9	-2.1*	
Kidney and Renal Pelvis	2.6	-1.3		Urinary Bladder	3.6	-2.5		Brain and ONS ^f	3.4	1.1*	
Brain and ONS ^f	2.5	1.8		Myeloma	3.4	-0.3		Myeloma	3.4	-0.6	
Myeloma	2.0	-1.8		Brain and ONS ^f	3.0	1.3		Oral Cavity and Pharynx	2.4	-0.2	
Soft Tissue including Heart	1.0	0.8		Soft Tissue including Heart	1.5	-		Larynx	1.5	-2.1*	

Source: US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

^a Top 15 cancer sites selected based on 2011-2015 age-adjusted rates for the race/ethnic group.

^b Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^c The APC is the Annual Percent Change over the time interval.

Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^d Rates for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^e Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives.

^f IBD = Intrahepatic Bile Duct. ONS = Other Nervous System.

* The APC is significantly different from zero (p<.05).

- Statistic not shown. Rate based on less than 16 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.

Table 1.29
Age-Adjusted U.S. Death Rates and Trends for the Top 15 Cancer Sites^a by Race/Ethnicity

Females

All Races	Rate ^b		APC ^c	All Sites	Rate ^b		APC ^c	All Sites	Rate ^b		APC ^c
	2011-2015	2006-2015			2011-2015	2006-2015			2011-2015	2006-2015	
All Sites	139.5	-1.4*		All Sites	140.0	-1.3*		All Sites	159.0	-1.6*	
Lung and Bronchus	35.4	-2.0*		Lung and Bronchus	36.6	-1.9*		Lung and Bronchus	33.5	-2.2*	
Breast	20.9	-1.6*		Breast	20.3	-1.6*		Breast	28.6	-1.5*	
Colon and Rectum	12.2	-2.5*		Colon and Rectum	11.9	-2.4*		Colon and Rectum	16.0	-3.3*	
Pancreas	9.5	0.0		Pancreas	9.4	0.1		Pancreas	12.2	-0.6*	
Ovary	7.2	-2.4*		Ovary	7.5	-2.5*		Corpus and Uterus, NOS	8.3	2.1*	
Leukemia	5.0	-1.4*		Leukemia	5.2	-1.3*		Ovary	6.3	-1.2*	
Corpus and Uterus, NOS	4.6	1.6*		Non-Hodgkin Lymphoma	4.6	-2.7*		Myeloma	5.5	0.0	
Non-Hodgkin Lymphoma	4.5	-2.7*		Corpus and Uterus, NOS	4.3	1.5*		Liver & IBD ^f	4.6	2.2*	
Liver & IBD ^f	3.8	2.5*		Brain and ONS ^f	3.9	0.6*		Leukemia	4.5	-1.6*	
Brain and ONS ^f	3.5	0.6*		Liver & IBD ^f	3.6	2.7*		Stomach	3.9	-3.5*	
Myeloma	2.7	-0.6		Kidney and Renal Pelvis	2.5	-1.4*		Cervix Uteri	3.7	-2.6*	
Kidney and Renal Pelvis	2.4	-1.5*		Myeloma	2.4	-0.7*		Non-Hodgkin Lymphoma	3.4	-1.5*	
Stomach	2.3	-2.0*		Urinary Bladder	2.2	-0.3		Urinary Bladder	2.4	-2.0*	
Cervix Uteri	2.3	-0.7*		Cervix Uteri	2.2	-0.2		Kidney and Renal Pelvis	2.4	-1.8*	
Urinary Bladder	2.2	-0.6*		Stomach	2.0	-1.7*		Brain and ONS ^f	2.1	1.2	
Asian/Pacific Islander				American Indian/Alaska Native^d				Hispanic^e			
	Rate ^b	APC ^c		Rate ^b	APC ^c		Rate ^b	APC ^c			
	2011-2015	2006-2015		2011-2015	2006-2015		2011-2015	2006-2015			
All Sites	87.7	-1.0*		All Sites	127.6	-1.5*		All Sites	96.7	-1.0*	
Lung and Bronchus	17.7	-0.9*		Lung and Bronchus	30.6	-1.6*		Breast	14.2	-0.8*	
Breast	11.3	-0.1		Breast	14.3	-1.6*		Lung and Bronchus	13.3	-1.2*	
Colon and Rectum	8.6	-2.4*		Colon and Rectum	13.6	-2.1		Colon and Rectum	9.0	-2.8*	
Pancreas	7.3	0.4		Pancreas	8.0	-2.5		Pancreas	7.7	0.1	
Liver & IBD ^f	6.0	-0.4		Liver & IBD ^f	7.0	2.7		Liver & IBD ^f	5.9	1.6*	
Ovary	4.3	-1.8*		Ovary	6.3	-1.4		Ovary	5.3	-1.8*	
Stomach	4.2	-3.7*		Kidney and Renal Pelvis	4.1	-1.8		Stomach	4.0	-1.7*	
Non-Hodgkin Lymphoma	3.2	-0.9		Corpus and Uterus, NOS	3.6	0.5		Non-Hodgkin Lymphoma	3.9	-2.5*	
Corpus and Uterus, NOS	2.9	2.2*		Stomach	3.5	-1.5		Leukemia	3.9	-0.7	
Leukemia	2.9	-1.9		Non-Hodgkin Lymphoma	3.4	-3.1		Corpus and Uterus, NOS	3.8	2.7*	
Brain and ONS ^f	1.8	3.7*		Leukemia	3.3	-0.4		Cervix Uteri	2.6	-2.2*	
Cervix Uteri	1.8	-1.7		Myeloma	2.7	2.7		Brain and ONS ^f	2.5	0.8	
Myeloma	1.3	-0.9		Cervix Uteri	2.6	-4.0		Myeloma	2.3	-1.0	
Oral Cavity and Pharynx	1.1	-0.6		Brain and ONS ^f	2.0	-0.6		Kidney and Renal Pelvis	2.3	-0.1	
Kidney and Renal Pelvis	1.1	-2.4*		Gallbladder	1.7	-5.2		Urinary Bladder	1.3	-0.3	

Source: US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

^a Top 15 cancer sites selected based on 2011-2015 age-adjusted rates for the race/ethnic group.

^b Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^c The APC is the Annual Percent Change over the time interval.

Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^d Rates for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^e Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives.

^f IBD = Intrahepatic Bile Duct. ONS = Other Nervous System.

* The APC is significantly different from zero (p<.05).

- Statistic not shown. Rate based on less than 16 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.

Surveillance, Epidemiology, and End Results (SEER) Program: SEER 9, 13, & 18 Geographic Areas National Cancer Institute, USA





Figure 1.1

Figure 1.2

Leading Causes of Death in US, 1975 vs 2015

Percent of All Causes of Death




Source: US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

Figure 1.3

Us Death Rates, 1975-2015

Heart Disease compared to Neoplasms, by age at death




Source: US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).


Figure 1.4

Trends in SEER Incidence and US Death Rates by Primary Cancer Site 2006-2015

Trends in SEER Incidence Rates



Trends in US Cancer Death Rates



Source: SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG) and US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

Underlying rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).


For sex-specific cancer sites, the population was limited to the population of the appropriate sex.

* The APC is significantly different from zero ($p < .05$).

^a Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

Figure 1.5

Trends in SEER Incidence Rates by Age Group and Primary Cancer Site 2006-2015



Source: SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

Underlying rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).


For sex-specific cancer sites, the population was limited to the population of the appropriate sex.

* The APC is significantly different from zero ($p < .05$).

^a Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

Figure 1.6


Trends in US Death Rates by Age Group and Primary Cancer Site 2006-2015



Source: US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.
 Underlying rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).
 For sex-specific cancer sites, the population was limited to the population of the appropriate sex.
 * The APC is significantly different from zero ($p < .05$).

Figure 1.7

Trends in SEER Incidence Rates by Sex and Primary Cancer Site 2006-2015



Source: SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG). Underlying rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).


For sex-specific cancer sites, the population was limited to the population of the appropriate sex.

* The APC is significantly different from zero ($p < .05$).

^a Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

Figure 1.8

Trends in US Death Rates by Sex and Primary Cancer Site 2006-2015




Source: US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Underlying rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).

For sex-specific cancer sites, the population was limited to the population of the appropriate sex.

* The APC is significantly different from zero ($p < .05$).

Figure 1.9

SEER Incidence^a and US Death Rates,^b 2011-2015 5-Year Relative Survival,^c 2008-2014 All Cancer Combined, by Race and Sex




^a Incidence rates are from the SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG) and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).

^b Death rates are from the US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).

^c Survival rates are from the SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

Figure 1.10


Trends in US Death Rates, 2006-2015 All Ages, by Race and Primary Cancer Site



Source: US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

Figure 1.11

5-Year Relative Survival (%) SEER Program, 2008-2014 Both Sexes, by Race and Cancer Site




Source: SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

^a Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

Figure 1.12

SEER Cancer Incidence and US Death Rates, 2011-2015 By Cancer Site and Race/Ethnicity



Source: SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG) and US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.


^a Rates for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives.

Incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

Figure 1.13

SEER Incidence 2006-2015 Males by Race/Ethnicity



Source: SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).

Regression lines are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.


^a Incidence rates for American Indian/Alaska Native (AI/AN) are based on the CHSDA(Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives.

Incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

Figure 1.14

SEER Incidence 2006-2015 Females by Race/Ethnicity



Source: SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG). Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).

Regression lines are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.


^a Incidence rates for American Indian/Alaska Native (AI/AN) are based on the CHSDA (Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives.

Incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

Figure 1.15

US Mortality 2006-2015 Males by Race/Ethnicity



Source: US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).


Regression lines are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.

^a Mortality rates for American Indian/Alaska Native (AI/AN) are based on the CHSDA (Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives.

Figure 1.16

US Mortality 2006-2015 Females by Race/Ethnicity



Source: US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).


Regression lines are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.

^a Mortality rates for American Indian/Alaska Native (AI/AN) are based on the CHSDA (Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives.

Figure 1.17

Incidence Percent Change between 2006 and 2015 Numbers (burden) vs Rates (risk) All Races, All Ages, Both Sexes



Source: SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG). Burden is the change in the number of incidence cases between 2006 and 2015. Risk is the change in the cancer incidence rates between 2006 and 2015.


^a Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

Figure 1.18

Mortality Percent Change between 2006 and 2015

Numbers (burden) vs Rates (risk)


All Races, All Ages, Both Sexes




US Mortality estimates based on US age-specific rates applied to US population.
 Burden is the change in the number of deaths between 2006 and 2015.
 Risk is the change in the cancer death rates between 2006 and 2015.

Figure 1.19

Person-Years of Life Lost Due to Cancer All Races, Both Sexes, 2015




Average Years of Life Lost Per Person Dying of Cancer All Races, Both Sexes, 2015




Source: US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention and 2014 Life Tables.

Figure 1.20

Person-Years of Life Lost Due to Major Causes of Death in US All Races, Both Sexes, 2015




Average Years of Life Lost Per Person Due to Major Causes of Death in US All Races, Both Sexes, 2015



Source: US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention and 2014 Life Tables.

Figure 1.21

SEER Observed Incidence and Delay Adjusted Incidence Rates^a All Cancer Sites, By Sex



Source: SEER 9 areas.


^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).

Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Figure 1.22

SEER Observed Incidence and Delay Adjusted Incidence Rates^a Both Sexes



Source: SEER 9 areas.


^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).

Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Figure 1.23

SEER Observed Incidence and Delay Adjusted Incidence Rates^a Males



Source: SEER 9 areas.


^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).

Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Figure 1.24

SEER Observed Incidence and Delay Adjusted Incidence Rates^a Females



Source: SEER 9 areas.

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).

Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).