

Table 1.23  
U.S. Complete Prevalence Counts, Invasive Cancers Only, January 1, 2015<sup>a</sup>  
By Age at Prevalence

Site/Sex	Age at Prevalence								
	All Ages <sup>c</sup>	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70+
All Sites									
Males	7,032,494	18,799	43,681	92,992	172,866	345,981	942,023	1,888,544	3,527,607
Females	8,079,604	17,434	38,006	96,195	248,166	638,524	1,448,944	2,081,139	3,511,196
Oral Cavity & Pharynx									
Males	240,247	60	336	1,691	3,835	13,583	51,730	81,471	87,541
Females	119,471	96	496	1,605	4,119	8,518	21,670	31,757	51,208
Esophagus									
Males	36,637	0	0	28	231	975	4,943	12,273	18,186
Females	10,647	0	0	11	55	198	1,426	3,121	5,836
Stomach									
Males	56,530	0	23	150	711	2,701	8,186	15,329	29,430
Females	41,385	0	61	172	792	2,373	6,357	9,414	22,216
Colon & Rectum									
Males	661,891	33	250	1,423	7,046	26,892	94,246	161,215	370,786
Females	670,194	0	371	1,716	7,187	25,306	82,985	135,011	417,617
Liver & Intrahep									
Males	50,546	578	624	681	635	1,774	10,577	22,897	12,780
Females	21,444	441	607	449	509	1,106	3,718	7,157	7,456
Pancreas									
Males	34,774	23	46	131	515	1,660	6,226	11,193	14,981
Females	33,841	11	46	330	783	1,891	6,174	9,420	15,186
Larynx									
Males	80,116	0	23	79	241	1,631	10,408	23,663	44,070
Females	19,640	0	0	17	152	770	3,601	6,083	9,016
Lung & Bronchus									
Males	238,904	45	110	427	1,280	5,040	25,457	66,067	140,479
Females	302,131	44	81	398	1,613	6,862	36,368	77,674	179,091
Melanoma of the Skin									
Males	613,293	68	636	5,049	18,308	45,387	105,775	168,413	269,658
Females	608,730	102	784	10,401	36,985	74,100	131,803	152,396	202,159

<sup>a</sup> U.S. 2015 cancer prevalence counts are based on 2015 cancer prevalence proportions from the SEER 9 registries (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta) and 1/1/2015 U.S. population estimates based on the average of 2014 and 2015 population estimates from the U.S. Bureau of the Census. Prevalence was calculated using the first invasive tumor for each cancer site diagnosed during the previous 40 years (1975-2014).

<sup>b</sup> Cases diagnosed more than 40 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000).

<sup>c</sup> Due to rounding, the sum of the age specific estimates may not equal the all ages estimate.

Table 1.23 - continued  
 U.S. Complete Prevalence Counts, Invasive Cancers Only, January 1, 2015<sup>a</sup>  
 By Age at Prevalence

Site/Sex	Age at Prevalence								
	All Ages <sup>c</sup>	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70+
Breast									
Males	19,095	0	0	11	153	624	2,094	5,126	11,088
Females	3,418,124	11	12	3,313	39,205	214,191	604,877	942,597	1,613,918
Cervix									
Females	257,524	0	81	2,063	14,486	40,266	60,508	63,055	77,065
Corpus & Uterus, NOS									
Females	727,200	11	23	844	6,718	27,300	101,936	215,845	374,522
Ovary <sup>d</sup>									
Females	224,940	113	948	3,985	7,585	18,327	46,034	62,323	85,625
Prostate									
Males	3,120,176	57	46	80	339	15,748	231,637	864,374	2,007,896
Urinary Bladder									
Males	529,318	62	105	595	2,505	9,096	43,303	120,318	353,336
Females	179,126	68	52	251	1,033	3,695	14,768	37,488	121,771
Kidney & Renal Pelvis									
Males	306,570	1,572	2,258	3,120	6,122	20,682	54,352	87,265	131,198
Females	198,810	1,676	2,482	3,197	5,496	13,754	33,887	50,099	88,220
Hodgkin Lymphoma									
Males	108,267	186	2,304	10,001	16,774	22,863	25,896	19,002	11,241
Females	100,538	45	1,912	9,703	16,856	22,201	23,482	15,606	10,735
Non-Hodgkin Lymphoma									
Males	364,491	828	4,278	8,341	14,678	30,274	62,974	91,757	151,361
Females	321,551	626	1,720	4,732	10,140	21,355	48,928	80,035	154,015
Myeloma									
Males	69,065	0	0	88	694	3,299	11,384	20,714	32,886
Females	55,668	0	0	67	464	2,560	8,464	16,640	27,472
Leukemia									
Males	230,623	6,548	13,735	14,815	13,747	17,054	29,724	50,174	84,826
Females	175,192	6,088	10,463	13,168	11,932	13,097	21,058	32,759	66,628
Acute Lymphocytic Leuk									
Males	45,907	5,461	11,360	11,466	7,880	5,297	2,407	1,257	780
Females	38,863	5,145	8,877	9,392	6,746	4,468	2,136	1,295	803

<sup>a</sup> U.S. 2015 cancer prevalence counts are based on 2015 cancer prevalence proportions from the SEER 9 registries (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta) and 1/1/2015 U.S. population estimates based on the average of 2014 and 2015 population estimates from the U.S. Bureau of the Census. Prevalence was calculated using the first invasive tumor for each cancer site diagnosed during the previous 40 years (1975-2014).

<sup>b</sup> Cases diagnosed more than 40 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000).

<sup>c</sup> Due to rounding, the sum of the age specific estimates may not equal the all ages estimate.