Table 1.24

Age-Adjusted SEER Incidence Rates and Trends for the Top 15 Cancer Sites by Race/Ethnicity

Both Sexes

White

Black

AII RACES			MILLE			Black		
	Rateb	APC°		Rateb	APC ^c		Rateb	APCc
	2011-2015			2011-2015	2006-2015		2011-2015	2006-2015
All Sites	439.2	-1.2*	All Sites	448.8	-1.2*	All Sites	453.4	-1.7*
Breast	67.5	0.0	Breast	68.1	-0.1	Prostate ^f	76.4	-5.2*
Lung and Bronchus	54.6	-2.4*	Lung and Bronchus	56.1	-2.4*	Breast	72.0	0.3
Prostate ^f	51.5	-5.7*	Prostate ^f	48.9	-6.2*	Lung and Bronchus	61.4	-2.6*
Colon and Rectum	39.4	-2.7*	Colon and Rectum	38.8	-2.6*	Colon and Rectum	47.6	-3.3*
Melanoma of the Skin	22.8	1.5*	Melanoma of the Skin	27.3	1.5*	Kidney and Renal Pelvis	18.3	0.7
Urinary Bladder	19.5	-1.3*	Urinary Bladder	21.5	-1.3*	Pancreas	15.5	-0.5
Non-Hodgkin Lymphoma	19.4	-0.5*	Non-Hodgkin Lymphoma	20.3	-0.7*	Non-Hodgkin Lymphoma	14.4	-0.2
Kidney and Renal Pelvis	15.9	0.5*	Kidney and Renal Pelvis	16.3	0.5*	Corpus and Uterus, NOSf	14.3	2.1*
Thyroid	14.5	3.3*	Thyroid	15.3	3.2*	Myeloma	13.4	0.8
Leukemia	13.8	0.3	Leukemia	14.6	0.2	Urinary Bladder	12.3	-0.8*
Corpus and Uterus, NOSf	13.8	1.2*	Corpus and Uterus, NOSf	14.0	1.0*	Leukemia	11.0	0.9
Pancreas	12.6	0.5*	Pancreas	12.6	0.6*	Liver & IBD ^g	10.5	2.5*
Oral Cavity and Pharynx	11.3	0.6*	Oral Cavity and Pharynx	11.9	0.9*	Stomach	10.1	-2.8*
Liver & IBDg	8.8	2.5*	Liver & IBD ^g	8.0	3.1*	Oral Cavity and Pharynx	9.0	-1.6*
Stomach	7.2	-0.9*	Brain and ONS ^g	7.1	-0.4	Thyroid	8.9	4.2*
Asian/Pacific Islander			American Indian/Alaska Native ^d			<u> Hispanic</u> e		
	Rateb	APC ^c		Rateb	APC ^c		Rate ^b	APCc
	2011-2015	2006-2015		2011-2015	2006-2015		2011-2015	2006-2015
All Sites	297.9	-1.2*	All Sites	315.1	-0.9*	All Sites	336.3	-1.1*
Breast	55.3	1.1*	Breast	44.4	0.5	Breast	50.0	0.5
Lung and Bronchus	35.6	-1.6*	Colon and Rectum	41.8	0.4	Prostate	40.9	-6.1*
Colon and Rectum	33.0	-2.9*	Lung and Bronchus	37.3	-2.7*	Colon and Rectum	33.5	-2.2*
Prostatef	25.9	-7.0*	Prostate ^f	24.3	-6.9*	Lung and Bronchus	27.8	-2.5*
Thyroid	14.0	2.6*	Kidney and Renal Pelvis		-3.1*	Non-Hodgkin Lymphoma	17.5	-0.2
Non-Hodgkin Lymphoma	13.6	0.0	Liver & IBD ^g	13.5	0.9	Kidney and Renal Pelvis		1.0*
Liver & IBD ^g	13.1	-2.0*	Non-Hodgkin Lymphoma	11.8	-1.9	Liver & IBD ^g	13.3	1.7*
Corpus and Uterus, NOSf	11.4	1.9*	Corpus and Uterus, NOS ^f		-0.2	Thyroid	12.5	4.0*
Stomach	10.3	-3.0*	Stomach	10.2	0.2	Corpus and Uterus, NOS ^f	11.9	3.1*
Pancreas	10.0	0.2	Pancreas	9.3	-1.9	Pancreas	11.2	0.2
Urinary Bladder	8.6	-1.5*	Thyroid	9.3	4.4*	Urinary Bladder	10.8	-1.6*
Kidney and Renal Pelvis				0 0	1.2	Leukemia	10.6	-0.2
-		0.7	Oral Cavity and Pharynx					
Leukemia	7.8	0.1	Urinary Bladder	8.8	0.2	Stomach	10.2	-1.6*
-	7.8						10.2	

Source: SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

- a Top 15 cancer sites selected based on 2011-2015 age-adjusted rates for the race/ethnic group.
- Pates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups Census P25-1130).
- The APC is the Annual Percent Change over the time interval.
- Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups Census P25-1130).
- d Rates for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.
- e Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives.
- Incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.
- f The rates for sex-specific cancer sites are calculated using the population for both sexes combined.
- g IBD = Intrahepatic Bile Duct. ONS = Other Nervous System.
- Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.
- * The APC is significantly different from zero (p<.05).

All Races

- Statistic not shown. Rate based on less than 16 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.