

Table 1.26
Age-Adjusted SEER Incidence Rates and Trends for the Top 15 Cancer Sites^a by Race/Ethnicity

Females

All Races	Rate ^b		APC ^c	All Sites	Rate ^b		APC ^c	All Sites	Rate ^b		APC ^c
	2011-2015	2006-2015			2011-2015	2006-2015			2011-2015	2006-2015	
All Sites	409.9	-0.2*		All Sites	423.1	-0.2*		All Sites	397.8	-0.4*	
Breast	126.0	0.2		Breast	128.6	0.0		Breast	126.9	0.4	
Lung and Bronchus	47.8	-1.8*		Lung and Bronchus	50.2	-1.7*		Lung and Bronchus	47.9	-1.8*	
Colon and Rectum	34.5	-2.6*		Colon and Rectum	34.0	-2.4*		Colon and Rectum	41.9	-3.4*	
Corpus and Uterus, NOS	26.0	1.4*		Corpus and Uterus, NOS	26.6	1.2*		Corpus and Uterus, NOS	25.4	2.3*	
Thyroid	21.4	3.3*		Thyroid	22.8	3.2*		Pancreas	14.3	-0.4	
Melanoma of the Skin	17.7	1.3*		Melanoma of the Skin	21.5	1.3*		Thyroid	13.4	4.6*	
Non-Hodgkin Lymphoma	15.9	-0.7*		Non-Hodgkin Lymphoma	16.8	-0.8*		Kidney and Renal Pelvis	12.8	0.6	
Ovary ^g	11.6	-1.8*		Ovary ^g	12.1	-1.9*		Non-Hodgkin Lymphoma	12.1	-0.2	
Pancreas	11.2	0.5*		Leukemia	11.4	0.1		Myeloma	11.6	0.9	
Kidney and Renal Pelvis	10.9	0.3		Kidney and Renal Pelvis	11.3	0.4		Ovary ^g	9.3	-1.4*	
Leukemia	10.8	0.2		Pancreas	11.1	0.6*		Leukemia	9.0	1.3*	
Urinary Bladder	8.3	-1.4*		Urinary Bladder	8.9	-1.4*		Cervix Uteri	8.4	-2.6*	
Cervix Uteri	7.4	-1.4*		Cervix Uteri	7.4	-1.2*		Stomach	7.7	-2.6*	
Oral Cavity and Pharynx	6.3	0.1		Oral Cavity and Pharynx	6.5	0.2		Urinary Bladder	6.7	-1.2*	
Brain and ONS ^f	5.4	-0.3		Brain and ONS ^f	6.0	-0.2		Oral Cavity and Pharynx	5.1	-1.3	
Asian/Pacific Islander				American Indian/Alaska Native^d				Hispanic^e			
	Rate ^b	APC ^c		Rate ^b	APC ^c		Rate ^b	APC ^c			
	2011-2015	2006-2015		2011-2015	2006-2015		2011-2015	2006-2015			
All Sites	298.3	-0.1		All Sites	311.4	-0.2		All Sites	322.9	0.3*	
Breast	100.6	1.2*		Breast	82.6	0.6		Breast	93.7	0.6*	
Lung and Bronchus	28.0	-0.8		Colon and Rectum	39.2	0.7		Colon and Rectum	28.6	-1.7*	
Colon and Rectum	28.0	-3.2*		Lung and Bronchus	31.2	-3.2		Lung and Bronchus	23.2	-1.5*	
Corpus and Uterus, NOS	20.8	1.9*		Corpus and Uterus, NOS	19.6	-0.1		Corpus and Uterus, NOS	22.5	3.3*	
Thyroid	20.2	2.2*		Thyroid	14.3	4.9*		Thyroid	19.6	4.3*	
Non-Hodgkin Lymphoma	11.1	-0.1		Kidney and Renal Pelvis	13.0	-2.5		Non-Hodgkin Lymphoma	15.3	0.2	
Ovary ^g	9.6	-0.3		Non-Hodgkin Lymphoma	10.8	-1.6		Kidney and Renal Pelvis	11.9	1.6*	
Pancreas	9.2	0.2		Ovary ^g	9.0	-5.0*		Pancreas	10.5	0.6	
Stomach	7.9	-3.1*		Liver & IBD ^f	8.9	2.0		Ovary ^g	10.4	-1.2	
Liver & IBD ^f	7.4	-2.2*		Cervix Uteri	8.1	1.7		Cervix Uteri	8.9	-3.6*	
Leukemia	6.4	-0.1		Pancreas	7.8	-5.8*		Leukemia	8.8	-0.3	
Cervix Uteri	6.1	-2.2*		Stomach	7.3	0.3		Stomach	8.3	-0.2	
Kidney and Renal Pelvis	5.6	0.0		Leukemia	6.5	-1.0		Liver & IBD ^f	7.7	2.6*	
Oral Cavity and Pharynx	5.1	0.2		Oral Cavity and Pharynx	6.1	-		Myeloma	5.0	1.0	
Urinary Bladder	3.9	-0.1		Myeloma	5.5	-		Urinary Bladder	4.8	-1.4*	

Source: SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

^a Top 15 cancer sites selected based on 2011-2015 age-adjusted rates for the race/ethnic group.

^b Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^c The APC is the Annual Percent Change over the time interval.

Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^d Rates for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^e Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives.

Incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

^f IBD = Intrahepatic Bile Duct. ONS = Other Nervous System.

^g Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

* The APC is significantly different from zero (p<.05).

- Statistic not shown. Rate based on less than 16 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.