Table 6.17 Cancer of the Colon and Rectum (Invasive)

> SEER^a Relative Survival (Percent) By Year of Diagnosis

> > All Races, Females

Year of Diagnosis

1975-1980-1985-1990-1979 1984 1989 1994 1995 1996 1997 1998 1999 2000 2001 2002 2003 2004 2005 2006 2007 2008 2009 2010 2011 2012 2013 2014

Survival Time

| 1-year | 73.9 | 76.3 | 78.9 | 80.0 | 78.0 | 79.9 | 79.3 | 82.0 | 81.2 | 80.2 | 81.0 | 83.4 | 81.7 | 82.5 | 83.4 | 82.5 | 82.5 | 83.1 | 83.0 | 82.9 | 83.3 | 83.4 | 83.6 | 83.3 |
|---------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|
| 2-year | 63.0 | 66.4 | 69.8 | 71.6 | 70.0 | 71.6 | 71.1 | 74.6 | 73.9 | 72.5 | 73.7 | 77.1 | 75.3 | 75.3 | 76.0 | 76.0 | 75.9 | 77.3 | 76.1 | 76.1 | 76.4 | 75.7 | 77.2 | |
| 3-year | 57.4 | 60.3 | 64.6 | 66.4 | 64.6 | 67.4 | 66.2 | 70.0 | 69.5 | 68.5 | 69.3 | 72.4 | 70.5 | 70.6 | 71.9 | 71.3 | 71.3 | 72.6 | 71.4 | 71.2 | 72.2 | 71.1 | | |
| 4-year | 53.8 | 56.6 | 61.2 | 63.0 | 60.8 | 64.7 | 63.1 | 67.2 | 66.2 | 64.9 | 66.5 | 69.3 | 67.6 | 67.5 | 68.9 | 68.1 | 68.6 | 69.8 | 68.4 | 68.6 | 68.4 | | | |
| 5-year | 51.4 | 54.1 | 58.9 | 60.9 | 59.4 | 62.7 | 60.6 | 64.1 | 64.1 | 63.5 | 64.7 | 66.9 | 65.3 | 65.3 | 66.7 | 65.9 | 66.6 | 67.2 | 66.2 | 66.2 | | | | |
| 6-year | 49.8 | 52.3 | 57.2 | 59.5 | 57.3 | 61.5 | 59.2 | 62.7 | 62.4 | 62.4 | 63.2 | 65.5 | 64.2 | 63.5 | 65.4 | 64.6 | 64.9 | 65.5 | 64.3 | | | | | |
| 7-year | 48.7 | 51.0 | 55.7 | 58.1 | 55.4 | 60.6 | 57.9 | 61.4 | 61.3 | 61.4 | 61.3 | 64.3 | 62.8 | 63.0 | 64.3 | 62.8 | 63.7 | 63.7 | | | | | | |
| 8-year | 47.9 | 50.1 | 54.6 | 57.1 | 54.7 | 59.4 | 57.6 | 61.0 | 60.1 | 60.4 | 60.1 | 63.7 | 62.5 | 62.1 | 63.2 | 61.7 | 62.4 | | | | | | | |
| 9-year | 47.3 | 49.3 | 53.9 | 56.3 | 54.5 | 58.5 | 57.0 | 59.9 | 59.5 | 59.4 | 59.2 | 62.2 | 61.4 | 61.1 | 62.7 | 61.0 | | | | | | | | |
| 10-year | 46.7 | 48.7 | 53.1 | 55.6 | 53.8 | 58.0 | 56.0 | 59.1 | 59.3 | 59.3 | 58.8 | 61.6 | 60.5 | 60.3 | 61.5 | | | | | | | | | |
| 11-year | 46.1 | 48.2 | 52.5 | 55.0 | 53.0 | 57.7 | 55.3 | 58.3 | 58.4 | 58.3 | 58.6 | 61.0 | 59.9 | 59.0 | | | | | | | | | | |
| 12-year | 45.9 | 47.7 | 52.3 | 54.4 | 52.4 | 57.6 | 54.7 | 58.3 | 58.0 | 57.8 | 57.3 | 60.2 | 59.6 | | | | | | | | | | | |
| 13-year | 45.5 | 47.4 | 51.4 | 54.1 | 51.6 | 56.9 | 54.4 | 57.9 | 58.0 | 57.2 | 56.9 | 59.7 | | | | | | | | | | | | |
| 14-year | 45.1 | 47.0 | 51.0 | 53.8 | 51.5 | 56.1 | 53.7 | 57.7 | 57.6 | 56.9 | 56.2 | | | | | | | | | | | | | |
| 15-year | 44.6 | 46.6 | 50.5 | 53.5 | 51.3 | 55.2 | 52.4 | 57.7 | 56.9 | 56.9 | | | | | | | | | | | | | | |
| 16-year | 44.2 | 45.9 | 50.1 | 53.4 | 50.6 | 54.6 | 52.0 | 57.3 | 56.1 | | | | | | | | | | | | | | | |
| 17-year | 44.0 | 45.6 | 49.5 | 52.7 | 50.0 | 53.4 | 52.0 | 57.0 | | | | | | | | | | | | | | | | |
| 18-year | 43.4 | 45.3 | 49.2 | 52.3 | 49.5 | 53.1 | 51.9 | | | | | | | | | | | | | | | | | |
| 19-year | 43.4 | 44.8 | 48.9 | 51.4 | 48.8 | 52.0 | | | | | | | | | | | | | | | | | | |
| 20-year | 43.1 | 44.6 | 48.5 | 50.5 | 48.4 | | | | | | | | | | | | | | | | | | | |
| 21-year | 42.7 | 44.0 | 48.5 | 50.0 | | | | | | | | | | | | | | | | | | | | |
| 22-year | 42.5 | 43.6 | 48.2 | | | | | | | | | | | | | | | | | | | | | |
| 23-year | 41.9 | 43.4 | 47.8 | | | | | | | | | | | | | | | | | | | | | |
| 24-year | 41.6 | 43.2 | 47.3 | | | | | | | | | | | | | | | | | | | | | |
| 25-year | 41.5 | 42.5 | 47.2 | | | | | | | | | | | | | | | | | | | | | |
| 26-year | 41.2 | 42.3 | 46.6 | | | | | | | | | | | | | | | | | | | | | |
| 27-year | 40.7 | 42.0 | | | | | | | | | | | | | | | | | | | | | | |
| 28-year | 40.7 | 41.4 | | | | | | | | | | | | | | | | | | | | | | |
| 29-year | 40.2 | 41.4 | | | | | | | | | | | | | | | | | | | | | | |
| 30-year | 40.1 | 41.1 | | | | | | | | | | | | | | | | | | | | | | |

Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). Expected survival rates are derived from the U.S. Annual Life Tables.