

Table 15.12
Cancer of the Lung and Bronchus (Invasive)

5-Year Relative and Period Survival (Percent) by Race, Sex, Diagnosis Year, Stage and Age

	All Races			Whites			Blacks		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
5-Year Relative Survival (Percent)									
Year of Diagnosis:									
1960-1963 ^a	-	-	-	8	7	11	5	5	6
1970-1973 ^a	-	-	-	10	9	14	7	6	10
1975-1977 ^b	12.2	11.1	15.3	12.3	11.1	15.4	11.2	10.5	13.8
1978-1980 ^b	12.8	11.3	16.3	12.9	11.5	16.2	11.7	9.7	17.8
1981-1983 ^b	13.1	11.5	16.4	13.3	11.7	16.6	11.4	10.2	14.9
1984-1986 ^b	12.8	11.2	16.0	13.0	11.2	16.3	11.1	10.3	12.8
1987-1989 ^b	13.0	11.9	15.0	13.3	12.0	15.3	10.9	10.8	11.1
1990-1992 ^b	13.6	12.1	15.9	14.0	12.5	16.2	10.5	9.3	12.7
1993-1995 ^b	14.3	12.5	16.9	14.5	12.6	17.1	12.8	11.2	15.7
1996-1998 ^b	14.6	12.9	16.8	14.8	13.1	17.0	12.3	10.7	15.0
1999-2001 ^b	15.3	13.1	17.9	15.5	13.3	18.0	12.7	10.8	15.4
2002-2004 ^b	16.1	14.2	18.4	16.5	14.4	18.9	13.2	12.8	13.9
2005-2007 ^b	17.8	15.2	20.6	18.3	15.7	21.1	14.0	12.2	16.1
2008-2014 ^b	19.9 ^f	16.7 ^f	23.3 ^f	20.0 ^f	16.9 ^f	23.2 ^f	18.3 ^f	15.0 ^f	21.9 ^f
5-Year Period Survival (Percent) ^{cd}									
2014	19.0	15.7	22.7	19.1	15.8	22.7	16.4	13.3	20.3
Stage Distribution (%) 2008-2014 ^{ce}									
All Stages									
Number of cases	264,275	138,788	125,487	214,482	111,164	103,318	30,645	16,916	13,729
Percent	100%	100%	100%	100%	100%	100%	100%	100%	100%
Localized	16	14	18	17	15	19	13	12	15
Regional	22	22	22	22	22	22	22	21	23
Distant	57	59	55	57	58	54	61	63	58
Unstaged/Unknown	5	4	5	5	5	5	4	4	4
5-Year Relative Survival (Percent), 2008-2014 ^c									
Age at Diagnosis:									
Ages <45	33.2	28.0	38.1	34.6	29.0	39.8	27.3	24.0	31.1
Ages 45-54	21.3	16.9	26.1	21.7	17.2	26.6	18.5	14.2	23.4
Ages 55-64	20.2	16.8	24.7	20.3	16.9	24.5	17.6	14.5	22.3
Ages 65-74	20.0	16.6	24.0	20.3	16.9	24.2	16.0	13.1	19.6
Ages 75+	14.1	11.8	16.2	14.3	12.0	16.3	11.3	8.1	14.0
Ages <65	21.2	17.3	25.9	21.4	17.5	26.0	18.3	14.9	23.1
Ages 65+	17.1	14.4	19.9	17.3	14.6	20.1	14.0	11.3	17.0
Stage ^e :									
All Stages	18.6	15.5	22.0	18.7	15.7	22.0	16.2	13.2	19.8
Localized	56.3	50.2	61.5	56.3	50.5	61.3	52.1	44.6	59.1
Regional	29.7	26.2	33.6	29.6	26.2	33.3	27.2	23.7	31.1
Distant	4.7	3.9	5.7	4.5	3.7	5.5	4.6	3.9	5.6
Unstaged/Unknown	7.8	7.2	8.4	7.0	6.4	7.6	10.2	8.3	12.7

^a Based on End Results data from a series of hospital registries and one population-based registry.

^b SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta).

Based on follow-up of patients into 2015. [Expected survival rates](#) are derived from the U.S. Annual Life Tables.

^c SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG). Expected survival rates are derived from life tables by socio-economic status, geography and race developed by the SEER program.

Based on follow-up of patients into 2015.

^d Period survival provides a 2014 estimate of survival by piecing together the most recent conditional survival estimates from several cohorts. It is computed here using three year calendar blocks (2010-2012: 0-1 year survival), (2009-2011: 1-2 year survival), (2008-2010: 2-3 year survival), (2007-2009: 3-4 year survival), (2006-2008: 4-5 years survival).

^e Stage at diagnosis is classified using SEER Summary Stage 2000. Stage distribution percentages may not sum to 100 due to rounding.

^f The difference between 1975-1977 and 2008-2014 is statistically significant (p<.05).

^g The standard error is between 5 and 10 percentage points.

^h The standard error is greater than 10 percentage points.

- Statistic could not be calculated due to fewer than 25 cases during the time period.