

Table 17.2
MesotheliomaAge-adjusted SEER Incidence^a Rates by Year, Race and Sex

	All Races			Whites			Blacks		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
Year of Diagnosis:									
1975-2015	0.99	1.84	0.39	1.08	2.01	0.43	0.51	0.92	0.25
1975-1979	0.69	1.21	0.30	0.71	1.28	0.31	0.39	0.49	-
1980	0.98	1.83	0.35	1.05	1.97	0.39	-	-	-
1981	0.75	1.42	0.27	0.80	1.50	0.29	-	-	-
1982	0.83	1.42	0.38	0.90	1.53	0.42	-	-	-
1983	1.05	1.86	0.49	1.09	1.95	0.51	-	-	-
1984	1.06	2.04	0.36	1.15	2.24	0.37	-	-	-
1985	0.92	1.67	0.37	0.97	1.73	0.39	-	-	-
1986	0.94	1.67	0.43	1.01	1.79	0.49	-	-	-
1987	0.97	1.73	0.43	1.03	1.82	0.47	-	-	-
1988	0.98	1.85	0.39	1.05	1.98	0.41	-	-	-
1989	1.13	2.21	0.37	1.20	2.36	0.39	-	-	-
1990	1.15	2.14	0.46	1.24	2.31	0.50	-	-	-
1991	1.12	2.11	0.42	1.18	2.21	0.46	-	-	-
1992	1.23	2.48	0.39	1.33	2.73	0.42	-	-	-
1993	1.06	2.04	0.38	1.14	2.24	0.37	-	-	-
1994	1.16	2.17	0.42	1.24	2.32	0.45	1.06	-	-
1995	1.16	2.17	0.46	1.23	2.30	0.48	-	-	-
1996	1.15	2.14	0.45	1.27	2.38	0.50	-	-	-
1997	1.01	1.92	0.37	1.06	2.06	0.35	0.92	-	-
1998	1.08	2.06	0.40	1.16	2.19	0.46	-	-	-
1999	1.12	2.17	0.38	1.20	2.30	0.40	-	-	-
2000	1.08	2.11	0.34	1.21	2.36	0.38	-	-	-
2001	1.06	1.98	0.41	1.18	2.19	0.45	-	-	-
2002	1.03	1.95	0.38	1.16	2.18	0.43	-	-	-
2003	0.96	1.74	0.40	1.06	1.91	0.44	-	-	-
2004	1.11	2.05	0.45	1.22	2.26	0.49	-	-	-
2005	1.05	1.95	0.43	1.19	2.18	0.50	-	-	-
2006	0.97	1.65	0.46	1.07	1.84	0.48	-	-	-
2007	0.98	1.77	0.43	1.11	2.00	0.48	-	-	-
2008	0.96	1.81	0.35	1.07	2.00	0.38	-	-	-
2009	0.99	1.83	0.39	1.16	2.12	0.46	-	-	-
2010	1.00	1.83	0.41	1.11	2.06	0.42	-	-	-
2011	0.96	1.72	0.40	1.10	1.91	0.47	-	-	-
2012	0.95	1.68	0.41	1.11	1.91	0.52	-	-	-
2013	0.83	1.50	0.35	0.94	1.65	0.42	-	-	-
2014	0.92	1.60	0.42	1.02	1.72	0.48	-	-	-
2015	0.88	1.65	0.32	1.03	1.94	0.35	-	-	-

Individual rates for diagnosis years 1975-1979 available in web browser version of [Cancer Statistics Review](#).

^a SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).
 Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
 - Statistic not shown. Rate based on less than 16 cases for the time interval.