

Table 17.3  
MesotheliomaSEER Incidence<sup>a</sup> Rates, Age-Adjusted and Age-Specific Rates, by Race and Sex

|                                       | All Races |       |         | Whites |       |         | Blacks |       |         |
|---------------------------------------|-----------|-------|---------|--------|-------|---------|--------|-------|---------|
|                                       | Total     | Males | Females | Total  | Males | Females | Total  | Males | Females |
| <u>SEER Incidence</u>                 |           |       |         |        |       |         |        |       |         |
| <u>Age at Diagnosis</u>               |           |       |         |        |       |         |        |       |         |
| Age-Adjusted Rates, 2011-2015         |           |       |         |        |       |         |        |       |         |
| All ages                              | 0.9       | 1.6   | 0.4     | 1.0    | 1.8   | 0.5     | 0.5    | 0.9   | 0.3     |
| Under 65                              | 0.2       | 0.3   | 0.2     | 0.2    | 0.3   | 0.2     | 0.1    | 0.2   | 0.1     |
| 65 and over                           | 5.8       | 11.0  | 2.1     | 6.6    | 12.4  | 2.4     | 2.8    | 5.5   | 1.3     |
| All ages (WHO world std) <sup>b</sup> | 0.6       | 1.0   | 0.3     | 0.7    | 1.1   | 0.3     | 0.3    | 0.6   | 0.2     |
| Age-Specific Rates, 2011-2015         |           |       |         |        |       |         |        |       |         |
| <1                                    | -         | -     | -       | -      | -     | -       | -      | -     | -       |
| 1-4                                   | -         | -     | -       | -      | -     | -       | -      | -     | -       |
| 5-9                                   | -         | -     | -       | -      | -     | -       | -      | -     | -       |
| 10-14                                 | -         | -     | -       | -      | -     | -       | -      | -     | -       |
| 15-19                                 | -         | -     | -       | -      | -     | -       | -      | -     | -       |
| 20-24                                 | -         | -     | -       | -      | -     | -       | -      | -     | -       |
| 25-29                                 | -         | -     | -       | -      | -     | -       | -      | -     | -       |
| 30-34                                 | 0.1       | -     | 0.1     | 0.1    | -     | -       | -      | -     | -       |
| 35-39                                 | 0.1       | 0.1   | 0.1     | 0.1    | -     | -       | -      | -     | -       |
| 40-44                                 | 0.1       | 0.1   | -       | 0.1    | 0.2   | -       | -      | -     | -       |
| 45-49                                 | 0.3       | 0.4   | 0.3     | 0.4    | 0.4   | 0.3     | -      | -     | -       |
| 50-54                                 | 0.5       | 0.6   | 0.4     | 0.5    | 0.7   | 0.4     | -      | -     | -       |
| 55-59                                 | 0.9       | 1.1   | 0.6     | 0.9    | 1.2   | 0.7     | 0.6    | -     | -       |
| 60-64                                 | 1.6       | 2.3   | 0.9     | 1.8    | 2.6   | 1.0     | 1.0    | 1.4   | -       |
| 65-69                                 | 2.9       | 4.8   | 1.1     | 3.2    | 5.3   | 1.3     | 1.9    | 3.2   | -       |
| 70-74                                 | 5.0       | 8.6   | 2.0     | 5.7    | 9.7   | 2.2     | 2.6    | 4.4   | -       |
| 75-79                                 | 7.2       | 12.9  | 2.7     | 8.3    | 14.8  | 3.0     | 3.0    | -     | -       |
| 80-84                                 | 8.6       | 16.8  | 2.9     | 9.7    | 18.8  | 3.3     | 3.1    | -     | -       |
| 85+                                   | 8.7       | 19.7  | 2.9     | 9.6    | 21.6  | 3.3     | 4.7    | 14.0  | -       |

<sup>a</sup> SEER 18 areas. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.

<sup>b</sup> Rates are per 100,000 and are age-adjusted to the world (WHO 2000-2025) standard million.

- Statistic not shown. Rate based on less than 16 cases for the time interval.