

Table 17.7  
MesotheliomaSEER<sup>a</sup> Relative Survival (Percent)  
By Year of Diagnosis

All Races, Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
Survival Time																								
1-year	40.8	43.7	42.3	48.0	38.8	32.3	46.3	54.5	43.7	37.1	52.3	35.7	51.2	46.0	41.0	57.0	46.7	57.8	42.9	42.5	52.3	41.9	42.7	38.7
2-year	31.1	28.3	24.5	31.2	23.1	26.8	27.9	34.8	39.3	19.0	34.8	19.4	32.9	22.3	27.9	45.8	27.9	26.1	27.2	18.4	38.4	31.8	30.4	
3-year	24.4	19.4	18.9	24.4	21.2	21.3	17.2	26.0	37.4	16.7	22.8	9.8	24.8	14.3	25.9	37.8	14.3	17.8	23.0	16.5	28.1	25.4		
4-year	21.5	14.4	15.4	21.3	19.3	11.8	14.4	21.6	35.4	8.6	18.8	9.8	16.0	14.3	23.7	29.3	10.3	15.2	23.0	14.2	28.1			
5-year	20.1	13.3	13.4	20.2	17.6	9.8	14.4	21.6	26.2	0.0	18.8	7.8	14.2	12.4	19.3	26.0	6.3	15.2	23.0	12.0				
6-year	17.1	10.5	11.9	16.9	15.2	7.9	14.4	21.6	23.8	0.0	16.9	7.8	12.2	12.4	19.3	24.9	4.3	12.2	22.2					
7-year	16.3	10.5	10.8	16.5	15.2	6.0	8.9	20.5	21.4	0.0	15.0	7.8	10.2	10.4	17.6	21.1	4.3	4.3						
8-year	15.7	9.9	8.7	15.8	15.2	6.0	8.9	20.5	21.4	0.0	15.0	7.8	5.2	8.4	15.5	20.5	4.3							
9-year	15.2	9.5	8.7	15.3	12.9	4.0	8.9	20.5	19.0	0.0	15.0	7.8	5.2	8.4	15.5	11.7								
10-year	14.4	7.7	8.4	14.0	12.9	4.0	8.9	20.1	19.0	0.0	11.1	6.8	2.9	8.4	15.5									
11-year	13.7	7.7	8.4	13.7	12.9	4.0	8.9	18.1	19.0	0.0	11.1	6.8	2.9	8.4										
12-year	13.7	7.7	8.1	12.7	10.5	4.0	8.9	12.6	19.0	0.0	11.1	6.8	2.9											
13-year	13.7	7.7	8.1	11.8	10.5	4.0	8.9	12.6	19.0	0.0	11.1	5.1												
14-year	13.7	7.7	8.1	11.8	10.5	4.0	8.9	12.6	19.0	0.0	8.9													
15-year	13.7	6.8	8.1	11.6	10.5	4.0	8.9	12.6	19.0	0.0														
16-year	12.9	6.8	8.1	11.6	10.5	4.0	8.9	12.6	16.9															
17-year	12.3	6.8	8.1	11.6	10.5	4.0	8.9	9.7																
18-year	12.3	6.8	7.9	10.5	10.5	2.2	8.9																	
19-year	12.3	6.8	7.9	8.7	10.5	2.2																		
20-year	12.3	6.5	7.4	8.7	10.5																			

<sup>a</sup> Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). [Expected survival rates](#) are derived from the U.S. Annual Life Tables.