

Table 17.9
Mesothelioma (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years,
Lifetime Risk of Being Diagnosed with Cancer Given Alive and Cancer-Free at Current Age, and
Lifetime Risk of Dying from Cancer Given Alive at Current Age
Males, 2013-2015 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer
		+10 yrs	+20 yrs	+30 yrs	Ever				+10 yrs	+20 yrs	+30 yrs	Ever	
All Races	0	0.00	0.00	0.01	0.07	0.15	Asian/ Pacific Islander	0	0.00	0.00	0.00	0.05	0.07
	10	0.00	0.01	0.02	0.07	0.15		10	0.00	0.00	0.01	0.05	0.07
	20	0.01	0.02	0.04	0.07	0.15		20	0.00	0.01	0.02	0.05	0.07
	30	0.01	0.03	0.04	0.07	0.15		30	0.01	0.01	0.02	0.05	0.07
	40	0.02	0.03	0.03	0.05	0.15		40	0.01	0.01	0.02	0.04	0.07
	50	0.01	0.02	0.03	0.04	0.16		50	0.00	0.01	0.02	0.03	0.07
	60	0.01	0.02	0.03	0.03	0.17		60	0.00	0.01	0.02	0.03	0.07
	70	0.01	0.02	-	0.03	0.17		70	0.01	0.02	-	0.03	0.07
White	80	0.01	-	-	0.02	0.14	American Indian/ Alaska Native ^a	80	0.01	-	-	0.02	0.05
	0	0.00	0.00	0.00	0.06	0.16		0	0.00	0.00	0.00	0.04	0.09
	10	0.00	0.00	0.01	0.06	0.16		10	0.00	0.00	0.00	0.04	0.09
	20	0.00	0.01	0.03	0.06	0.16		20	0.00	0.00	0.01	0.04	0.09
	30	0.01	0.02	0.03	0.06	0.17		30	0.00	0.01	0.01	0.05	0.09
	40	0.01	0.02	0.03	0.05	0.17		40	0.01	0.01	0.02	0.05	0.09
	50	0.01	0.02	0.03	0.04	0.17		50	0.01	0.02	0.04	0.04	0.10
	60	0.01	0.02	0.03	0.03	0.18		60	0.01	0.04	0.04	0.04	0.11
Black	70	0.01	0.02	-	0.03	0.18	Hispanic ^b	70	0.03	0.03	-	0.03	0.11
	80	0.01	-	-	0.02	0.14		80	0.00	-	-	0.00	0.12
	0	0.00	0.00	0.04	0.13	0.06		0	0.00	0.00	0.01	0.11	0.10
	10	0.00	0.04	0.08	0.13	0.06		10	0.00	0.01	0.03	0.11	0.10
	20	0.04	0.08	0.10	0.13	0.06		20	0.01	0.02	0.04	0.11	0.10
	30	0.05	0.07	0.08	0.10	0.06		30	0.02	0.04	0.05	0.10	0.10
	40	0.03	0.04	0.05	0.06	0.06		40	0.02	0.03	0.04	0.09	0.11
	50	0.01	0.03	0.03	0.03	0.07		50	0.01	0.03	0.05	0.07	0.11
	60	0.01	0.02	0.02	0.02	0.07		60	0.01	0.03	0.05	0.06	0.11
	70	0.01	0.01	-	0.01	0.07		70	0.03	0.04	-	0.05	0.10
	80	0.00	-	-	0.00	0.06		80	0.02	-	-	0.04	0.08

Devcan 6.7.6, April 2018, National Cancer Institute (<https://surveillance.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, and Georgia excluding ATL/RG). Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives.

- Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

- Statistic could not be calculated.

A percent of 0.00 represents a value that is below 0.005.