

Table 17.12
MesotheliomaAge-Adjusted SEER Incidence Rates^a
By Registry, Race and Sex

	All Races			Whites			Blacks		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
<u>SEER Incidence Rates^a, 2011-2015</u>									
California	0.90	1.55	0.41	1.04	1.78	0.48	0.60	1.12	-
Greater Bay Area	0.90	1.52	0.44	1.12	1.91	0.54	-	-	-
San Francisco-Oakland	0.92	1.55	0.47	1.18	1.96	0.62	-	-	-
San Jose-Monterey	0.85	1.46	0.39	1.02	1.83	0.39	-	-	-
Los Angeles	0.76	1.27	0.38	0.90	1.50	0.45	0.57	1.04	-
Greater California	0.96	1.68	0.41	1.07	1.84	0.47	0.64	1.32	-
Connecticut	1.03	1.86	0.44	1.11	2.00	0.46	-	-	-
Detroit	0.91	1.70	0.37	1.07	1.98	0.42	0.36	-	-
Georgia	0.60	1.12	0.24	0.72	1.31	0.27	0.25	0.43	-
Atlanta	0.57	1.16	0.18	0.74	1.42	-	-	-	-
Rural Georgia	-	-	-	-	-	-	-	-	-
Greater Georgia	0.62	1.09	0.27	0.71	1.27	0.28	0.29	-	-
Hawaii	0.48	0.91	-	-	-	-	-	-	-
Iowa	0.84	1.40	0.43	0.86	1.44	0.42	-	-	-
Kentucky	0.78	1.42	0.31	0.80	1.47	0.31	-	-	-
Louisiana	1.31	2.26	0.64	1.47	2.52	0.69	0.86	1.49	0.50
New Jersey	1.15	2.05	0.53	1.30	2.29	0.59	0.48	0.88	-
New Mexico	0.79	1.38	0.32	0.82	1.38	0.36	-	-	-
Seattle-Puget Sound	1.30	2.35	0.50	1.39	2.47	0.56	-	-	-
Utah	0.83	1.44	0.33	0.83	1.43	0.33	-	-	-
SEER 9 Areas ^b	0.91	1.63	0.38	1.04	1.83	0.45	0.35	0.63	-
SEER 13 Areas ^b	0.87	1.54	0.38	1.01	1.76	0.44	0.40	0.72	0.23
SEER 18 Areas ^b	0.92	1.63	0.41	1.04	1.83	0.46	0.48	0.86	0.26

^a Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130)^b The SEER 9 areas are San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah and Atlanta.

The SEER 13 areas comprise the SEER 9 areas plus San Jose-Monterey, Los Angeles, the Alaska Native Registry and Rural Georgia.

The SEER 18 areas comprise the SEER 13 areas plus California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG.

- Statistic not shown. Rate based on less than 16 cases for the time interval.