

Table 18.9
MyelomaSEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Males and Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
Survival Time																								
1-year	68.0	69.5	72.3	71.9	74.0	70.6	74.3	75.0	73.6	72.5	72.7	73.6	77.2	74.8	76.2	74.2	77.2	77.8	82.0	81.4	79.8	81.4	81.3	82.8
2-year	52.1	54.5	57.6	57.4	58.7	57.6	60.6	62.5	59.8	60.0	59.8	62.9	65.0	66.5	64.9	66.4	66.8	69.0	72.8	72.1	71.7	73.3	72.0	
3-year	39.6	42.8	45.3	46.2	47.8	45.0	48.7	51.3	48.0	50.0	49.6	53.5	55.8	58.7	57.5	58.3	58.1	61.8	65.4	65.5	64.9	66.2		
4-year	31.5	34.2	35.7	36.9	40.1	38.2	38.7	40.0	38.9	41.3	44.1	46.9	47.7	50.9	50.8	51.3	50.6	55.9	59.7	58.8	57.7			
5-year	25.1	26.9	27.5	29.9	33.3	31.9	31.4	34.3	31.4	34.6	37.6	40.0	42.0	45.5	46.6	46.3	46.4	50.2	52.9	53.0				
6-year	20.1	21.0	22.1	24.5	28.3	27.2	26.4	29.9	27.2	29.3	32.8	35.3	37.6	40.1	41.6	41.2	41.3	44.5	48.6					
7-year	16.2	16.7	17.5	20.4	23.7	23.8	22.6	24.6	23.5	25.5	29.3	31.8	33.1	35.8	36.9	36.5	37.1	41.5						
8-year	13.5	13.4	14.9	17.3	20.4	21.4	19.9	22.4	20.2	22.9	25.8	28.7	29.8	32.1	33.7	32.8	33.9							
9-year	11.5	11.2	11.9	15.4	18.2	19.5	17.2	19.7	18.4	20.7	23.2	25.7	26.9	30.2	31.1	30.8								
10-year	9.8	9.6	10.4	13.7	17.1	17.3	15.6	18.1	16.5	19.6	21.3	23.1	23.9	27.2	28.4									
11-year	8.7	8.6	9.2	12.7	15.6	15.7	14.4	17.0	14.2	18.6	20.0	20.1	21.6	25.0										
12-year	7.8	7.6	8.4	11.2	14.3	14.9	13.1	15.2	13.8	17.1	18.9	19.1	20.3											
13-year	6.7	6.6	7.6	10.3	13.6	13.4	11.9	14.3	12.8	16.1	16.9	18.2												
14-year	6.1	6.0	7.0	9.6	12.4	12.7	11.3	14.0	11.5	15.1	16.1													
15-year	5.3	5.7	6.2	9.2	12.0	11.6	11.0	12.3	11.2	14.4														
16-year	4.9	5.1	5.7	8.7	10.8	9.9	10.6	11.4	10.3															
17-year	4.6	4.7	5.4	8.1	10.1	9.0	9.9	10.7																
18-year	4.0	4.3	5.1	7.6	9.9	8.4	8.7																	
19-year	3.8	3.8	4.9	6.9	9.0	7.8																		
20-year	3.3	3.5	4.7	6.8	7.7																			

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). [Expected survival rates](#) are derived from the U.S. Annual Life Tables.