

Table 18.15  
Myeloma

SEER Incidence and U.S. Mortality  
Age-Adjusted Rates and Trends<sup>a</sup>  
By Race/Ethnicity and Sex

SEER Incidence	SEER 18 Areas <sup>b</sup>			SEER 13 Areas <sup>b</sup>			SEER 9 Areas <sup>b</sup>		
	Rate 2011-2015			Trend 2006-2015 <sup>c</sup>			Trend 2006-2015 <sup>d</sup>		
	Rate per 100,000 persons			AAPC (%)			AAPC (%)		
RACE/ETHNICITY	Total	Males	Females	Total	Males	Females	Total	Males	Females
All Races	6.7	8.4	5.3	0.9	1.1	0.5*	0.9*	0.9*	0.7*
White	6.1	7.9	4.7	0.7*	0.8*	0.4*	0.8*	0.9*	0.6*
White Hispanic <sup>e</sup>	6.3	7.8	5.1	0.3	0.4	0.1	-	-	-
White Non-Hispanic <sup>e</sup>	6.0	7.8	4.6	0.8*	0.9*	0.4*	-	-	-
Black	13.4	15.9	11.6	0.7*	1.0*	0.4	0.1	-0.6	0.9*
Asian/Pacific Islander	3.9	4.9	3.1	0.7*	0.8*	0.7	-	-	-
Amer Ind/Alaska Nat <sup>f</sup>	5.8	6.2	5.5	0.8	1.3	0.3	-	-	-
Hispanic <sup>e</sup>	6.2	7.6	5.0	0.4*	0.4	0.2	-	-	-
U.S. Mortality <sup>g</sup>	Rate 2011-2015			Trend 2006-2015 <sup>h</sup>			AAPC (%)		
RACE/ETHNICITY	Total	Males	Females	Total	Males	Females	Total	Males	Females
All Races	3.3	4.2	2.7	-0.7	-0.6*	-0.7*			
White	3.1	4.0	2.4	-0.5*	-0.9*	-0.8*			
White Hispanic <sup>e</sup>	2.9	3.6	2.4	-0.6*	-0.5*	-1.3*			
White Non-Hispanic <sup>e</sup>	3.1	4.0	2.4	-0.5*	-0.9*	-0.8*			
Black	6.2	7.5	5.5	-0.5	-1.3*	-0.1			
Asian/Pacific Islander	1.6	2.0	1.3	-0.7*	-0.4	-0.9			
Amer Ind/Alaska Nat									
Total U.S.	2.1	2.4	1.9	-2.0*	-2.1*	-2.1*			
CHSDA Counties	3.0	3.4	2.7	-1.5*	-1.3*	-1.7			
Non-CHSDA Counties	1.2	1.3	1.1	-2.1*	-3.0*	-			
Hispanic <sup>e</sup>	2.8	3.4	2.3	-0.7*	-0.6*	-0.9*			

The AAPC is the Average Annual Percent Change over the time interval. The AAPCs are calculated by the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.

- Statistic not shown. Rate based on less than 16 cases for the time interval.

a Trend based on less than 10 cases for at least one year within the time interval.

b Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130). Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

c The SEER 9 areas are San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta.

d The SEER 13 areas comprise the SEER 9 areas plus San Jose-Monterey, Los Angeles, the Alaska Native Registry, and Rural Georgia.

e The SEER 18 areas comprise the SEER 13 areas plus California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, and Georgia excluding ATL/RG.

f The 2006-2015 AAPC estimates are based on a Joinpoint analysis with up to 4 Joinpoints over diagnosis years 1992-2015.

g The 2006-2015 AAPC estimates are based on a Joinpoint analysis with up to 5 Joinpoints over diagnosis years 1975-2015.

h Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry. The 2006-2015 Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.

i Incidence data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

j US Mortality Files, National Center for Health Statistics, CDC.

k The 2006-2015 mortality AAPCs are based on a Joinpoint analysis using years of death 1992-2015.

\* The APC is significantly different from zero (p<.05).