

Table 22.21
Cancer of the Pancreas (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2015
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <23 ^e	0 to <40 ^e	>=40 ^g	Complete ^h
Race	Sex										
All Races ^b	Both Sexes	51,209	9,768	3,649	1,734	1,027	501	67,393	68,335	280	68,615
	Males	26,610	4,996	1,653	739	331	175	34,332	34,642	132	34,774
	Females	24,599	4,772	1,996	995	696	326	33,061	33,693	148	33,841
White ^b	Both Sexes	42,537	8,172	3,081	1,262	822	329	55,876	56,558	234	56,792
	Males	22,389	4,226	1,507	600	277	120	29,000	29,227	120	29,347
	Females	20,148	3,946	1,574	662	545	209	26,876	27,331	114	27,445
Black ^b	Both Sexes	6,153	1,028	357	261	106	122	7,907	8,086	117	8,203
	Males	2,950	505	83	69	0	50	3,608	3,669	52	3,721
	Females	3,203	523	274	192	106	72	4,299	4,417	65	4,482
Asian/ Pacific Islander ^c	Both Sexes	1,988	379	154	+	+	+	2,645	+	+	+
	Males	960	187	59	+	+	+	1,242	+	+	+
	Females	1,028	192	95	+	+	+	1,403	+	+	+
Hispanic ^d	Both Sexes	3,542	731	253	+	+	+	4,767	+	+	+
	Males	1,614	321	106	+	+	+	2,103	+	+	+
	Females	1,928	410	147	+	+	+	2,664	+	+	+

Estimated prevalence percent^a on January 1, 2015, of the SEER^c population diagnosed in the previous 23 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	Age-Adjusted ^f All Ages
Race	Sex	Age Specific (Crude)										
All Races ^c	Both Sexes	0.0189%	-	0.0002%	0.0010%	0.0027%	0.0082%	0.0259%	0.0550%	0.0922%	0.0888%	0.0171%
	Males	0.0192%	-	0.0003%	0.0004%	0.0022%	0.0078%	0.0265%	0.0618%	0.1095%	0.1007%	0.0189%
	Females	0.0185%	-	0.0002%	0.0016%	0.0032%	0.0086%	0.0253%	0.0488%	0.0780%	0.0816%	0.0157%
White ^c	Both Sexes	0.0200%	-	0.0002%	0.0010%	0.0026%	0.0081%	0.0254%	0.0560%	0.0956%	0.0885%	0.0173%
	Males	0.0206%	-	0.0002%	0.0004%	0.0021%	0.0076%	0.0266%	0.0636%	0.1137%	0.1003%	0.0192%
	Females	0.0193%	-	-	0.0015%	0.0032%	0.0086%	0.0242%	0.0488%	0.0804%	0.0812%	0.0157%
Black ^c	Both Sexes	0.0170%	-	-	0.0011%	0.0028%	0.0110%	0.0341%	0.0628%	0.0955%	0.0772%	0.0188%
	Males	0.0159%	-	-	-	0.0026%	0.0101%	0.0314%	0.0741%	0.1074%	0.0615%	0.0193%
	Females	0.0179%	-	-	0.0016%	0.0030%	0.0117%	0.0365%	0.0539%	0.0872%	0.0850%	0.0184%
Asian/ Pacific Islander ^c	Both Sexes	0.0155%	-	-	0.0010%	0.0030%	0.0071%	0.0211%	0.0453%	0.0710%	0.0934%	0.0146%
	Males	0.0151%	-	-	-	0.0028%	0.0076%	0.0206%	0.0433%	0.0864%	0.1141%	0.0160%
	Females	0.0158%	-	-	0.0018%	0.0032%	0.0066%	0.0215%	0.0469%	0.0589%	0.0807%	0.0137%
Hispanic ^d	Both Sexes	0.0089%	-	0.0003%	0.0011%	0.0022%	0.0059%	0.0196%	0.0413%	0.0721%	0.0746%	0.0134%
	Males	0.0077%	-	-	-	0.0013%	0.0045%	0.0184%	0.0414%	0.0873%	0.0631%	0.0134%
	Females	0.0100%	-	-	0.0020%	0.0032%	0.0075%	0.0208%	0.0412%	0.0607%	0.0815%	0.0136%

^a US 2015 cancer prevalence counts are based on 2015 cancer prevalence proportions from the SEER registries and 1/1/2015 US population estimates based on the average of 2014 and 2015 population estimates from the US Bureau of the Census.
^{b c d} Prevalence was calculated using the first invasive tumor for each cancer site diagnosed during the previous 40/23 years. Statistics based on (b) SEER 9 Areas (c) SEER 13 Areas excluding the Alaska Native Registry
^e (d) NHIA for Hispanic for SEER 13 Areas excluding the Alaska Native Registry.
^f Maximum limited-duration prevalence: 40 years for 1975-2015 SEER 9 data; 23 years for 1992-2015 SEER 13 data (excluding the Alaska Native Registry) used to calculate prevalence for Hispanics and Asian Pacific Islanders.
^{g h i} Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups. (g) Cases diagnosed more than 40 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <40 and >=40. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.
 - Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.
 + Not available.