

Table 23.7
Cancer of the Prostate (Invasive)

SEER Incidence^a and U.S. Death^b Rates, Age-Adjusted and Age-Specific Rates, by Race

	SEER Incidence ^a			U.S. Mortality ^b		
	All Races Males	White Males	Black Males	All Races Males	White Males	Black Males
<u>Age-Adjusted Rates, 2011-2015</u>						
Age at Diagnosis/Death:						
All ages	112.6	105.7	178.3	19.5	18.2	39.9
Under 65	44.9	40.9	83.4	1.7	1.4	4.0
65 and over	580.7	553.1	834.3	142.9	134.0	288.4
All ages (WHO world std) ^c	85.7	80.2	139.0	10.9	10.0	23.0
<u>Age-Specific Rates, 2011-2015</u>						
Age at Diagnosis/Death:						
<1	-	-	-	-	-	-
1-4	-	-	-	-	-	-
5-9	-	-	-	-	-	-
10-14	-	-	-	-	-	-
15-19	-	-	-	-	-	-
20-24	-	-	-	-	-	-
25-29	-	-	-	-	-	-
30-34	0.1	-	-	-	-	-
35-39	0.7	0.6	1.6	0.0	-	-
40-44	8.1	6.9	19.9	0.2	0.1	0.4
45-49	36.0	29.9	89.4	0.8	0.7	1.8
50-54	116.9	104.8	228.7	3.1	2.7	7.3
55-59	251.0	228.8	460.8	8.5	7.1	20.5
60-64	423.2	394.7	721.5	19.7	17.0	47.5
65-69	651.7	616.1	1,017.6	38.4	34.0	92.0
70-74	673.1	641.7	959.3	70.7	63.6	166.4
75-79	589.3	560.5	796.1	126.3	116.2	273.3
80-84	444.2	425.2	582.5	219.9	207.3	447.2
85+	376.9	366.5	529.0	461.8	446.0	816.2

^a SEER 18 areas. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.

^b US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

^c Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.

- Rates are per 100,000 and are age-adjusted to the world (WHO 2000-2025) standard million.

- Statistic not shown. Rate based on less than 16 cases for the time interval.