

Table 29.6

5-Year Relative Survival (Percent), 2008-2014^aBy International Classification of Childhood Cancer (ICCC)^b Selected Group and Subgroup^c and Sex and Age
Excluding benign brain and myelodysplastic syndromes

ICCC Group & Subgroup	Ages 0-19 by Sex			Both Sexes by Age					
	Total	Male	Female	<1	1-4	5-9	10-14	15-19	0-14
All ICCC Groups Combined	83.8	82.7	85.0	78.4	85.5	82.8	82.9	84.6	83.4
I Leukemia	83.9	83.7	84.2	60.4	91.1	90.1	78.9	72.1	86.5
I(a) Lymphoid leukemia	88.5	88.1	89.0	57.5	94.5	92.8	83.4	73.8	90.8
I(b) Acute myeloid leukemia	65.8	65.3	66.3	63.0	70.9	70.4	60.6	64.2	66.4
II Lymphomas and reticuloendothelial neo.	93.3	93.1	93.6	84.1 ^d	88.7	94.7	93.4	93.5	93.0
II(a) Hodgkin lymphoma	96.7	96.5	96.8	-	94.4 ^d	99.4	97.4	96.1	97.8
II(b,c,e) Non-Hodgkin lymphoma	89.8	90.2	88.8	-	87.4	92.8	89.7	89.1	90.2
III CNS & misc intracranial & intraspinal neo.	73.9	73.2	74.6	61.3	74.5	68.9	79.1	77.9	72.9
III(a) Ependymomas and choroid plexus tumor	80.2	80.4	80.1	74.3 ^d	76.9	79.5 ^d	86.9 ^d	89.3	78.5
III(b) Astrocytoma	81.1	80.3	82.0	85.7	90.2	77.3	80.4	75.2	82.6
III(c) Intracranial and intraspinal embryonal tumors	63.5	62.1	65.6	28.4 ^d	57.7	70.5	77.3	71.2	62.6
III(d) Other gliomas	60.7	64.2	57.0	-	53.0	42.9	73.4	83.0	54.8
IV Neuroblastoma and other peripheral nervous cell tumors	79.3	77.7	81.0	91.1	75.3	73.6	83.1 ^d	54.1 ^d	80.2
IV(a) Neuroblastoma and ganglioneuroblastoma	79.3	77.7	81.0	91.1	75.4	72.8	83.6 ^d	-	80.2
V Retinoblastoma	95.2	94.4	95.9	96.7	94.1	-	-	-	95.2
VI Renal tumors	90.5	87.9	92.9	91.6	93.4	89.3	85.1 ^d	73.8 ^d	91.9
VI(a) Nephroblastoma and other nonepithelial renal tumor	92.5	91.0	93.8	91.5	93.6	91.0	88.8 ^d	-	92.7
VII Hepatic tumors	77.3	76.3	78.7	90.1	82.0	63.0 ^d	61.0 ^d	52.4 ^d	80.4
VII(a) Hepatoblastoma	84.3	83.1	86.4	90.0	81.9	-	-	-	84.6
VIII Malignant bone tumors	72.0	71.4	72.9	-	79.7 ^d	74.2	73.0	69.3	73.8
VIII(a) Osteosarcoma	68.0	65.8	70.9	-	-	68.2	70.2	65.7	69.6
VIII(c) Ewing tumor and related sarcomas of bone	73.1	74.3	71.2	-	97.1	80.3 ^d	72.7	64.3	77.7
IX Soft tissue and other extraosseous sarcomas	72.3	69.7	75.4	65.9	78.0	77.8	73.4	67.1	75.0
IX(a) Rhabdomyosarcoma	65.5	65.6	65.4	68.0 ^d	73.5	77.5	58.2	46.2	70.3
X Germ cell & trophoblastic tumors & neo. of gonads	92.3	92.7	91.5	85.9	92.1	98.4	91.8	92.6	91.6
X(a) Intracranial and intraspinal germ-cell tumor	90.0	90.6	87.5	-	-	98.6	88.1	91.5	88.9
X(c) Malignant gonadal germ cell tumor	96.5	96.3	97.1	100.0	100.0	100.0	98.4	95.8	99.2
XI Other malignant epithelial neo. and melanomas ^f	92.5	88.5	94.4	-	92.1	92.9	93.1	92.5	92.7
XI(b) Thyroid carcinoma	99.3	99.6	99.2	-	-	100.0	99.7	99.2	99.7
XI(d) Malignant melanoma	94.5	92.8	95.8	-	88.5 ^d	96.5	95.5	94.9	94.0

^a SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG). [Expected survival rates](#) are derived from life tables by socio-economic status, geography and race developed by the SEER program.

^b Based on follow-up of patients into 2015.

^c International Classification of Childhood Cancer is based on ICD-O-3. Steliarova-Foucher E, Stiller C, Lacour B, Kaatsch P. International Classification of Childhood Cancer, Third Edition. Cancer. April 1, 2005: Vol 103, No. 7, pg 1457-1467.

^d Classifications are shown for invasive cases only. For ICCC groups and subgroups that include *in situ* behavior, only invasive cases are shown except as noted.

^e The standard error is between 5 and 10 percentage points.

^f The standard error is greater than 10 percentage points.

^f Includes *in situ* urinary bladder tumors.

⁻ Statistic could not be calculated due to fewer than 25 cases during the time period.

Neoplasms are abbreviated as neo.