

Table 24.1
Cancer of the Stomach (Invasive)

Trends in SEER Incidence^{ab} and U.S. Mortality^c Using the Joinpoint Regression Program,
1975-2016 With up to Five Joinpoints, 2000-2016 With up to Three Joinpoints,
Both Sexes by Race/Ethnicity

	JP Trend 1		JP Trend 2		JP Trend 3		JP Trend 4		JP Trend 5		JP Trend 6		AAPC ^d	
	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	2007-16	2012-16
<u>SEER 9 Delay-Adjusted Incidence^a, 1975-2016</u>														
All Races	1975-16	-1.5*											-1.5*	-1.5*
White	1975-97	-2.0*	1997-16	-1.2*									-1.2*	-1.2*
Black	1975-82	1.1	1982-16	-1.6*									-1.6*	-1.6*
<u>SEER 21 Delay-Adjusted Incidence^b, 2000-2016</u>														
All Races	2000-16	-1.0*											-1.0*	-1.0*
White	2000-16	-0.9*											-0.9*	-0.9*
White NH ^{ef}	2000-06	-2.0*	2006-16	-0.8*									-0.8*	-0.8*
Black	2000-16	-1.8*											-1.8*	-1.8*
Black NH ^{ef}	2000-16	-1.8*											-1.8*	-1.8*
API ^e	2000-16	-2.7*											-2.7*	-2.7*
AI/AN ^{eg}	2000-16	-2.1*											-2.1*	-2.1*
Hispanic ^f	2000-16	-1.6*											-1.6*	-1.6*
<u>U.S. Cancer Mortality^c, 1975-2016</u>														
All Races	1975-87	-2.5*	1987-90	-0.3	1990-09	-3.0*	2009-16	-1.8*					-2.1*	-1.8*
White	1975-87	-2.8*	1987-90	-0.8	1990-08	-3.2*	2008-16	-1.9*					-2.0*	-1.9*
Black	1975-95	-1.3*	1995-16	-3.2*									-3.2*	-3.2*
<u>U.S. Cancer Mortality^c, 2000-2016</u>														
All Races	2000-08	-3.2*	2008-16	-1.9*									-2.1*	-1.9*
White	2000-08	-3.3*	2008-16	-1.9*									-2.0*	-1.9*
White NH ^{ef}	2000-08	-3.6*	2008-16	-2.5*									-2.6*	-2.5*
Black	2000-16	-3.2*											-3.2*	-3.2*
Black NH ^{ef}	2000-16	-3.1*											-3.1*	-3.1*
API ^e	2000-14	-3.9*	2014-16	1.4									-2.8*	-1.3
AI/AN ^{eg}	2000-16	-3.2*											-3.2*	-3.2*
Hispanic ^f	2000-16	-2.4*											-2.4*	-2.4*

Joinpoint Regression Program Version 4.7, February 2019, National Cancer Institute. (<https://surveillance.cancer.gov/joinpoint/>).

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^a Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

^b Trends are from the SEER 21 areas (SEER 9 areas, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, Georgia excluding ATL/RG, Idaho, New York and Massachusetts).

^c Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

^d The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

^e API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic

^f Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

^g Data for American Indian/Alaska Native are based on the Purchased/Referred Care Delivery Area (PRCDA) counties.

* The APC/AAPC is significantly different from zero (p<.05).

- Joinpoint regression line analysis could not be performed on data series.

Table 24.2
Cancer of the Stomach (Invasive)

Trends in SEER Incidence^{ab} and U.S. Mortality^c Using the Joinpoint Regression Program,
1975-2016 With up to Five Joinpoints, 2000-2016 With up to Three Joinpoints,
Males by Race/Ethnicity

	JP Trend 1		JP Trend 2		JP Trend 3		JP Trend 4		JP Trend 5		JP Trend 6		AAPC ^d	
	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	2007-16	2012-16
<u>SEER 9 Delay-Adjusted Incidence^a, 1975-2016</u>														
All Races	1975-80	-0.2	1980-16	-1.8*									-1.8*	-1.8*
White	1975-16	-1.8*											-1.8*	-1.8*
Black	1975-82	2.4	1982-16	-2.1*									-2.1*	-2.1*
<u>SEER 21 Delay-Adjusted Incidence^b, 2000-2016</u>														
All Races	2000-16	-1.4*											-1.4*	-1.4*
White	2000-06	-2.1*	2006-14	-0.7*	2014-16	-3.6*							-1.4*	-2.2*
White NH ^{ef}	2000-06	-2.6*	2006-14	-0.7*	2014-16	-4.1*							-1.5*	-2.4*
Black	2000-16	-2.0*											-2.0*	-2.0*
Black NH ^{ef}	2000-08	-0.8	2008-11	-6.3	2011-16	0.0							-2.2	0.0
API ^e	2000-16	-2.8*											-2.8*	-2.8*
AI/AN ^{eg}	2000-16	-2.5*											-2.5*	-2.5*
Hispanic ^f	2000-16	-2.0*											-2.0*	-2.0*
<u>U.S. Cancer Mortality^c, 1975-2016</u>														
All Races	1975-87	-2.3*	1987-91	-1.0	1991-09	-3.4*	2009-16	-2.1*					-2.4*	-2.1*
White	1975-93	-2.3*	1993-08	-3.6*	2008-16	-2.2*							-2.4*	-2.2*
Black	1975-93	-1.2*	1993-16	-3.2*									-3.2*	-3.2*
<u>U.S. Cancer Mortality^c, 2000-2016</u>														
All Races	2000-11	-3.2*	2011-16	-1.7*									-2.4*	-1.7*
White	2000-08	-3.6*	2008-16	-2.3*									-2.4*	-2.3*
White NH ^{ef}	2000-09	-3.8*	2009-16	-2.6*									-2.9*	-2.6*
Black	2000-16	-3.1*											-3.1*	-3.1*
Black NH ^{ef}	2000-16	-3.0*											-3.0*	-3.0*
API ^e	2000-16	-3.7*											-3.7*	-3.7*
AI/AN ^{eg}	2000-16	-3.3*											-3.3*	-3.3*
Hispanic ^f	2000-16	-2.9*											-2.9*	-2.9*

Joinpoint Regression Program Version 4.7, February 2019, National Cancer Institute. (<https://surveillance.cancer.gov/joinpoint/>).

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^a Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

^b Trends are from the SEER 21 areas (SEER 9 areas, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, Georgia excluding ATL/RG, Idaho, New York and Massachusetts).

^c Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

^d The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

^e API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic

^f Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

^g Data for American Indian/Alaska Native are based on the Purchased/Referred Care Delivery Area (PRCDA) counties.

* The APC/AAPC is significantly different from zero ($p < .05$).

- Joinpoint regression line analysis could not be performed on data series.

Table 24.3
Cancer of the Stomach (Invasive)

Trends in SEER Incidence^{ab} and U.S. Mortality^c Using the Joinpoint Regression Program,
1975-2016 With up to Five Joinpoints, 2000-2016 With up to Three Joinpoints,
Females by Race/Ethnicity

	JP Trend 1		JP Trend 2		JP Trend 3		JP Trend 4		JP Trend 5		JP Trend 6		AAPC ^d	
	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	2007-16	2012-16
<u>SEER 9 Delay-Adjusted Incidence^a, 1975-2016</u>														
All Races	1975-95	-1.9*	1995-16	-0.9*									-0.9*	-0.9*
White	1975-95	-2.4*	1995-16	-0.9*									-0.9*	-0.9*
Black	1975-16	-0.9*											-0.9*	-0.9*
<u>SEER 21 Delay-Adjusted Incidence^b, 2000-2016</u>														
All Races	2000-16	-0.6*											-0.6*	-0.6*
White	2000-16	-0.6*											-0.6*	-0.6*
White NH ^{ef}	2000-16	-1.0*											-1.0*	-1.0*
Black	2000-16	-1.6*											-1.6*	-1.6*
Black NH ^{ef}	2000-16	-1.6*											-1.6*	-1.6*
API ^e	2000-16	-2.6*											-2.6*	-2.6*
AI/AN ^{eg}	2000-16	-1.9											-1.9	-1.9
Hispanic ^f	2000-16	-1.2*											-1.2*	-1.2*
<u>U.S. Cancer Mortality^c, 1975-2016</u>														
All Races	1975-87	-2.8*	1987-90	-0.4	1990-11	-2.7*	2011-16	-1.4*					-1.9*	-1.4*
White	1975-87	-3.1*	1987-90	-1.0	1990-08	-2.9*	2008-16	-1.6*					-1.7*	-1.6*
Black	1975-98	-1.2*	1998-16	-3.5*									-3.5*	-3.5*
<u>U.S. Cancer Mortality^c, 2000-2016</u>														
All Races	2000-08	-3.1*	2008-16	-1.7*									-1.9*	-1.7*
White	2000-08	-3.1*	2008-16	-1.5*									-1.7*	-1.5*
White NH ^{ef}	2000-16	-3.0*											-3.0*	-3.0*
Black	2000-16	-3.6*											-3.6*	-3.6*
Black NH ^{ef}	2000-16	-3.5*											-3.5*	-3.5*
API ^e	2000-16	-3.4*											-3.4*	-3.4*
AI/AN ^{eg}	2000-16	-3.5*											-3.5*	-3.5*
Hispanic ^f	2000-16	-1.9*											-1.9*	-1.9*

Joinpoint Regression Program Version 4.7, February 2019, National Cancer Institute. (<https://surveillance.cancer.gov/joinpoint/>).

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^a Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

^b Trends are from the SEER 21 areas (SEER 9 areas, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, Georgia excluding ATL/RG, Idaho, New York and Massachusetts).

^c Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

^d The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

^e API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic

^f Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

^g Data for American Indian/Alaska Native are based on the Purchased/Referred Care Delivery Area (PRCDA) counties.

* The APC/AAPC is significantly different from zero ($p < .05$).

- Joinpoint regression line analysis could not be performed on data series.

Table 24.4
Cancer of the Stomach (Invasive)

Delay-adjusted SEER Incidence^a Rates by Year, Race and Sex

Year of Diagnosis:	All Races			Whites			Blacks		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
1975-1979	11.78	17.02	8.05	10.64	15.50	7.22	17.30	24.86	11.72
1980	11.29	17.13	7.17	10.32	15.68	6.56	15.12	26.32	7.32
1981	11.14	16.60	7.39	10.05	15.07	6.68	17.60	27.84	10.39
1982	10.92	16.29	7.12	9.71	14.58	6.34	19.03	31.17	10.33
1983	10.94	15.86	7.47	9.54	13.69	6.59	18.08	27.78	11.29
1984	10.57	15.32	7.31	9.42	13.90	6.41	16.59	21.84	12.53
1985	10.22	15.09	6.87	8.92	13.32	5.96	16.44	23.50	11.34
1986	10.21	15.02	6.81	9.06	13.53	5.88	16.47	23.17	11.88
1987	10.20	15.06	6.81	8.98	13.26	5.96	17.06	26.68	10.69
1988	10.20	15.50	6.47	8.81	13.58	5.48	16.22	24.75	10.09
1989	10.01	15.15	6.34	8.74	13.47	5.36	15.78	23.16	10.66
1990	9.28	13.74	6.19	7.95	11.91	5.20	14.44	21.78	9.43
1991	9.71	14.20	6.58	8.30	12.32	5.52	16.49	25.77	10.42
1992	9.20	13.37	6.16	7.97	11.68	5.25	14.02	19.90	9.84
1993	9.03	13.61	5.80	7.67	11.61	4.88	14.44	23.70	8.49
1994	9.01	13.60	5.70	7.69	11.75	4.76	15.03	24.74	8.76
1995	8.34	12.58	5.23	7.24	11.23	4.34	13.41	18.97	9.58
1996	8.49	12.61	5.53	7.32	11.03	4.68	14.48	22.59	8.91
1997	8.61	12.71	5.63	7.15	10.79	4.50	15.73	22.35	11.01
1998	8.59	12.21	5.94	7.18	10.52	4.76	14.98	20.16	11.45
1999	8.58	12.06	6.02	7.37	10.50	5.01	13.18	18.26	10.08
2000	8.11	11.73	5.42	6.79	9.87	4.47	12.95	19.97	8.25
2001	7.79	11.29	5.14	6.50	9.69	4.05	13.05	18.60	9.01
2002	7.98	11.30	5.50	6.70	9.88	4.32	13.11	15.68	11.06
2003	7.81	10.88	5.48	6.59	9.33	4.44	13.43	18.52	10.04
2004	7.91	11.33	5.32	6.97	10.00	4.61	11.28	17.19	7.45
2005	7.49	10.71	5.02	6.46	9.23	4.26	11.56	18.21	7.43
2006	7.60	10.71	5.20	6.53	9.33	4.30	12.61	17.16	9.36
2007	7.29	10.51	4.80	6.22	9.18	3.85	11.59	17.77	7.61
2008	7.26	10.18	4.98	6.09	8.81	3.87	13.01	18.46	9.58
2009	7.46	10.48	5.13	6.30	9.22	3.96	11.76	15.72	9.38
2010	7.04	9.64	4.96	6.11	8.65	4.00	10.48	13.68	8.30
2011	7.27	9.95	5.14	6.30	8.87	4.21	11.67	14.31	9.71
2012	7.09	9.45	5.24	6.11	8.31	4.36	10.77	15.01	8.05
2013	6.84	9.27	4.89	5.88	8.21	3.96	10.07	13.77	7.37
2014	6.68	9.38	4.48	5.69	8.19	3.59	10.23	13.61	7.71
2015	6.60	9.02	4.69	5.64	7.84	3.85	10.65	13.65	8.50
2016	6.66	8.68	5.10	5.76	7.62	4.27	9.83	13.90	7.33

Reference: [Cancer Incidence Rates Adjusted for Reporting Delay](#).

Individual rates for diagnosis years 1975-1979 available in web browser version of [Cancer Statistics Review](#).

SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

Delay-adjusted rate is not shown for observed rates based on less than 16 cases for the time interval.

^a

-

Table 24.5
Cancer of the Stomach (Invasive)

Age-adjusted SEER Incidence^a Rates by Year, Race and Sex

Year of Diagnosis:	All Races			Whites			Blacks		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
1975-2016	8.62	12.32	5.85	7.52	10.91	4.96	13.46	19.41	9.38
1975-1979	11.78	17.02	8.05	10.64	15.50	7.22	17.30	24.86	11.72
1980	11.29	17.13	7.17	10.32	15.68	6.56	15.12	26.32	7.32
1981	11.14	16.60	7.39	10.05	15.07	6.68	17.60	27.84	10.39
1982	10.92	16.29	7.12	9.71	14.58	6.34	19.03	31.17	10.33
1983	10.94	15.86	7.47	9.54	13.69	6.59	18.08	27.78	11.29
1984	10.57	15.32	7.31	9.42	13.90	6.41	16.59	21.84	12.53
1985	10.22	15.09	6.87	8.92	13.32	5.96	16.44	23.50	11.34
1986	10.21	15.02	6.81	9.06	13.53	5.88	16.47	23.17	11.88
1987	10.20	15.06	6.81	8.98	13.26	5.96	17.06	26.68	10.69
1988	10.20	15.50	6.47	8.81	13.58	5.48	16.22	24.75	10.09
1989	10.01	15.15	6.34	8.74	13.47	5.36	15.78	23.16	10.66
1990	9.28	13.74	6.19	7.95	11.91	5.20	14.44	21.78	9.43
1991	9.71	14.20	6.58	8.30	12.32	5.52	16.49	25.77	10.42
1992	9.20	13.37	6.16	7.97	11.68	5.25	14.02	19.90	9.84
1993	9.03	13.61	5.80	7.67	11.61	4.88	14.44	23.70	8.49
1994	9.01	13.60	5.70	7.69	11.75	4.76	15.03	24.74	8.76
1995	8.34	12.58	5.23	7.24	11.23	4.34	13.41	18.97	9.58
1996	8.49	12.61	5.53	7.32	11.03	4.68	14.48	22.59	8.91
1997	8.61	12.71	5.63	7.15	10.79	4.50	15.73	22.35	11.01
1998	8.59	12.21	5.94	7.18	10.52	4.76	14.98	20.16	11.45
1999	8.58	12.06	6.02	7.37	10.50	5.01	13.18	18.26	10.08
2000	8.11	11.73	5.42	6.79	9.87	4.47	12.95	19.97	8.25
2001	7.79	11.29	5.14	6.50	9.69	4.05	13.05	18.60	9.01
2002	7.98	11.30	5.50	6.70	9.88	4.32	13.11	15.68	11.06
2003	7.81	10.88	5.48	6.59	9.33	4.44	13.43	18.52	10.04
2004	7.91	11.33	5.32	6.97	10.00	4.61	11.28	17.19	7.45
2005	7.49	10.71	5.02	6.46	9.23	4.26	11.56	18.21	7.43
2006	7.60	10.71	5.20	6.53	9.33	4.30	12.61	17.16	9.36
2007	7.29	10.51	4.80	6.22	9.18	3.85	11.59	17.77	7.61
2008	7.26	10.18	4.98	6.09	8.81	3.86	12.99	18.44	9.57
2009	7.45	10.47	5.12	6.29	9.21	3.96	11.74	15.70	9.36
2010	7.03	9.62	4.96	6.10	8.63	4.00	10.45	13.64	8.28
2011	7.25	9.93	5.13	6.29	8.85	4.20	11.63	14.26	9.67
2012	7.07	9.43	5.23	6.10	8.29	4.35	10.72	14.93	8.01
2013	6.83	9.24	4.88	5.87	8.18	3.95	10.01	13.69	7.33
2014	6.65	9.33	4.46	5.66	8.15	3.57	10.15	13.50	7.65
2015	6.55	8.94	4.66	5.59	7.77	3.82	10.51	13.48	8.39
2016	6.55	8.54	5.02	5.65	7.47	4.18	9.58	13.55	7.14

^a Individual rates for diagnosis years 1975-1979 available in web browser version of [Cancer Statistics Review](#).
SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).
Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table 24.6
Cancer of the Stomach (Invasive)

Age-adjusted U.S. Death^a Rates by Year, Race and Sex

Year of Death:	All Races			Whites			Blacks		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
1975-2016	4.93	6.93	3.47	4.44	6.24	3.11	8.91	13.21	6.06
1975-1979	7.99	11.54	5.56	7.43	10.71	5.20	13.23	19.60	8.64
1980	7.36	10.57	5.15	6.82	9.76	4.80	12.16	18.27	7.96
1981	7.25	10.58	4.98	6.68	9.80	4.56	12.48	18.06	8.60
1982	7.02	10.26	4.86	6.46	9.37	4.53	12.17	19.24	7.44
1983	6.83	9.90	4.76	6.24	9.02	4.38	12.27	18.39	8.08
1984	6.78	9.89	4.68	6.21	9.07	4.28	11.92	17.95	7.93
1985	6.51	9.54	4.44	5.96	8.75	4.07	11.54	17.03	7.82
1986	6.36	9.44	4.24	5.77	8.58	3.83	11.39	17.39	7.42
1987	6.18	9.11	4.18	5.62	8.32	3.77	11.29	16.97	7.55
1988	6.08	8.81	4.17	5.51	7.92	3.82	11.05	17.02	7.07
1989	6.21	9.18	4.16	5.60	8.28	3.77	11.49	17.73	7.33
1990	6.07	8.86	4.16	5.45	7.95	3.73	11.46	17.24	7.71
1991	6.00	8.86	4.01	5.39	7.97	3.59	11.55	17.64	7.55
1992	5.63	8.14	3.90	5.01	7.30	3.43	10.71	15.55	7.53
1993	5.62	8.19	3.83	5.03	7.34	3.41	10.70	16.26	7.11
1994	5.41	7.87	3.70	4.81	7.04	3.26	10.26	15.37	6.97
1995	5.35	7.70	3.71	4.72	6.89	3.21	10.36	14.89	7.37
1996	5.13	7.43	3.54	4.53	6.60	3.10	9.99	15.09	6.80
1997	4.94	7.16	3.37	4.34	6.33	2.92	9.66	14.64	6.50
1998	4.81	6.83	3.39	4.25	6.06	2.97	9.29	13.72	6.42
1999	4.64	6.59	3.25	4.05	5.78	2.79	9.25	13.73	6.47
2000	4.55	6.38	3.24	3.97	5.63	2.79	8.95	12.77	6.49
2001	4.37	6.11	3.10	3.84	5.39	2.70	8.32	12.06	5.86
2002	4.26	5.98	3.01	3.71	5.29	2.56	8.48	12.20	6.15
2003	4.15	5.75	2.98	3.65	5.05	2.60	7.89	11.56	5.48
2004	4.01	5.59	2.87	3.52	4.91	2.50	7.61	11.31	5.26
2005	3.82	5.29	2.74	3.32	4.59	2.38	7.29	10.89	5.02
2006	3.69	5.08	2.68	3.22	4.47	2.29	6.90	9.97	4.96
2007	3.64	5.05	2.60	3.17	4.39	2.24	6.94	10.52	4.69
2008	3.55	4.87	2.52	3.07	4.23	2.16	6.91	10.03	4.90
2009	3.43	4.71	2.45	2.98	4.09	2.11	6.41	9.41	4.41
2010	3.42	4.64	2.49	2.99	4.03	2.19	6.33	9.32	4.37
2011	3.25	4.42	2.35	2.83	3.87	2.02	6.11	8.79	4.33
2012	3.22	4.31	2.37	2.80	3.76	2.02	5.93	8.42	4.23
2013	3.17	4.34	2.26	2.80	3.82	1.99	5.63	8.23	3.83
2014	3.12	4.22	2.25	2.74	3.69	1.98	5.47	8.16	3.66
2015	3.07	4.11	2.25	2.69	3.58	1.98	5.42	8.06	3.63
2016	3.02	4.04	2.20	2.61	3.46	1.92	5.32	7.91	3.52

^a Individual rates for years of death 1975-1979 available in web browser version of [Cancer Statistics Review](#).
US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.
Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table 24.7
Cancer of the Stomach (Invasive)SEER Incidence^a and U.S. Death^b Rates, Age-Adjusted and Age-Specific Rates, by Race and Sex

	All Races			Whites			Blacks		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
<u>SEER Incidence</u>									
<u>Age at Diagnosis</u>									
Age-Adjusted Rates, 2012-2016									
All ages	7.4	10.0	5.3	6.6	8.9	4.6	10.3	14.1	7.7
Under 65	3.1	3.9	2.4	2.8	3.5	2.2	4.1	5.2	3.2
65 and over	36.7	52.1	25.3	32.4	46.8	21.6	53.0	75.8	38.9
All ages (WHO world std) ^c	5.3	7.1	3.9	4.7	6.4	3.4	7.3	10.0	5.4
Age-Specific Rates, 2012-2016									
<1	-	-	-	-	-	-	-	-	-
1-4	-	-	-	-	-	-	-	-	-
5-9	-	-	-	-	-	-	-	-	-
10-14	-	-	-	-	-	-	-	-	-
15-19	0.1	0.1	-	0.1	-	-	-	-	-
20-24	0.3	0.3	0.3	0.3	0.2	0.3	0.3	-	-
25-29	0.5	0.5	0.5	0.5	0.5	0.4	0.4	-	-
30-34	1.1	1.1	1.1	1.1	1.1	1.1	1.0	1.3	0.8
35-39	2.0	2.0	2.1	1.9	1.9	1.9	2.0	1.8	2.2
40-44	3.3	3.7	2.9	3.1	3.5	2.7	3.6	3.9	3.4
45-49	5.4	6.1	4.7	4.9	5.6	4.2	6.4	7.3	5.6
50-54	8.4	10.5	6.5	7.5	9.3	5.8	11.2	14.2	8.5
55-59	12.2	16.3	8.4	10.6	14.2	7.0	17.9	24.7	12.1
60-64	17.0	23.2	11.2	14.8	20.4	9.6	25.2	34.9	17.4
65-69	23.8	33.2	15.6	21.3	29.9	13.4	34.4	49.4	23.0
70-74	32.9	46.2	21.8	29.4	41.7	18.8	45.8	64.8	32.5
75-79	42.3	59.8	28.7	36.9	53.1	24.1	60.4	86.6	43.6
80-84	49.0	69.5	34.8	43.3	62.0	30.1	71.1	98.7	55.8
85+	49.1	72.7	37.0	43.0	66.0	31.1	74.7	111.6	59.3
<u>U.S. Mortality</u>									
<u>Age at Death</u>									
Age-Adjusted Rates, 2012-2016									
All ages	3.1	4.2	2.3	2.7	3.7	2.0	5.5	8.2	3.8
Under 65	1.2	1.5	0.9	1.0	1.3	0.8	1.9	2.6	1.2
65 and over	16.6	22.9	11.9	14.5	20.0	10.3	30.8	46.3	21.2
All ages (WHO world std) ^c	1.9	2.5	1.3	1.6	2.1	1.2	3.2	4.8	2.1
Age-Specific Rates, 2012-2016									
<1	-	-	-	-	-	-	-	-	-
1-4	-	-	-	-	-	-	-	-	-
5-9	-	-	-	-	-	-	-	-	-
10-14	-	-	-	-	-	-	-	-	-
15-19	0.0	-	-	-	-	-	-	-	-
20-24	0.1	0.1	0.1	0.1	0.1	0.1	-	-	-
25-29	0.2	0.2	0.2	0.2	0.2	0.1	0.2	0.2	0.2
30-34	0.4	0.4	0.4	0.4	0.4	0.4	0.6	0.7	0.4
35-39	0.8	0.8	0.8	0.7	0.8	0.7	0.9	1.0	0.9
40-44	1.3	1.4	1.1	1.2	1.4	1.0	1.5	1.8	1.3
45-49	2.0	2.3	1.6	1.7	2.0	1.4	3.1	3.9	2.3
50-54	3.2	4.2	2.3	2.8	3.5	2.1	5.2	7.3	3.3
55-59	4.6	6.3	3.1	4.0	5.3	2.6	8.1	11.8	5.0
60-64	6.3	8.8	4.0	5.4	7.4	3.4	11.5	18.1	6.2
65-69	8.7	12.0	5.7	7.5	10.3	4.9	16.2	24.8	9.5
70-74	12.4	17.4	8.1	10.7	14.8	7.2	24.2	38.9	13.6
75-79	18.0	24.8	12.7	15.8	21.6	11.1	32.9	49.5	22.1
80-84	23.9	33.2	17.3	20.8	29.2	14.8	44.1	66.2	31.7
85+	31.5	43.2	25.5	27.8	38.5	22.2	57.9	80.3	48.6

^a SEER 21 areas. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.

^b US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

^c Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.

- Rates are per 100,000 and are age-adjusted to the world (WHO 2000-2025) standard million.

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table 24.8
Cancer of the Stomach (Invasive)

5-Year Relative and Period Survival (Percent) by Race, Sex, Diagnosis Year, Stage and Age

	All Races			Whites			Blacks		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
5-Year Relative Survival (Percent)									
Year of Diagnosis:									
1960-1963 ^a	-	-	-	11	10	13	8	5	14
1970-1973 ^a	-	-	-	13	12	14	13	15	10
1975-1977 ^b	15.2	14.4	16.5	14.1	13.1	15.8	16.1	16.1	16.1
1978-1980 ^b	16.1	14.3	18.9	15.4	13.6	18.2	16.4	15.6	18.0
1981-1983 ^b	17.3	16.4	18.7	16.2	15.4	17.5	16.5	16.2	17.1
1984-1986 ^b	17.6	15.6	20.9	16.8	14.3	20.9	19.1	16.9	22.1
1987-1989 ^b	19.7	17.0	24.3	18.3	15.5	23.1	18.8	16.6	22.5
1990-1992 ^b	20.9	18.1	25.4	18.8	16.1	23.2	22.7	22.0	23.7
1993-1995 ^b	21.0	20.1	22.6	20.0	18.9	21.9	19.3	17.5	21.9
1996-1998 ^b	22.3	21.1	24.3	20.4	18.7	23.3	22.6	20.4	25.3
1999-2001 ^b	24.1	23.4	25.2	22.3	21.1	24.4	23.1	24.9	20.6
2002-2004 ^b	27.5	25.2	31.1	26.1	24.1	29.5	29.2	24.8	34.5
2005-2008 ^b	29.6	27.7	32.6	28.7	27.5	30.9	28.6	22.4	37.5
2009-2015 ^b	32.1 ^f	29.9 ^f	35.7 ^f	31.2 ^f	29.0 ^f	35.0 ^f	32.7 ^f	27.0 ^f	39.5 ^f
5-Year Period Survival (Percent) ^{cd}									
2015	31.8	28.7	36.5	30.5	27.3	35.9	33.0	27.9	39.3
Stage Distribution (%) 2009-2015 ^{ce}									
All Stages									
Number of cases	37,318	22,651	14,667	26,032	16,351	9,681	5,140	2,857	2,283
Percent	100%	100%	100%	100%	100%	100%	100%	100%	100%
Localized	28	25	33	27	24	33	29	25	34
Regional	26	28	23	26	28	22	25	26	24
Distant	36	38	33	37	39	34	36	40	31
Unstaged/Unknown	10	9	12	10	8	12	9	8	11
5-Year Relative Survival (Percent), 2009-2015 ^c									
Age at Diagnosis:									
Ages <45	34.0	30.0	38.3	33.0	28.5	38.4	35.7	32.2	38.8
Ages 45-54	35.4	31.7	41.6	33.8	30.4	40.1	37.7	30.3	48.5
Ages 55-64	34.7	30.7	42.8	33.3	29.7	41.8	35.4	28.1	47.3
Ages 65-74	34.5	30.7	41.6	32.6	29.3	39.4	32.8	25.2	43.8
Ages 75+	23.9	22.6	25.1	22.9	21.1	24.9	22.4	19.6	24.4
Ages <65	34.8	30.9	41.4	33.4	29.7	40.5	36.2	29.4	46.2
Ages 65+	28.8	26.8	31.6	27.4	25.4	30.6	27.6	22.9	32.4
Stage ^e :									
All Stages	31.5	28.7	35.7	30.1	27.4	34.7	32.0	26.5	38.9
Localized	68.8	64.8	73.4	67.6	64.3	71.6	66.4	56.2	75.8
Regional	31.0	30.8	31.5	28.9	28.6	29.6	33.4	33.0	33.8
Distant	5.3	5.4	5.0	5.2	5.5	4.4	6.2	5.1	7.8
Unstaged/Unknown	23.0	20.1	26.4	23.0	18.8	28.2	21.5	14.9	27.0

^a Based on End Results data from a series of hospital registries and one population-based registry.^b SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta).Based on follow-up of patients into 2016. [Expected survival rates](#) are derived from the U.S. Annual Life Tables.^c SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG). Expected survival rates are derived from life tables by socio-economic status, geography and race developed by the SEER program.^d Based on follow-up of patients into 2016.

Period survival provides a 2015 estimate of survival by piecing together the most recent conditional survival estimates from several cohorts. It is computed here using three year calendar blocks (2010-2012: 0-1 year survival), (2009-2011: 1-2 year survival), (2008-2010: 2-3 year survival), (2007-2009: 3-4 year survival), (2006-2008: 4-5 years survival).

^e Stage at diagnosis is classified using SEER Summary Stage 2000. Stage distribution percentages may not sum to 100 due to rounding.^f The difference between 1975-1977 and 2009-2015 is statistically significant (p<.05).^g The standard error is between 5 and 10 percentage points.^h The standard error is greater than 10 percentage points.

- Statistic could not be calculated due to fewer than 25 cases during the time period.

Table 24.9
Cancer of the Stomach (Invasive)
SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Males and Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
Survival Time																									
1-year	38.7	41.5	43.8	44.8	45.8	45.5	45.4	47.9	46.6	47.1	48.7	49.9	50.5	53.1	50.1	52.4	51.9	55.7	55.8	54.6	58.1	57.6	59.6	60.3	58.1
2-year	24.8	26.7	29.2	31.2	31.4	32.0	32.6	33.7	34.0	32.7	33.4	35.3	38.4	38.5	36.4	40.0	39.4	41.9	40.5	40.3	43.5	45.1	44.6	45.3	
3-year	19.7	21.0	23.4	25.1	26.5	26.8	26.3	28.4	27.7	26.9	29.1	30.3	33.5	33.0	31.6	34.5	33.7	36.6	36.1	35.2	36.2	38.5	38.0		
4-year	17.3	18.3	20.6	22.5	23.3	23.7	23.4	25.2	24.9	24.6	27.6	27.1	30.5	30.1	28.0	33.2	31.1	34.0	33.6	32.1	33.0	35.5			
5-year	15.6	16.6	19.3	20.8	21.9	22.2	21.9	22.8	23.1	22.9	26.5	26.2	28.0	28.3	25.9	31.3	29.0	32.0	31.6	30.5	30.9				
6-year	14.6	15.9	18.4	19.3	20.7	20.6	20.8	21.9	21.2	22.1	25.3	24.5	26.6	26.6	24.8	29.9	27.0	31.0	30.5	29.5					
7-year	14.0	15.1	17.6	18.4	19.6	20.0	20.1	21.1	20.1	21.3	24.3	23.9	25.8	25.4	23.9	28.9	25.8	29.9	29.5						
8-year	13.4	14.4	17.0	17.5	19.4	19.8	19.4	20.6	19.1	20.6	22.9	22.6	24.6	24.5	23.4	28.3	24.9	29.1							
9-year	13.1	14.1	16.5	16.9	19.1	19.3	19.3	19.7	18.6	20.0	22.4	21.7	24.2	23.7	22.2	27.7	24.1								
10-year	12.7	13.9	15.8	16.1	18.2	18.9	18.3	18.7	18.0	19.5	21.6	21.1	23.6	22.6	21.4	26.3									
11-year	12.3	13.4	15.4	15.8	17.8	17.7	17.8	18.6	17.2	18.5	21.2	20.2	23.0	22.1	21.2										
12-year	12.0	12.9	15.0	15.6	17.2	17.2	17.8	17.8	17.1	18.0	20.5	19.4	22.4	21.1											
13-year	11.6	12.9	14.9	15.2	16.7	16.1	17.8	17.0	15.7	17.5	20.5	19.2	21.8												
14-year	11.2	12.8	14.6	14.9	16.4	15.8	17.3	17.0	15.6	17.4	20.2	18.1													
15-year	10.8	12.3	14.3	14.6	16.3	15.5	17.3	17.0	15.6	17.2	19.3														
16-year	10.5	11.9	14.0	14.1	16.3	14.7	16.6	16.6	14.8	17.1															
17-year	10.2	11.6	13.8	13.8	16.3	14.1	15.6	15.7	14.2																
18-year	9.8	11.4	13.6	13.3	16.1	13.4	15.2	15.5																	
19-year	9.5	11.2	13.2	13.0	15.7	13.1	15.2																		
20-year	9.2	10.8	13.0	12.8	14.7	12.8																			

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). [Expected survival rates](#) are derived from the U.S. Annual Life Tables.

Table 24.10
Cancer of the Stomach (Invasive)

SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Males

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
Survival Time																									
1-year	38.1	41.1	42.8	43.6	46.5	43.6	45.6	49.3	46.4	46.3	49.4	48.1	49.8	53.5	51.7	52.2	50.8	54.1	55.7	54.1	57.6	56.1	58.8	60.2	58.5
2-year	23.8	25.8	27.5	29.3	31.5	28.8	31.0	35.5	33.8	32.8	33.1	32.6	36.4	37.9	37.0	37.7	36.9	39.7	39.3	38.9	42.3	43.1	42.9	44.9	
3-year	18.5	20.0	21.4	23.3	25.9	23.7	24.9	29.5	27.4	27.2	28.6	27.1	31.6	31.0	31.5	33.1	31.6	33.4	34.2	33.7	34.8	36.5	36.5		
4-year	16.1	17.1	18.2	20.7	22.4	21.5	21.6	25.6	24.2	24.5	26.7	24.7	28.1	27.7	27.5	31.6	28.6	30.4	31.9	29.9	31.6	32.5			
5-year	14.5	15.4	16.7	18.8	20.8	20.2	19.9	23.2	22.2	22.4	25.7	23.5	25.5	26.4	25.3	29.8	26.7	29.1	29.3	27.6	29.3				
6-year	13.4	14.6	15.9	17.2	19.7	19.2	18.7	22.4	20.4	21.3	23.8	22.0	24.2	24.4	24.3	28.4	24.9	28.3	28.1	26.8					
7-year	12.6	13.9	15.0	16.2	18.7	18.8	17.8	21.3	18.7	20.2	23.0	21.1	23.8	23.4	23.7	27.4	23.2	27.5	27.1						
8-year	12.3	13.3	14.5	15.2	18.6	18.7	17.6	20.8	17.7	19.4	21.6	20.2	22.4	22.6	23.1	26.9	22.4	27.3							
9-year	12.2	13.0	13.9	14.6	18.5	18.0	17.4	20.1	16.8	18.2	21.4	19.8	22.2	21.8	21.8	26.1	21.9								
10-year	11.8	12.9	13.4	13.6	17.3	17.5	16.4	19.2	16.5	17.9	20.8	18.7	22.2	20.5	21.3	25.0									
11-year	11.4	12.5	12.9	13.4	17.1	15.7	15.8	19.2	15.9	16.6	20.5	17.8	22.2	20.0	20.5										
12-year	11.3	11.8	12.7	13.1	16.6	15.0	15.7	18.4	15.6	16.0	19.9	17.2	21.6	19.2											
13-year	11.0	11.8	12.6	12.7	16.0	14.8	15.7	17.7	14.6	15.5	19.8	16.5	21.1												
14-year	10.4	11.8	12.3	12.0	15.4	14.7	15.2	17.7	14.1	15.1	19.8	15.7													
15-year	10.0	11.3	12.0	11.4	15.4	14.7	15.1	17.7	14.1	14.6	18.7														
16-year	9.8	10.8	11.7	11.2	15.4	13.5	15.1	16.5	14.0	14.4															
17-year	9.5	10.5	11.5	11.1	15.3	12.2	13.5	15.9	13.1																
18-year	8.8	10.4	11.3	10.7	15.0	11.5	12.6	15.9																	
19-year	8.7	10.3	10.9	10.4	14.3	11.5	12.5																		
20-year	8.6	9.9	10.8	10.3	13.3	10.9																			

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). [Expected survival rates](#) are derived from the U.S. Annual Life Tables.

Table 24.11
Cancer of the Stomach (Invasive)

SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
Survival Time																									
1-year	39.7	42.3	45.5	46.9	44.6	48.8	45.0	45.8	46.9	48.5	47.4	52.6	51.7	52.4	47.2	52.7	53.7	58.4	56.0	55.4	58.9	59.9	60.8	60.5	57.4
2-year	26.6	28.1	32.0	34.3	31.3	37.4	35.1	30.9	34.4	32.4	33.9	39.4	41.5	39.5	35.2	43.5	43.4	45.5	42.4	42.5	45.3	48.0	47.4	45.9	
3-year	21.5	22.6	26.7	28.1	27.6	31.7	28.5	26.6	28.3	26.3	29.7	35.0	36.5	36.0	31.8	36.9	36.9	42.0	38.9	37.5	38.3	41.7	40.3		
4-year	19.2	20.1	24.6	25.4	24.7	27.2	26.3	24.5	25.9	24.7	29.0	30.6	34.1	34.0	29.0	35.6	35.0	39.9	36.2	35.3	35.2	40.0			
5-year	17.4	18.7	23.6	24.1	23.7	25.7	25.1	22.2	24.4	23.7	27.6	30.0	31.8	31.6	26.8	33.5	32.8	36.8	35.2	34.7	33.7				
6-year	16.6	18.0	22.5	22.8	22.4	22.9	24.1	21.0	22.3	23.4	27.4	28.2	30.3	30.2	25.5	32.3	30.4	35.5	34.1	33.4					
7-year	16.1	17.2	21.8	22.0	21.2	21.7	23.4	20.8	22.1	22.9	26.6	27.9	28.9	28.4	24.3	31.3	30.0	34.0	33.1						
8-year	15.2	16.3	21.3	21.2	20.7	21.3	22.2	20.2	21.1	22.5	24.8	26.1	27.9	27.4	23.9	30.5	28.7	32.0							
9-year	14.5	15.9	20.8	20.7	19.8	21.3	22.2	18.9	21.1	22.5	23.7	24.4	27.0	26.9	22.9	30.1	27.6								
10-year	14.3	15.4	19.8	20.2	19.6	21.1	21.2	17.7	20.1	21.9	22.9	24.3	25.2	25.9	21.4	28.1									
11-year	13.7	15.0	19.6	19.7	19.0	21.0	20.8	17.6	19.1	21.4	22.1	23.8	24.3	25.4	21.4										
12-year	13.2	14.6	18.9	19.6	18.2	20.7	20.8	16.8	19.0	20.9	21.5	22.7	23.6	24.2											
13-year	12.6	14.6	18.7	19.3	17.8	18.0	20.8	15.8	17.3	20.8	21.4	22.6	23.1												
14-year	12.4	14.4	18.5	19.3	17.8	17.5	20.4	15.7	17.3	20.8	20.8	21.5													
15-year	12.2	13.9	18.1	19.3	17.8	16.5	20.4	15.7	17.3	20.8	20.2														
16-year	11.7	13.6	17.8	18.8	17.8	16.4	18.8	15.7	16.0	20.8															
17-year	11.4	13.4	17.5	18.3	17.8	16.4	18.7	15.1	15.4																
18-year	11.4	13.2	17.5	17.6	17.8	16.3	18.7	14.8																	
19-year	10.9	12.8	17.2	17.3	17.8	15.7	18.7																		
20-year	10.2	12.4	16.4	16.8	17.0	15.6																			

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). [Expected survival rates](#) are derived from the U.S. Annual Life Tables.

Table 24.12
Cancer of the Stomach (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years,
Lifetime Risk of Being Diagnosed with Cancer Given Alive and Cancer-Free at Current Age, and
Lifetime Risk of Dying from Cancer Given Alive at Current Age
Both Sexes, 2014-2016 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer
		+10 yrs	+20 yrs	+30 yrs	Ever				+10 yrs	+20 yrs	+30 yrs	Ever	
All Races	0	0.00	0.00	0.00	0.86	0.37	Asian/ Pacific Islander	0	0.00	0.00	0.01	1.64	0.99
	10	0.00	0.00	0.02	0.86	0.38		10	0.00	0.01	0.03	1.65	1.00
	20	0.00	0.02	0.06	0.87	0.38		20	0.01	0.03	0.08	1.66	1.00
	30	0.02	0.06	0.16	0.87	0.38		30	0.02	0.08	0.21	1.66	1.00
	40	0.04	0.14	0.32	0.86	0.38		40	0.06	0.19	0.44	1.64	1.00
	50	0.10	0.28	0.55	0.84	0.37		50	0.14	0.39	0.84	1.61	0.98
	60	0.19	0.47	0.71	0.78	0.36		60	0.26	0.72	1.22	1.51	0.95
	70	0.32	0.58	-	0.67	0.32		70	0.49	1.02	-	1.34	0.90
80	0.36	-	-	0.46	0.26	80	0.63	-	-	1.00	0.78		
White	0	0.00	0.00	0.00	0.75	0.32	American Indian/ Alaska Native ^a	0	0.00	0.00	0.00	0.95	0.52
	10	0.00	0.00	0.02	0.76	0.33		10	0.00	0.00	0.03	0.96	0.53
	20	0.00	0.02	0.06	0.76	0.33		20	0.00	0.03	0.09	0.96	0.53
	30	0.02	0.05	0.14	0.76	0.33		30	0.03	0.09	0.19	0.98	0.54
	40	0.04	0.13	0.28	0.76	0.33		40	0.06	0.17	0.35	0.98	0.54
	50	0.09	0.25	0.48	0.73	0.32		50	0.11	0.30	0.52	0.95	0.54
	60	0.17	0.42	0.62	0.68	0.31		60	0.20	0.44	0.71	0.89	0.52
	70	0.28	0.51	-	0.58	0.28		70	0.28	0.58	-	0.79	0.48
80	0.31	-	-	0.40	0.23	80	0.41	-	-	0.69	0.40		
Black	0	0.00	0.00	0.00	1.10	0.60	Hispanic ^b	0	0.00	0.00	0.01	1.36	0.73
	10	0.00	0.00	0.02	1.12	0.61		10	0.00	0.01	0.04	1.37	0.73
	20	0.00	0.02	0.07	1.12	0.61		20	0.01	0.03	0.10	1.38	0.73
	30	0.01	0.06	0.20	1.13	0.62		30	0.03	0.09	0.23	1.38	0.74
	40	0.05	0.19	0.43	1.14	0.62		40	0.06	0.20	0.45	1.36	0.73
	50	0.14	0.39	0.72	1.13	0.62		50	0.14	0.39	0.78	1.32	0.72
	60	0.27	0.64	0.96	1.07	0.61		60	0.26	0.67	1.06	1.23	0.68
	70	0.43	0.82	-	0.95	0.58		70	0.44	0.88	-	1.07	0.63
80	0.54	-	-	0.73	0.50	80	0.54	-	-	0.77	0.51		

Devcan 6.7.7, April 2019, National Cancer Institute (<https://surveillance.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 21 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, Georgia excluding ATL/RG, Idaho, New York and Massachusetts). Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the PRCA (Purchased/Referred Care Delivery Areas) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

- Statistic could not be calculated.

A percent of 0.00 represents a value that is below 0.005.

Table 24.13
Cancer of the Stomach (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years,
Lifetime Risk of Being Diagnosed with Cancer Given Alive and Cancer-Free at Current Age, and
Lifetime Risk of Dying from Cancer Given Alive at Current Age
Males, 2014-2016 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer
		+10 yrs	+20 yrs	+30 yrs	Ever				+10 yrs	+20 yrs	+30 yrs	Ever	
All Races	0	0.00	0.00	0.00	1.07	0.45	Asian/ Pacific Islander	0	0.00	0.00	0.01	2.01	1.16
	10	0.00	0.00	0.02	1.08	0.45		10	0.00	0.01	0.03	2.02	1.17
	20	0.00	0.02	0.07	1.08	0.45		20	0.01	0.03	0.08	2.03	1.18
	30	0.02	0.06	0.19	1.09	0.46		30	0.02	0.08	0.25	2.03	1.18
	40	0.05	0.17	0.41	1.09	0.46		40	0.06	0.23	0.58	2.03	1.18
	50	0.13	0.37	0.72	1.08	0.46		50	0.18	0.52	1.11	2.00	1.18
	60	0.26	0.64	0.94	1.02	0.44		60	0.36	0.97	1.58	1.89	1.14
	70	0.44	0.78	-	0.88	0.40		70	0.67	1.33	-	1.66	1.08
80	0.49	-	-	0.61	0.33	80	0.82	-	-	1.22	0.94		
White	0	0.00	0.00	0.00	0.95	0.39	American Indian/ Alaska Native ^a	0	0.00	0.00	0.00	1.07	0.60
	10	0.00	0.00	0.02	0.96	0.39		10	0.00	0.00	0.03	1.09	0.61
	20	0.00	0.02	0.06	0.96	0.39		20	0.00	0.03	0.11	1.09	0.61
	30	0.02	0.06	0.17	0.97	0.40		30	0.03	0.12	0.25	1.12	0.63
	40	0.04	0.15	0.36	0.97	0.40		40	0.09	0.23	0.48	1.13	0.64
	50	0.11	0.33	0.65	0.96	0.40		50	0.14	0.41	0.70	1.08	0.65
	60	0.23	0.57	0.84	0.90	0.38		60	0.29	0.61	0.92	1.03	0.62
	70	0.39	0.70	-	0.78	0.35		70	0.38	0.73	-	0.86	0.57
80	0.43	-	-	0.54	0.28	80	0.50	-	-	0.68	0.43		
Black	0	0.00	0.00	0.00	1.31	0.75	Hispanic ^b	0	0.00	0.00	0.01	1.61	0.84
	10	0.00	0.00	0.02	1.33	0.77		10	0.00	0.01	0.04	1.63	0.85
	20	0.00	0.02	0.07	1.34	0.77		20	0.01	0.04	0.10	1.63	0.85
	30	0.01	0.07	0.24	1.36	0.78		30	0.03	0.09	0.25	1.64	0.86
	40	0.06	0.23	0.56	1.38	0.80		40	0.06	0.22	0.52	1.64	0.85
	50	0.18	0.52	0.95	1.38	0.81		50	0.16	0.47	0.97	1.61	0.84
	60	0.38	0.86	1.23	1.33	0.80		60	0.32	0.86	1.34	1.53	0.82
	70	0.59	1.06	-	1.19	0.77		70	0.60	1.15	-	1.36	0.76
80	0.72	-	-	0.92	0.65	80	0.72	-	-	0.99	0.63		

Devcan 6.7.7, April 2019, National Cancer Institute (<https://surveillance.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 21 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, Georgia excluding ATL/RG, Idaho, New York and Massachusetts). Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the PRCD (Purchased/Referred Care Delivery Areas) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

- Statistic could not be calculated.

A percent of 0.00 represents a value that is below 0.005.

Table 24.14
Cancer of the Stomach (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years,
Lifetime Risk of Being Diagnosed with Cancer Given Alive and Cancer-Free at Current Age, and
Lifetime Risk of Dying from Cancer Given Alive at Current Age
Females, 2014-2016 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer
		+10 yrs	+20 yrs	+30 yrs	Ever				+10 yrs	+20 yrs	+30 yrs	Ever	
All Races	0	0.00	0.00	0.00	0.66	0.30	Asian/ Pacific Islander	0	0.00	0.00	0.01	1.34	0.85
	10	0.00	0.00	0.02	0.67	0.30		10	0.00	0.01	0.03	1.35	0.86
	20	0.00	0.02	0.06	0.67	0.31		20	0.01	0.03	0.08	1.35	0.86
	30	0.02	0.06	0.13	0.67	0.31		30	0.02	0.07	0.17	1.34	0.86
	40	0.04	0.11	0.23	0.66	0.30		40	0.05	0.15	0.32	1.33	0.85
	50	0.07	0.20	0.38	0.63	0.29		50	0.10	0.27	0.60	1.29	0.83
	60	0.13	0.32	0.51	0.58	0.28		60	0.17	0.51	0.92	1.21	0.80
	70	0.21	0.42	-	0.50	0.26		70	0.35	0.79	-	1.09	0.76
80	0.26	-	-	0.36	0.22	80	0.50	-	-	0.84	0.68		
White	0	0.00	0.00	0.00	0.57	0.26	American Indian/ Alaska Native ^a	0	0.00	0.00	0.00	0.82	0.43
	10	0.00	0.00	0.02	0.57	0.27		10	0.00	0.01	0.03	0.82	0.44
	20	0.00	0.02	0.05	0.57	0.27		20	0.01	0.03	0.07	0.83	0.44
	30	0.02	0.05	0.11	0.57	0.27		30	0.02	0.06	0.14	0.83	0.44
	40	0.03	0.10	0.20	0.56	0.26		40	0.04	0.12	0.22	0.82	0.43
	50	0.06	0.17	0.33	0.53	0.26		50	0.08	0.19	0.36	0.81	0.43
	60	0.11	0.28	0.44	0.49	0.24		60	0.11	0.28	0.53	0.76	0.43
	70	0.18	0.36	-	0.42	0.22		70	0.19	0.46	-	0.72	0.39
80	0.22	-	-	0.30	0.19	80	0.34	-	-	0.67	0.37		
Black	0	0.00	0.00	0.00	0.93	0.47	Hispanic ^b	0	0.00	0.00	0.01	1.16	0.64
	10	0.00	0.00	0.02	0.94	0.48		10	0.00	0.01	0.04	1.17	0.64
	20	0.00	0.02	0.06	0.94	0.48		20	0.01	0.03	0.10	1.17	0.64
	30	0.02	0.06	0.16	0.95	0.48		30	0.03	0.09	0.21	1.17	0.64
	40	0.05	0.15	0.31	0.94	0.48		40	0.06	0.18	0.39	1.14	0.63
	50	0.11	0.28	0.54	0.92	0.48		50	0.12	0.33	0.62	1.09	0.61
	60	0.18	0.46	0.75	0.87	0.47		60	0.21	0.51	0.84	1.00	0.58
	70	0.32	0.65	-	0.79	0.45		70	0.32	0.68	-	0.85	0.53
80	0.43	-	-	0.62	0.42	80	0.42	-	-	0.63	0.43		

Devcan 6.7.7, April 2019, National Cancer Institute (<https://surveillance.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 21 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, Georgia excluding ATL/RG, Idaho, New York and Massachusetts). Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the PRCD (Purchased/Referred Care Delivery Areas) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

- Statistic could not be calculated.

A percent of 0.00 represents a value that is below 0.005.

Table 24.15
Cancer of the Stomach (Invasive)

SEER Incidence and U.S. Mortality
Age-Adjusted Rates and Trends^a
By Race/Ethnicity and Sex

SEER Incidence	SEER 21 Areas ^b			SEER 21 Areas ^b			SEER 9 Areas ^b		
	Rate 2012-2016			Trend 2007-2016 ^c			Trend 2007-2016 ^d		
	Rate per 100,000 persons			AAPC (%)			AAPC (%)		
Race/Ethnicity	Total	Males	Females	Total	Males	Females	Total	Males	Females
All Races	7.4	10.0	5.3	-1.1*	-1.5*	-0.7*	-1.5*	-1.8*	-0.9*
White	6.6	8.9	4.6	-1.0*	-1.6*	-0.7*	-1.3*	-1.8*	-1.0*
White Hispanic ^e	10.8	13.4	8.8	-1.8*	-2.2*	-1.4*	-	-	-
White Non-Hispanic ^e	5.8	8.2	3.8	-1.3*	-1.7*	-1.1*	-	-	-
Black	10.3	14.1	7.7	-1.9*	-2.8	-1.7*	-1.7*	-2.1*	-0.9*
Asian/Pacific Islander	10.9	14.3	8.2	-2.8*	-2.9*	-2.7*	-	-	-
Amer Ind/Alaska Nat ^f	8.7	11.5	6.4	-2.4*	-2.7*	-2.2	-	-	-
Hispanic ^e	10.4	13.0	8.5	-1.7*	-2.1*	-1.3*	-	-	-
<u>U.S. Mortality^g</u>									
Race/Ethnicity	Rate 2012-2016			Trend 2007-2016 ^h					
	Rate per 100,000 persons			AAPC (%)					
	Total	Males	Females	Total	Males	Females			
All Races	3.1	4.2	2.3	-2.1*	-2.4*	-1.9*			
White	2.7	3.7	2.0	-2.0*	-2.4*	-1.7*			
White Hispanic ^e	5.4	6.9	4.3	-2.3*	-2.8*	-1.8*			
White Non-Hispanic ^e	2.4	3.3	1.7	-2.6*	-2.9*	-3.0*			
Black	5.5	8.2	3.8	-3.2*	-3.1*	-3.6*			
Asian/Pacific Islander	5.3	6.8	4.2	-2.8*	-3.7*	-3.4*			
Amer Ind/Alaska Nat									
Total U.S.	3.4	4.6	2.5	-3.5*	-3.9*	-3.4*			
PRCDA Counties	5.0	6.9	3.6	-3.2*	-3.3*	-3.5*			
Non-PRCDA Counties	1.6	2.0	1.3	-2.2*	-4.1*	-0.7			
Hispanic ^e	5.1	6.5	4.0	-2.4*	-2.9*	-1.9*			

The AAPC is the Average Annual Percent Change over the time interval. The AAPCs are calculated by the Joinpoint Regression Program Version 4.7, February 2019, National Cancer Institute.

Statistic not shown. Rate based on less than 16 cases for the time interval.

Trend based on less than 10 cases for at least one year within the time interval.

Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130). Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

The SEER 9 areas are San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta.

The SEER 21 areas comprise the SEER 9 areas plus San Jose-Monterey, Los Angeles, the Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, Georgia excluding ATL/RG, Idaho, New York and Massachusetts.

The 2007-2016 AAPC estimates are based on a Joinpoint analysis with up to 3 Joinpoints over diagnosis years 2000-2016.

The 2007-2016 AAPC estimates are based on a Joinpoint analysis with up to 5 Joinpoints over diagnosis years 1975-2016.

Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

Incidence data for American Indian/Alaska Native are based on the Purchased/Referred Care Delivery Area (PRCDA) counties.

US Mortality Files, National Center for Health Statistics, CDC.

The 2007-2016 mortality AAPCs are based on a Joinpoint analysis using years of death 2000-2016.

The APC is significantly different from zero (p<.05).

Table 24.16
Cancer of the Stomach (Invasive)

Age-Adjusted SEER Incidence Rates^a
By Registry, Race and Sex

	All Races			Whites			Blacks		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
<u>SEER Incidence Rates^a, 2012-2016</u>									
California	7.51	9.89	5.58	6.91	9.14	5.07	8.81	12.18	6.39
Greater Bay Area	7.37	9.70	5.53	6.32	8.34	4.68	9.95	13.52	7.34
San Francisco-Oakland	7.37	9.70	5.54	6.28	8.30	4.63	9.87	13.55	7.22
San Jose-Monterey	7.37	9.70	5.49	6.40	8.42	4.77	10.43	13.54	-
Los Angeles	9.42	12.30	7.16	8.85	11.22	6.93	8.95	12.92	6.36
Greater California	6.67	8.87	4.86	6.31	8.55	4.44	7.94	10.60	5.82
Connecticut	8.02	11.16	5.59	7.33	10.39	4.92	13.35	18.24	10.25
Detroit	7.31	10.07	5.20	6.44	9.07	4.42	10.04	13.55	7.58
Georgia	6.46	8.76	4.69	5.28	7.18	3.76	9.44	13.40	6.89
Atlanta	7.14	9.85	5.16	5.43	7.52	3.80	9.61	14.07	6.95
Rural Georgia	6.64	8.38	5.59	4.29	-	-	10.69	15.22	-
Greater Georgia	6.21	8.37	4.48	5.27	7.13	3.75	9.40	13.03	6.88
Hawaii	8.70	11.23	6.67	4.74	6.35	3.02	-	-	-
Idaho	5.17	7.04	3.50	5.09	6.95	3.43	-	-	-
Iowa	4.63	6.60	3.03	4.44	6.45	2.79	9.24	-	9.34
Kentucky	6.77	9.18	4.85	6.57	8.95	4.64	9.59	13.27	7.12
Louisiana	7.08	10.01	4.73	5.56	8.10	3.48	11.04	15.47	7.87
Massachusetts	6.70	9.43	4.56	6.07	8.65	4.04	11.91	17.28	8.21
New Jersey	7.93	10.77	5.81	7.29	10.08	5.16	10.16	13.87	7.91
New Mexico	6.49	8.24	5.00	6.00	7.64	4.57	7.32	-	-
New York	8.74	11.97	6.25	7.52	10.40	5.26	11.31	15.54	8.64
Seattle-Puget Sound	6.19	8.57	4.20	5.48	7.85	3.47	10.75	14.27	7.91
Utah	4.61	5.79	3.58	4.31	5.53	3.28	-	-	-
SEER 9 Areas ^b	6.72	9.09	4.84	5.77	7.97	3.97	10.17	13.80	7.68
SEER 13 Areas ^b	7.39	9.86	5.42	6.45	8.66	4.64	9.90	13.63	7.39
SEER 18 Areas ^b	7.15	9.60	5.20	6.40	8.68	4.54	9.82	13.50	7.29
SEER 21 Areas ^b	7.38	9.97	5.33	6.55	8.95	4.61	10.28	14.14	7.68

^a Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130)

^b The SEER 9 areas are San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah and Atlanta.

The SEER 13 areas comprise the SEER 9 areas plus San Jose-Monterey, Los Angeles, the Alaska Native Registry and Rural Georgia.

The SEER 18 areas comprise the SEER 13 areas plus California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia

excluding ATL/RG.

The SEER 21 areas comprise the SEER 18 areas plus Idaho, New York and Massachusetts.

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table 24.17
Cancer of the Stomach (Invasive)

Age-Adjusted SEER Death Rates^a
By Registry, Race and Sex

	All Races			Whites			Blacks		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
<u>SEER Death Rates^a, 2012-2016</u>									
California	4.00	5.19	3.07	3.66	4.75	2.77	5.27	7.72	3.50
Greater Bay Area	3.88	5.08	2.97	3.15	4.11	2.41	6.14	8.29	4.51
San Francisco-Oakland	3.91	5.16	2.96	3.11	4.10	2.33	6.54	8.89	4.79
San Jose-Monterey	3.83	4.93	2.99	3.23	4.15	2.55	-	-	-
Los Angeles	5.27	6.95	3.99	5.05	6.57	3.84	5.40	8.08	3.62
Greater California	3.47	4.46	2.66	3.26	4.24	2.44	4.72	7.12	2.86
Connecticut	3.16	4.44	2.18	2.93	4.07	2.06	5.78	8.56	3.83
Detroit	3.45	4.60	2.57	2.81	3.82	2.06	5.45	7.18	4.18
Georgia	3.33	4.58	2.36	2.53	3.38	1.86	5.56	8.70	3.54
Atlanta	3.40	4.62	2.51	2.32	2.78	1.97	5.17	8.58	3.11
Rural Georgia	3.40	-	-	-	-	-	6.02	-	-
Greater Georgia	3.30	4.62	2.26	2.62	3.62	1.81	5.85	8.94	3.78
Hawaii	4.42	5.69	3.49	2.53	2.77	2.33	-	-	-
Idaho	2.35	3.25	1.58	2.35	3.22	1.59	-	-	-
Iowa	2.07	2.93	1.39	1.97	2.79	1.32	6.70	-	-
Kentucky	2.89	4.00	2.02	2.80	3.89	1.94	4.33	5.73	3.31
Louisiana	3.65	4.96	2.60	2.37	3.23	1.72	7.13	10.24	4.88
Massachusetts	3.07	4.16	2.25	2.88	3.92	2.09	4.92	6.48	3.82
New Jersey	3.52	4.98	2.46	3.27	4.63	2.27	5.35	7.94	3.69
New Mexico	3.41	4.24	2.72	3.04	3.75	2.43	-	-	-
New York	3.69	5.15	2.59	3.19	4.40	2.26	5.32	7.77	3.74
Seattle-Puget Sound	2.91	3.85	2.12	2.42	3.31	1.65	5.62	8.74	-
Utah	2.14	2.62	1.74	1.97	2.46	1.57	-	-	-
SEER 9 Areas ^b	3.19	4.25	2.37	2.59	3.44	1.91	5.56	7.89	3.94
SEER 13 Areas ^b	3.71	4.89	2.78	3.12	4.10	2.35	5.49	7.89	3.84
SEER 18 Areas ^b	3.56	4.75	2.64	3.09	4.12	2.28	5.64	8.21	3.84
SEER 21 Areas ^b	3.54	4.76	2.59	3.08	4.14	2.25	5.53	8.04	3.82
Total U.S.	3.12	4.20	2.26	2.73	3.66	1.98	5.54	8.15	3.76

^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b The SEER 9 areas are San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah and Atlanta. The SEER 13 areas comprise the SEER 9 areas plus San Jose-Monterey, Los Angeles, the Alaska Native Registry and Rural Georgia. The SEER 18 areas comprise the SEER 13 areas plus California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG.

The SEER 21 areas comprise the SEER 18 areas plus Idaho, New York and Massachusetts.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 24.18
Cancer of the Stomach (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2012-2016

Males and Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	3.12	0.01		
High Five States				
D.C.	5.08	0.41	01(01,05)	63.02 ^c
Alaska	4.66	0.42	02(01,11)	49.46 ^c
Hawaii	4.42	0.23	03(01,05)	41.63 ^c
California	4.00	0.05	04(02,05)	28.37 ^c
Mississippi	3.79	0.15	05(03,14)	21.66 ^c
Low Five States				
Montana	2.12	0.18	47(29,51)	-31.90 ^c
New Hampshire	2.09	0.16	48(36,51)	-33.14 ^c
Iowa	2.07	0.11	49(39,51)	-33.60 ^c
Nebraska	1.92	0.14	50(41,51)	-38.47 ^c
Wyoming	1.82	0.25	51(34,51)	-41.53 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	3.37	0.11	13(06,19)	7.99	Montana	2.12	0.18	47(29,51)	-31.90 ^c
Alaska	4.66	0.42	02(01,11)	49.46 ^c	Nebraska	1.92	0.14	50(41,51)	-38.47 ^c
Arizona	2.68	0.08	31(21,40)	-14.00	Nevada	2.73	0.14	29(16,42)	-12.42
Arkansas	2.85	0.13	25(15,39)	-8.66	New Hampshire	2.09	0.16	48(36,51)	-33.14 ^c
California	4.00	0.05	04(02,05)	28.37 ^c	New Jersey	3.52	0.08	10(05,15)	12.72
Colorado	2.52	0.10	37(26,45)	-19.21 ^c	New Mexico	3.41	0.17	11(05,22)	9.45
Connecticut	3.16	0.12	16(09,26)	1.44	New York	3.69	0.06	06(05,11)	18.18 ^c
Delaware	2.97	0.23	20(07,43)	-4.92	North Carolina	2.94	0.07	21(15,30)	-5.59
D.C.	5.08	0.41	01(01,05)	63.02 ^c	North Dakota	2.34	0.25	42(17,51)	-24.99 ^c
Florida	2.75	0.05	28(22,36)	-11.72	Ohio	2.66	0.06	33(24,39)	-14.78
Georgia	3.33	0.08	15(09,19)	6.62	Oklahoma	2.71	0.11	30(18,41)	-13.17
Hawaii	4.42	0.23	03(01,05)	41.63 ^c	Oregon	2.63	0.11	35(20,42)	-15.64 ^c
Idaho	2.35	0.17	41(24,51)	-24.54 ^c	Pennsylvania	2.68	0.06	32(23,39)	-14.02
Illinois	3.40	0.07	12(08,17)	9.03	Rhode Island	3.33	0.23	14(04,32)	6.83
Indiana	2.64	0.09	34(22,41)	-15.43 ^c	South Carolina	3.61	0.12	08(04,15)	15.68 ^c
Iowa	2.07	0.11	49(39,51)	-33.60 ^c	South Dakota	2.20	0.22	44(24,51)	-29.47 ^c
Kansas	2.24	0.12	43(35,51)	-28.32 ^c	Tennessee	2.93	0.09	22(15,33)	-6.21
Kentucky	2.89	0.11	23(15,36)	-7.42	Texas	3.60	0.05	09(05,13)	15.31 ^c
Louisiana	3.65	0.12	07(04,15)	16.87 ^c	Utah	2.14	0.14	46(36,51)	-31.25 ^c
Maine	2.16	0.16	45(33,51)	-30.86 ^c	Vermont	2.38	0.25	39(16,51)	-23.63 ^c
Maryland	3.08	0.10	17(12,28)	-1.21	Virginia	2.97	0.08	19(15,30)	-4.86
Massachusetts	3.07	0.09	18(13,26)	-1.48	Washington	2.79	0.09	26(18,38)	-10.69
Michigan	2.86	0.07	24(17,34)	-8.23	West Virginia	2.78	0.15	27(15,42)	-10.88
Minnesota	2.37	0.09	40(33,47)	-23.91 ^c	Wisconsin	2.60	0.09	36(23,43)	-16.77 ^c
Mississippi	3.79	0.15	05(03,14)	21.66 ^c	Wyoming	1.82	0.25	51(34,51)	-41.53 ^c
Missouri	2.47	0.08	38(29,45)	-20.79 ^c					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 24.19
Cancer of the Stomach (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2012-2016

Males

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	4.20	0.02		
High Five States				
D.C.	6.99	0.75	01(01,08)	66.62 ^c
Alaska	5.86	0.68	02(01,24)	39.57 ^c
Hawaii	5.69	0.39	03(01,13)	35.63 ^c
California	5.19	0.08	04(02,09)	23.73 ^c
New York	5.15	0.10	05(02,10)	22.67 ^c
Low Five States				
Montana	2.64	0.30	47(31,51)	-37.03 ^c
Utah	2.62	0.22	48(39,51)	-37.60 ^c
Maine	2.58	0.25	49(39,51)	-38.42 ^c
South Dakota	2.47	0.33	50(31,51)	-41.14 ^c
Wyoming	2.34	0.42	51(30,51)	-44.21 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	4.66	0.20	11(04,20)	11.10	Montana	2.64	0.30	47(31,51)	-37.03 ^c
Alaska	5.86	0.68	02(01,24)	39.57 ^c	Nebraska	2.93	0.25	45(28,51)	-30.14 ^c
Arizona	3.36	0.14	39(27,45)	-19.93 ^c	Nevada	3.44	0.23	38(18,47)	-17.99 ^c
Arkansas	3.69	0.22	30(15,44)	-12.20	New Hampshire	3.02	0.29	43(22,51)	-28.08 ^c
California	5.19	0.08	04(02,09)	23.73 ^c	New Jersey	4.98	0.15	07(03,14)	18.56 ^c
Colorado	3.45	0.17	37(22,45)	-17.77 ^c	New Mexico	4.24	0.29	16(05,37)	0.96
Connecticut	4.44	0.22	14(06,27)	5.78	New York	5.15	0.10	05(02,10)	22.67 ^c
Delaware	3.87	0.40	24(05,48)	-7.89	North Carolina	3.92	0.13	23(15,35)	-6.52
D.C.	6.99	0.75	01(01,08)	66.62 ^c	North Dakota	2.94	0.40	44(17,51)	-30.07 ^c
Florida	3.58	0.08	32(24,40)	-14.63	Ohio	3.56	0.11	34(23,42)	-15.11 ^c
Georgia	4.58	0.15	13(06,19)	9.00	Oklahoma	3.86	0.20	25(14,41)	-7.98
Hawaii	5.69	0.39	03(01,13)	35.63 ^c	Oregon	3.55	0.19	35(19,44)	-15.33 ^c
Idaho	3.25	0.29	41(18,50)	-22.51 ^c	Pennsylvania	3.79	0.10	27(18,37)	-9.67
Illinois	4.66	0.12	12(06,17)	10.96	Rhode Island	4.39	0.40	15(02,40)	4.55
Indiana	3.74	0.15	28(17,41)	-10.99	South Carolina	5.06	0.21	06(02,15)	20.61 ^c
Iowa	2.93	0.19	46(35,51)	-30.29 ^c	South Dakota	2.47	0.33	50(31,51)	-41.14 ^c
Kansas	3.13	0.21	42(27,50)	-25.44 ^c	Tennessee	4.05	0.16	21(13,34)	-3.51
Kentucky	4.00	0.19	22(12,38)	-4.77	Texas	4.78	0.09	10(06,15)	13.81
Louisiana	4.96	0.22	08(02,16)	18.28 ^c	Utah	2.62	0.22	48(39,51)	-37.60 ^c
Maine	2.58	0.25	49(39,51)	-38.42 ^c	Vermont	4.07	0.49	20(02,47)	-3.02
Maryland	4.19	0.18	17(10,31)	-0.15	Virginia	4.08	0.15	19(12,32)	-2.85
Massachusetts	4.16	0.16	18(12,30)	-0.91	Washington	3.68	0.15	31(18,42)	-12.23
Michigan	3.84	0.12	26(16,37)	-8.56	West Virginia	3.57	0.26	33(15,47)	-15.03 ^c
Minnesota	3.31	0.16	40(25,46)	-21.05 ^c	Wisconsin	3.46	0.15	36(23,45)	-17.45 ^c
Mississippi	4.78	0.26	09(02,21)	14.00	Wyoming	2.34	0.42	51(30,51)	-44.21 ^c
Missouri	3.70	0.16	29(17,42)	-11.90					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 24.20
Cancer of the Stomach (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2012-2016

Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	2.26	0.02		
High Five States				
D.C.	3.73	0.47	01(01,12)	64.78 ^c
Hawaii	3.49	0.27	02(01,06)	54.09 ^c
Alaska	3.44	0.50	03(01,26)	52.00 ^c
California	3.07	0.05	04(02,06)	35.41 ^c
Mississippi	2.92	0.18	05(01,14)	28.93 ^c
Low Five States				
Wyoming	1.40	0.29	47(14,51)	-38.40 ^c
Iowa	1.39	0.12	48(38,51)	-38.42 ^c
New Hampshire	1.32	0.18	49(34,51)	-41.81 ^c
Nebraska	1.10	0.14	50(44,51)	-51.52 ^c
Vermont	1.05	0.21	51(37,51)	-53.42 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	2.40	0.13	14(06,29)	5.94	Montana	1.66	0.23	42(13,51)	-26.70 ^c
Alaska	3.44	0.50	03(01,26)	52.00 ^c	Nebraska	1.10	0.14	50(44,51)	-51.52 ^c
Arizona	2.10	0.10	24(12,37)	-7.23	Nevada	2.15	0.17	22(07,44)	-5.09
Arkansas	2.17	0.15	21(07,40)	-4.09	New Hampshire	1.32	0.18	49(34,51)	-41.81 ^c
California	3.07	0.05	04(02,06)	35.41 ^c	New Jersey	2.46	0.09	11(06,22)	8.48
Colorado	1.77	0.11	40(22,47)	-21.83 ^c	New Mexico	2.72	0.21	06(02,25)	19.93 ^c
Connecticut	2.18	0.14	19(07,38)	-3.75	New York	2.59	0.06	09(05,15)	14.31
Delaware	2.29	0.27	16(04,45)	1.11	North Carolina	2.17	0.08	20(12,34)	-4.03
D.C.	3.73	0.47	01(01,12)	64.78 ^c	North Dakota	1.82	0.30	36(06,51)	-19.66 ^c
Florida	2.06	0.05	27(18,35)	-9.23	Ohio	1.95	0.07	31(21,41)	-14.03
Georgia	2.36	0.09	15(07,27)	4.16	Oklahoma	1.78	0.13	37(21,48)	-21.23 ^c
Hawaii	3.49	0.27	02(01,06)	54.09 ^c	Oregon	1.85	0.13	34(18,46)	-18.16 ^c
Idaho	1.58	0.19	44(20,51)	-30.43 ^c	Pennsylvania	1.82	0.06	35(27,44)	-19.43 ^c
Illinois	2.43	0.08	12(07,22)	7.13	Rhode Island	2.57	0.27	10(02,37)	13.32
Indiana	1.78	0.10	39(24,46)	-21.35 ^c	South Carolina	2.42	0.13	13(05,28)	6.65
Iowa	1.39	0.12	48(38,51)	-38.42 ^c	South Dakota	1.88	0.28	33(06,51)	-16.94 ^c
Kansas	1.53	0.13	45(31,50)	-32.28 ^c	Tennessee	2.05	0.10	28(15,40)	-9.66
Kentucky	2.02	0.12	30(14,42)	-10.77	Texas	2.66	0.06	07(04,13)	17.38 ^c
Louisiana	2.60	0.14	08(04,21)	14.70	Utah	1.74	0.17	41(17,49)	-23.06 ^c
Maine	1.78	0.19	38(14,49)	-21.31 ^c	Vermont	1.05	0.21	51(37,51)	-53.42 ^c
Maryland	2.25	0.11	18(09,33)	-0.73	Virginia	2.08	0.09	26(14,38)	-8.26
Massachusetts	2.25	0.10	17(09,33)	-0.54	Washington	2.03	0.10	29(15,40)	-10.18
Michigan	2.09	0.08	25(14,37)	-7.57	West Virginia	2.11	0.18	23(07,44)	-6.75
Minnesota	1.64	0.10	43(31,48)	-27.68 ^c	Wisconsin	1.92	0.10	32(18,44)	-15.15 ^c
Mississippi	2.92	0.18	05(01,14)	28.93 ^c	Wyoming	1.40	0.29	47(14,51)	-38.40 ^c
Missouri	1.49	0.09	46(37,50)	-34.25 ^c					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table 24.21
Cancer of the Stomach (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2016
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <24	0 to <24 ^b	>=24 ^d	Complete ^e
<u>Race</u>	<u>Sex</u>								
All Races	Both Sexes	53,400	25,301	13,855	7,885	3,686	104,126	8,928	113,054
	Males	31,498	14,259	7,709	4,287	1,970	59,722	5,115	64,837
	Females	21,901	11,042	6,147	3,597	1,716	44,404	3,813	48,217
White	Both Sexes	40,321	19,117	10,308	5,880	2,783	78,409	6,979	86,508
	Males	24,380	11,125	5,946	3,236	1,422	46,109	3,972	50,671
	Females	15,942	7,992	4,361	2,645	1,361	32,301	3,007	35,836
Black	Both Sexes	7,845	3,338	1,909	1,092	410	14,594	1,156	15,750
	Males	4,110	1,557	905	528	253	7,352	614	7,966
	Females	3,735	1,781	1,005	564	157	7,241	542	7,784
Asian/ Pacific Islander	Both Sexes	4,048	2,310	1,377	783	400	8,918	+	+
	Males	2,382	1,251	739	441	239	5,052	+	+
	Females	1,666	1,059	638	341	161	3,865	+	+
Hispanic	Both Sexes	7,356	3,456	1,883	996	475	14,166	+	+
	Males	3,866	1,810	987	506	208	7,377	+	+
	Females	3,490	1,646	896	490	267	6,789	+	+

Estimated prevalence percent^a on January 1, 2016, of the SEER population diagnosed in the previous 24 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted ^c	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races	Both Sexes	0.0323	-	0.0001	0.0008	0.0039	0.0130	0.0354	0.0759	0.1462	0.2055	0.0277
	Males	0.0376	-	0.0001	0.0007	0.0037	0.0133	0.0408	0.0959	0.2002	0.2791	0.0355
	Females	0.0271	-	0.0002	0.0010	0.0041	0.0128	0.0302	0.0578	0.1012	0.1603	0.0215
White	Both Sexes	0.0311	-	0.0002	0.0008	0.0038	0.0122	0.0314	0.0679	0.1317	0.1830	0.0249
	Males	0.0369	-	0.0001	0.0006	0.0038	0.0126	0.0364	0.0862	0.1849	0.2506	0.0323
	Females	0.0254	-	0.0002	0.0009	0.0038	0.0117	0.0264	0.0510	0.0864	0.1406	0.0189
Black	Both Sexes	0.0324	-	-	0.0010	0.0036	0.0135	0.0504	0.1072	0.1999	0.2867	0.0375
	Males	0.0340	-	-	0.0007	0.0036	0.0126	0.0557	0.1330	0.2512	0.3732	0.0461
	Females	0.0309	-	-	0.0012	0.0035	0.0144	0.0458	0.0865	0.1642	0.2440	0.0319
Asian/ Pacific Islander	Both Sexes	0.0438	-	0.0003	0.0012	0.0040	0.0185	0.0503	0.1205	0.2676	0.4688	0.0496
	Males	0.0518	-	-	0.0012	0.0033	0.0200	0.0632	0.1610	0.3478	0.6499	0.0650
	Females	0.0364	-	0.0006	0.0011	0.0046	0.0171	0.0391	0.0878	0.2032	0.3523	0.0381
Hispanic	Both Sexes	0.0248	-	0.0001	0.0008	0.0057	0.0173	0.0463	0.1072	0.2083	0.3279	0.0401
	Males	0.0256	-	0.0001	0.0006	0.0057	0.0171	0.0482	0.1220	0.2769	0.3913	0.0477
	Females	0.0240	-	0.0001	0.0010	0.0058	0.0175	0.0444	0.0942	0.1556	0.2887	0.0345

^a US 2016 cancer prevalence counts are based on 2016 cancer prevalence proportions from the SEER 13 Areas (excluding the Alaska Native Registry) and 1/1/2016 US population estimates based on the average of 2015 and 2016 population estimates from the US Bureau of the Census.

^b Prevalence was calculated using the first invasive tumor for each cancer site diagnosed during the previous 24 years. Maximum limited-duration prevalence is 24 years for 1992-2016 data.

^c Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^d Cases diagnosed more than 24 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000).

^e Complete prevalence is obtained by summing 0 to <24 and >=24.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 24.22

Cancer of the Stomach (Invasive)

Percent Distribution and Counts by Histology among Histologically Confirmed Cases, 2012-2016
Both Sexes by Race

Histology ^a	All Races		White		Black		Asian/Pacific Islander		American Indian/ Alaska Native ^b		Hispanic ^c	
	Count	Percent	Count	Percent	Count	Percent	Count	Percent	Count	Percent	Count	Percent
Carcinoma	42,127	91.1%	29,986	92.1%	5,701	85.2%	5,657	92.2%	267	96.4%	7,751	94.0%
Epidermoid carcinoma ^d (8051-8131)	383	0.8%	268	0.8%	74	1.1%	36	0.6%	-	-	46	0.6%
Squamous cell carcinoma (8070-8078,8083-8084)	362	0.8%	253	0.8%	71	1.1%	33	0.5%	-	-	40	0.5%
Adenocarcinoma (8050,8140-8147, 8160-8162,8180-8221, 8250-8507,8514,8520-8551, 8560,8570-8574, 8576,8940-8941)	37,148	80.4%	26,311	80.8%	4,950	74.0%	5,283	86.1%	238	85.9%	6,764	82.0%
Adenocarcinoma, NOS (8140)	21,351	46.2%	15,809	48.6%	2,727	40.8%	2,489	40.6%	131	47.3%	3,234	39.2%
Adenocarcinoma, intestinal type (8144)	3,820	8.3%	2,208	6.8%	665	9.9%	883	14.4%	29	10.5%	729	8.8%
Carcinoma, diffuse type (8145)	1,729	3.7%	1,073	3.3%	198	3.0%	426	6.9%	-	-	462	5.6%
Linitus plastica (8142)	194	0.4%	147	0.5%	19	0.3%	25	0.4%	-	-	45	0.5%
Mucinous adenocarcinoma (8470-8473,8480,8482)	535	1.2%	370	1.1%	89	1.3%	65	1.1%	-	-	93	1.1%
Mucin-producing adenocarcinoma (8481)	124	0.3%	88	0.3%	23	0.3%	-	-	-	-	26	0.3%
Signet ring adenocarcinoma (8490)	7,410	16.0%	5,262	16.2%	910	13.6%	1,098	17.9%	57	20.6%	1,800	21.8%
Other adenocarcinomas ^e	1,985	4.3%	1,354	4.2%	319	4.8%	284	4.6%	-	-	375	4.5%
Other specific carcinomas (8012-8015,8030-8046, 8150-8155,8170-8175, 8230-8249,8508,8510-8513, 8561-8562,8575,8580-8671)	3,732	8.1%	2,820	8.7%	520	7.8%	228	3.7%	24	8.7%	797	9.7%
Unspecified, Carcinoma, NOS (8010-8011,8020-8022)	864	1.9%	587	1.8%	157	2.3%	110	1.8%	-	-	144	1.7%

Source: SEER 21 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, Georgia excluding ATL/RG, Idaho, New York and Massachusetts).
Percents may not sum to 100 due to rounding.

^a Excludes Kaposi Sarcoma, mesothelioma, lymphomas, leukemias, myelomas, lymphoreticular, and immunoproliferative diseases.

^b Estimates for American Indian/Alaska Native are based on the Purchased/Referred Care Delivery Area (PRCDA) counties.

^c Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives.

^d Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

^e Epidermoid carcinoma includes squamous, basal, and transitional cell carcinomas.

^e Other adenocarcinomas includes histologies 8050, 8141, 8143, 8146-8147, 8160-8162, 8180-8221, 8250-8469, 8474-8479, 8483-8489, 8491-8507, 8514, 8520-8551, 8560, 8570-8574, 8576, 8940-8941.

- Statistic not shown due to fewer than 16 cases during the time period.

Table 24.22 - continued

Cancer of the Stomach (Invasive)

Percent Distribution and Counts by Histology among Histologically Confirmed Cases, 2012-2016
Both Sexes by Race

Histology ^a	All Races		White		Black		Asian/Pacific Islander		American Indian/ Alaska Native ^b		Hispanic ^c	
	Count	Percent	Count	Percent	Count	Percent	Count	Percent	Count	Percent	Count	Percent
Sarcoma and other soft tissue tumors (8680-8713,8800-8921, 8990-8991,9040-9044, 9120-9136,9150-9252, 9370-9373,9540-9582)	114	0.2%	94	0.3%	-	-	-	-	-	-	-	-
Other specific types (8720-8790,8930-8936, 8950-8983,9000-9030, 9060-9110,9260-9365, 9380-9539)	3,763	8.1%	2,324	7.1%	949	14.2%	433	7.1%	-	-	437	5.3%
Unspecified (8000-8005)	220	0.5%	149	0.5%	29	0.4%	35	0.6%	-	-	50	0.6%
Total ^d	46,224	100.0%	32,553	100.0%	6,688	100.0%	6,135	100.0%	277	100.0%	8,250	100.0%

Source: SEER 21 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, Georgia excluding ATL/RG, Idaho, New York and Massachusetts).

Percents may not sum to 100 due to rounding.

^a Excludes Kaposi Sarcoma, mesothelioma, lymphomas, leukemias, myelomas, lymphoreticular, and immunoproliferative diseases.

^b Estimates for American Indian/Alaska Native are based on the Purchased/Referred Care Delivery Area (PRCDA) counties.

^c Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives.


Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

^d Total includes carcinoma cases from the previous page.

- Statistic not shown due to fewer than 16 cases during the time period.

Figure 24.1

SEER Observed Incidence, SEER Delay Adjusted Incidence and US Death Rates^a Cancer of the Stomach, by Race and Sex



Source: SEER 9 areas and US Mortality Files (National Center for Health Statistics, CDC).

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).

Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.7, February 2019, National Cancer Institute.

The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.


* The APC is significantly different from zero ($p < 0.05$).

Figure 24.2

SEER Incidence and US Death Rates^a Cancer of the Stomach, Both Sexes

Joinpoint Analyses for Whites and Blacks from 1975-2016

and for Asian/Pacific Islanders, American Indians/Alaska Natives and Hispanics from 2000-2016



Source: Incidence data for whites and blacks are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta). Incidence data for Asian/Pacific Islanders, American Indians/Alaska Natives and Hispanics are from the SEER 21 areas (SEER 9 areas, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, Georgia excluding ATL/RG, Idaho, New York and Massachusetts).

Mortality data are from US Mortality Files, National Center for Health Statistics, CDC.

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).

Regression lines are calculated using the Joinpoint Regression Program Version 4.7, February 2019, National Cancer Institute. Joinpoint analyses for Whites and Blacks during the 1975-2016 period allow a maximum of 5 joinpoints. Analyses for other ethnic groups during the period 2000-2016 allow a maximum of 3 joinpoints.

^b API = Asian/Pacific Islander.

^c AI/AN = American Indian/Alaska Native. Rates for American Indian/Alaska Native are based on the Purchased/Referred Care Delivery Area (PRCDA) counties.


^d Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

Figure 24.3

Cancer of the Stomach

5-Year SEER Conditional Relative Survival and 95% Confidence Intervals

Probability of surviving the next 5 years given the cohort
has already survived 0, 1, or 3 years
2000-2015 by stage at diagnosis



Source: SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).
Percent surviving is not shown if based on less than 25 cases.
Confidence intervals are not shown if length of the confidence interval is greater than 5 times the standard error.