

Table 1.21
U.S. Prevalence Counts, Invasive Cancers Only, January 1, 2016^a
Using Different Tumor Inclusion Criteria^b

Site	Sex	5-Year Limited Duration			24-year Limited Duration	
		1st Invasive Tumor Ever ^c	1st Per Site in Previous 24 Years ^d	1st Per Site in Previous 5 Years ^e	1st Invasive Tumor Ever ^c	1st Per Site in Previous 24 Years ^d
All Sites	Both Sexes	4,618,309	4,788,249	5,268,178	13,238,684	13,725,579
	Male	2,265,828	2,321,794	2,561,190	6,465,129	6,615,379
	Female	2,352,481	2,466,455	2,706,988	6,773,555	7,110,200
Oral Cavity & Pharynx	Both Sexes	116,614	137,601	142,958	285,683	322,983
	Male	83,745	97,382	100,597	197,295	220,172
	Female	32,869	40,219	42,361	88,388	102,811
Esophagus	Both Sexes	21,742	27,114	27,273	37,573	44,872
	Male	16,966	21,041	21,144	29,074	34,504
	Female	4,776	6,073	6,129	8,499	10,368
Stomach	Both Sexes	44,232	53,399	54,255	89,342	104,126
	Male	25,922	31,498	31,907	51,076	59,722
	Female	18,310	21,901	22,348	38,266	44,404
Colon & Rectum	Both Sexes	384,576	444,429	452,381	1,077,905	1,211,974
	Male	200,293	230,124	234,263	550,903	613,912
	Female	184,283	214,305	218,118	527,002	598,062
Liver & Intrahepatic Bile Duct	Both Sexes	48,352	56,128	56,342	70,556	80,067
	Male	34,831	39,846	40,003	50,338	56,224
	Female	13,521	16,282	16,339	20,218	23,843
Pancreas	Both Sexes	43,231	53,929	54,044	57,753	70,735
	Male	21,568	27,124	27,209	28,410	35,088
	Female	21,663	26,805	26,835	29,343	35,647
Larynx	Both Sexes	29,186	34,757	35,246	74,772	85,315
	Male	23,886	28,382	28,773	61,289	69,580
	Female	5,300	6,375	6,473	13,483	15,735
Lung & Bronchus	Both Sexes	224,901	296,258	307,825	387,582	494,958
	Male	102,077	135,269	139,706	170,233	217,057
	Female	122,824	160,989	168,119	217,349	277,901
Melanoma of the Skin	Both Sexes	291,465	340,956	359,793	897,922	1,003,650
	Male	161,053	191,955	204,571	469,112	530,596
	Female	130,412	149,001	155,222	428,810	473,054
Breast	Female	915,040	1,009,922	1,069,620	2,862,940	3,112,731
Cervix	Female	40,564	43,027	43,191	172,170	181,314
Corpus & Uterus, NOS	Female	200,697	225,923	226,147	575,275	636,687
Ovary ^f	Female	62,449	71,655	71,737	158,048	180,026

^a U.S. 2016 cancer prevalence counts are based on 2016 cancer prevalence proportions from the SEER 13 registries (not including the Alaska Natives Registry) and 1/1/2016 U.S. population estimates based on the average of 2015 and 2016 population estimates from the U.S. Bureau of the Census.

^b Prevalence estimates are ambiguous for those with multiple cancers, unless the tumor inclusion criteria are understood. Depending on the application, different inclusion criteria may be appropriate. This table provides three different methods of tumor inclusion:

^c (c) First invasive tumor ever

^d (d) First invasive tumor for each cancer site diagnosed during the previous 24 years (1992-2015)

^e (e) First invasive tumor for each cancer site diagnosed during the previous 5 years (2011-2015)

For definitions (d) and (e) all sites is treated as a separate cancer "site".

Consider a woman who had three invasive cancers: Melanoma in 1981; Breast cancer in 2011; Melanoma in 2012.

In method (c) the melanoma is the woman's first cancer, and is counted for the melanoma and all sites 24-year limited duration prevalence. For 5-year limited duration prevalence, the woman is not counted at all since her first cancer occurred more than 5 years prior to 1/1/2016.

In method (d) the 1981 melanoma is counted for the melanoma and all sites 24-year limited duration prevalence. The 2011 breast cancer is counted for the breast 5-year and 24-year limited duration prevalence.

In method (e) the 2011 breast cancer is counted for the breast cancer and all sites 5-year limited duration prevalence. The 2012 melanoma is counted for 5-year limited duration prevalence for melanoma.

^f Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

Table 1.21 - continued
 U.S. Prevalence Counts, Invasive Cancers Only, January 1, 2016^a
 Using Different Tumor Inclusion Criteria^b

Site	Sex	5-Year Limited Duration			24-year Limited Duration	
		1st Invasive Tumor Ever ^c	1st Per Site in Previous 24 Years ^d	1st Per Site in Previous 5 Years ^e	1st Invasive Tumor Ever ^c	1st Per Site in Previous 24 Years ^d
Prostate	Male	852,616	925,846	925,901	2,883,784	3,069,398
Testis	Male	43,378	44,309	44,823	180,132	183,792
Urinary Bladder	Both Sexes	201,203	254,235	260,387	528,898	631,413
	Male	154,809	195,876	200,971	400,452	477,786
	Female	46,394	58,359	59,416	128,446	153,627
Kidney & Renal Pelvis	Both Sexes	168,243	204,952	208,257	406,205	479,439
	Male	105,668	129,796	132,082	248,043	294,719
	Female	62,575	75,156	76,175	158,162	184,720
Brain & Nervous System	Both Sexes	44,838	48,165	48,639	121,093	126,388
	Male	24,984	26,796	27,023	64,923	67,482
	Female	19,854	21,369	21,616	56,170	58,906
Thyroid	Both Sexes	203,126	229,536	230,264	587,959	644,571
	Male	46,377	55,079	55,254	127,419	144,092
	Female	156,749	174,457	175,010	460,540	500,479
Hodgkin Lymphoma	Both Sexes	35,058	37,437	37,487	140,786	146,789
	Male	19,459	20,890	20,914	74,193	77,233
	Female	15,599	16,547	16,573	66,593	69,556
Non-Hodgkin Lymphoma	Both Sexes	209,115	248,946	254,357	566,981	644,761
	Male	113,015	135,137	138,113	300,944	342,069
	Female	96,100	113,809	116,244	266,037	302,692
Myeloma	Both Sexes	66,177	78,454	78,949	111,277	128,969
	Male	36,691	44,371	44,700	61,269	72,052
	Female	29,486	34,083	34,249	50,008	56,917
Leukemia	Both Sexes	130,295	153,253	153,837	336,170	376,508
	Male	76,451	90,192	90,566	193,982	217,145
	Female	53,844	63,061	63,271	142,188	159,363
Acute Lymphocytic Leukemia	Both Sexes	19,515	20,277	20,277	69,109	70,308
	Male	10,797	11,203	11,203	38,607	39,179
	Female	8,718	9,074	9,074	30,502	31,129
Childhood (Ages 0-19)	Both Sexes	67,096	67,129	67,682	264,287	264,940
	Male	35,300	35,299	35,549	139,152	139,459
	Female	31,796	31,830	32,133	125,135	125,481
Kaposi Sarcoma	Both Sexes	7,567	8,281	8,281	32,104	33,825
	Male	6,918	7,549	7,549	30,486	32,021
	Female	649	732	732	1,618	1,804
Mesothelioma	Both Sexes	3,199	4,257	4,257	4,536	5,767
	Male	2,203	2,983	2,983	2,834	3,687
	Female	996	1,274	1,274	1,702	2,080

^a U.S. 2016 cancer prevalence counts are based on 2016 cancer prevalence proportions from the SEER 9 registries and 1/1/2016 U.S. population estimates based on the average of 2015 and 2016 population estimates from the U.S. Bureau of the Census.

^b Prevalence estimates are ambiguous for those with multiple cancers, unless the tumor inclusion criteria are understood. Depending on the application, different inclusion criteria may be appropriate. This table provides three different methods of tumor inclusion:

^c (c) First invasive tumor ever
^d (d) First invasive tumor for each cancer site diagnosed during the previous 24 years (1992-2015)
^e (e) First invasive tumor for each cancer site diagnosed during the previous 5 years (2011-2015)
 For definitions (d) and (e) all sites is treated as a separate cancer "site".

Consider a woman who had three invasive cancers: Melanoma in 1981; Breast cancer in 2011; Melanoma in 2012.

In method (c) the melanoma is the woman's first cancer, and is counted for the melanoma and all sites 24-year limited duration prevalence. For 5-year limited duration prevalence, the woman is not counted at all since her first cancer occurred more than 5 years prior to 1/1/2016. In method (d) the 1981 melanoma is counted for the melanoma and all sites 24-year limited duration prevalence. The 2011 breast cancer is counted for the breast 5-year and 24-year limited duration prevalence. In method (e) the 2011 breast cancer is counted for the breast cancer and all sites 5-year limited duration prevalence. The 2012 melanoma is counted for 5-year limited duration prevalence for melanoma.

Table 1.22
U.S. Complete Prevalence Counts, Invasive Cancers Only, January 1, 2016^a
By Age at Prevalence

Site/Sex	Age at Prevalence								
	All Ages ^c	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70+
All Sites									
Males	7,046,820	19,013	45,658	92,312	168,663	324,888	900,095	1,901,064	3,595,127
Females	8,292,168	17,282	38,768	93,009	252,052	631,369	1,452,263	2,179,010	3,628,415
Oral Cavity & Pharynx									
Males	250,059	79	483	1,435	3,554	12,626	49,283	84,123	98,476
Females	120,250	80	486	1,591	3,870	8,272	20,688	31,925	53,339
Esophagus									
Males	35,559	0	0	29	183	903	4,460	11,824	18,160
Females	10,918	0	0	11	54	242	1,320	3,146	6,144
Stomach									
Males	64,837	0	24	160	789	2,727	9,103	17,395	34,640
Females	48,217	0	34	215	859	2,676	7,066	11,599	25,768
Colon & Rectum									
Males	661,406	19	310	1,521	6,867	26,306	95,856	171,638	358,889
Females	663,516	39	358	1,832	6,831	25,137	85,953	146,080	397,285
Liver & Intrahep									
Males	57,099	624	769	529	434	1,690	10,569	26,699	15,787
Females	25,982	541	654	392	562	1,249	3,991	9,069	9,525
Pancreas									
Males	36,418	8	58	120	525	1,740	5,902	11,974	16,091
Females	37,136	12	70	338	716	2,183	6,294	10,710	16,812
Larynx									
Males	78,929	6	8	102	246	1,565	8,956	22,799	45,246
Females	17,422	0	16	16	151	658	3,099	5,295	8,188
Lung & Bronchus									
Males	239,856	51	127	370	1,223	4,483	23,185	66,309	144,108
Females	298,387	42	71	411	1,456	5,877	32,254	76,338	181,938
Melanoma of the Skin									
Males	611,158	58	530	4,277	16,027	39,516	98,628	166,460	285,663
Females	584,451	109	657	8,033	31,954	66,800	124,763	146,768	205,367

^a U.S. 2016 cancer prevalence counts are based on 2016 cancer prevalence proportions from the SEER 13 areas not including the Alaska Natives Registry (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, Rural Georgia, San Jose-Monterey, and Los Angeles) and 1/1/2016 U.S. population estimates based on the average of 2015 and 2016 population estimates from the U.S. Bureau of the Census.

^b Prevalence was calculated using the first invasive tumor for each cancer site diagnosed during the previous 24 years. Cases diagnosed more than 24 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000).

^c Due to rounding, the sum of the age specific estimates may not equal the all ages estimate.

Table 1.22 - continued
 U.S. Complete Prevalence Counts, Invasive Cancers Only, January 1, 2016^a
 By Age at Prevalence

Site/Sex	Age at Prevalence								
	All Ages ^c	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70+
Breast									
Males	19,496	0	0	4	135	584	2,200	4,872	11,703
Females	3,477,866	16	25	3,276	39,571	207,774	602,250	975,728	1,649,227
Cervix									
Females	289,696	0	42	1,803	14,401	41,426	66,591	72,523	92,910
Corpus & Uterus, NOS									
Females	772,245	8	14	867	7,585	28,997	101,642	228,398	404,735
Ovary ^d									
Females	229,875	106	1,124	4,479	8,509	19,167	46,532	64,680	85,279
Prostate									
Males	3,110,403	24	52	54	206	14,018	217,572	865,480	2,012,997
Urinary Bladder									
Males	525,713	60	99	541	2,145	8,784	39,848	120,226	354,010
Females	173,737	49	64	251	979	3,027	13,312	36,306	119,749
Kidney & Renal Pelvis									
Males	326,154	1,538	2,140	2,609	9,271	21,911	54,924	92,324	141,438
Females	207,050	1,706	2,575	3,132	6,768	14,900	33,535	52,972	91,463
Hodgkin Lymphoma									
Males	108,576	179	2,567	9,822	15,519	21,451	26,921	19,909	12,208
Females	102,397	70	1,843	8,849	16,460	21,101	24,748	17,379	11,947
Non-Hodgkin Lymphoma									
Males	367,558	900	4,029	8,311	13,996	27,628	61,422	94,982	156,290
Females	327,146	521	1,769	4,696	10,164	21,105	48,512	82,379	157,999
Myeloma									
Males	72,999	0	8	83	680	3,390	11,695	22,552	34,593
Females	58,392	3	9	60	426	2,610	8,912	17,672	28,699
Leukemia									
Males	235,556	6,996	15,050	16,790	12,272	17,182	30,974	51,848	84,446
Females	179,217	6,251	11,682	13,694	10,821	13,891	22,189	34,672	66,018
Acute Lymphocytic Leuk									
Males	52,176	5,892	12,608	13,226	8,345	6,762	3,097	1,536	710
Females	43,589	5,242	9,971	10,290	7,475	5,610	2,594	1,532	874

^a U.S. 2016 cancer prevalence counts are based on 2016 cancer prevalence proportions from the SEER 13 areas not including the Alaska Natives Registry (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, Rural Georgia, San Jose-Monterey, and Los Angeles) and 1/1/2016 U.S. population estimates based on the average of 2015 and 2016 population estimates from the U.S. Bureau of the Census.

^b Prevalence was calculated using the first invasive tumor for each cancer site diagnosed during the previous 24 years. Cases diagnosed more than 24 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000).

^c Due to rounding, the sum of the age specific estimates may not equal the all ages estimate.

Table 2.21
All Cancer Sites (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2016
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <24	0 to <24 ^b	>=24 ^d	Complete ^e
<u>Race</u>	<u>Sex</u>								
All Races	Both Sexes	4,788,249	3,608,607	2,608,324	1,793,597	926,801	13,725,578	1,613,409	15,338,988
	Males	2,321,794	1,829,893	1,272,515	808,408	382,770	6,615,379	431,441	7,046,820
	Females	2,466,455	1,778,715	1,335,809	985,189	544,031	7,110,200	1,181,968	8,292,168
White	Both Sexes	3,968,466	3,053,020	2,255,275	1,568,276	818,502	11,663,539	1,415,076	13,324,536
	Males	1,916,728	1,536,815	1,094,048	703,143	338,020	5,588,755	397,269	6,143,984
	Females	2,051,738	1,516,204	1,161,227	865,133	480,481	6,074,784	1,017,807	7,180,552
Black	Both Sexes	511,191	353,620	232,335	146,476	69,448	1,313,069	113,724	1,426,793
	Males	253,883	191,846	122,833	72,371	30,400	671,332	22,538	693,871
	Females	257,308	161,774	109,502	74,105	39,048	641,737	91,186	732,922
Asian/ Pacific Islander	Both Sexes	175,153	114,975	76,704	47,271	22,714	436,817	+	+
	Males	70,211	46,774	29,571	16,332	7,072	169,960	+	+
	Females	104,942	68,200	47,133	30,939	15,642	266,857	+	+
Hispanic	Both Sexes	389,479	259,791	174,874	108,086	53,984	986,213	+	+
	Males	171,164	120,631	79,132	44,942	20,093	435,962	+	+
	Females	218,314	139,160	95,742	63,144	33,891	550,250	+	+

Estimated prevalence percent^a on January 1, 2016, of the SEER population diagnosed in the previous 24 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)										Age-Adjusted ^c
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races	Both Sexes	4.2597	0.0899	0.2023	0.3922	0.8989	2.1532	4.9326	10.4420	18.3168	21.5939	3.6659
	Males	4.1699	0.0922	0.2141	0.3839	0.7190	1.4723	3.9242	10.6514	22.0759	29.4584	3.9311
	Females	4.3467	0.0875	0.1899	0.4009	1.0798	2.8229	5.8958	10.2522	15.1851	16.7558	3.5353
White	Both Sexes	4.6254	0.0951	0.2182	0.4325	0.9779	2.2654	5.0324	10.5285	18.5682	21.9709	3.7456
	Males	4.4728	0.0966	0.2321	0.4256	0.7883	1.5534	3.9221	10.4574	21.8930	29.3620	3.9319
	Females	4.7752	0.0935	0.2036	0.4397	1.1737	2.9850	6.1157	10.5944	15.7385	17.3436	3.6809
Black	Both Sexes	2.9117	0.0668	0.1406	0.2434	0.5916	1.6512	4.4428	10.3411	17.2760	18.1672	3.2840
	Males	3.1033	0.0708	0.1416	0.2341	0.4508	1.1414	4.1261	12.7006	24.6323	29.6595	4.1495
	Females	2.7350	0.0627	0.1395	0.2528	0.7206	2.1031	4.7187	8.4565	12.1550	12.4940	2.7687
Asian/ Pacific Islander	Both Sexes	2.1454	0.0723	0.1485	0.2388	0.5246	1.4613	3.3932	6.5148	10.4674	13.3000	2.2837
	Males	1.7424	0.0765	0.1606	0.2210	0.3500	0.7578	2.0343	5.4264	11.4323	17.2834	2.1151
	Females	2.5161	0.0679	0.1362	0.2567	0.6822	2.0847	4.5689	7.3937	9.6921	10.7354	2.4628
Hispanic	Both Sexes	1.7268	0.0837	0.1898	0.3320	0.6683	1.5085	3.4100	7.1864	12.5767	14.9162	2.5729
	Males	1.5106	0.0874	0.2050	0.3350	0.5272	0.9328	2.3723	6.7440	15.2423	21.1008	2.6949
	Females	1.9476	0.0798	0.1739	0.3288	0.8210	2.1006	4.4323	7.5777	10.5338	11.0982	2.5702

^a US 2016 cancer prevalence counts are based on 2016 cancer prevalence proportions from the SEER 13 Areas (excluding the Alaska Native Registry) and 1/1/2016 US population estimates based on the average of 2015 and 2016 population estimates from the US Bureau of the Census.

^b Prevalence was calculated using the first invasive tumor for each cancer site diagnosed during the previous 24 years. Maximum limited-duration prevalence is 24 years for 1992-2016 data.

^c Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^d Cases diagnosed more than 24 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000).

^e Complete prevalence is obtained by summing 0 to <24 and >=24.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 3.21
Brain and Other Nervous System (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2016
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <24	0 to <24 ^b	>=24 ^d	Complete ^e
<u>Race</u>	<u>Sex</u>								
All Races	Both Sexes	48,165	27,446	22,184	17,297	11,295	126,388	39,426	165,813
	Males	26,796	14,537	11,213	8,983	5,952	67,482	18,482	85,964
	Females	21,369	12,909	10,971	8,314	5,343	58,906	20,944	79,850
White	Both Sexes	41,387	23,514	19,265	15,074	9,729	108,971	38,377	149,087
	Males	23,187	12,576	9,881	7,889	5,085	58,618	19,472	79,085
	Females	18,200	10,939	9,385	7,185	4,645	50,353	18,905	70,002
Black	Both Sexes	3,998	2,399	1,832	1,454	983	10,666	4,539	15,205
	Males	2,150	1,114	800	707	545	5,316	1,807	7,123
	Females	1,848	1,285	1,032	747	438	5,350	2,732	8,082
Asian/ Pacific Islander	Both Sexes	1,727	968	754	481	295	4,224	+	+
	Males	882	512	349	212	158	2,112	+	+
	Females	845	456	405	268	137	2,111	+	+
Hispanic	Both Sexes	5,549	3,632	2,739	2,095	1,058	15,073	+	+
	Males	2,920	1,950	1,472	1,086	495	7,922	+	+
	Females	2,629	1,682	1,267	1,009	563	7,151	+	+

Estimated prevalence percent^a on January 1, 2016, of the SEER population diagnosed in the previous 24 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)										Age-Adjusted ^c
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races	Both Sexes	0.0392	0.0160	0.0350	0.0415	0.0410	0.0421	0.0509	0.0497	0.0440	0.0262	0.0382
	Males	0.0425	0.0160	0.0365	0.0440	0.0436	0.0458	0.0557	0.0566	0.0518	0.0316	0.0414
	Females	0.0360	0.0160	0.0334	0.0388	0.0385	0.0384	0.0462	0.0434	0.0374	0.0228	0.0352
White	Both Sexes	0.0432	0.0170	0.0383	0.0457	0.0454	0.0468	0.0566	0.0536	0.0471	0.0272	0.0419
	Males	0.0469	0.0168	0.0402	0.0488	0.0476	0.0507	0.0620	0.0604	0.0556	0.0338	0.0453
	Females	0.0396	0.0172	0.0363	0.0425	0.0432	0.0429	0.0513	0.0472	0.0399	0.0231	0.0387
Black	Both Sexes	0.0237	0.0122	0.0229	0.0289	0.0277	0.0248	0.0226	0.0288	0.0247	0.0208	0.0236
	Males	0.0246	0.0131	0.0224	0.0299	0.0287	0.0267	0.0218	0.0361	0.0261	0.0133	0.0246
	Females	0.0228	0.0113	0.0235	0.0280	0.0268	0.0231	0.0233	0.0229	0.0237	0.0245	0.0227
Asian/ Pacific Islander	Both Sexes	0.0207	0.0101	0.0227	0.0212	0.0197	0.0214	0.0274	0.0246	0.0237	0.0138	0.0205
	Males	0.0217	0.0106	0.0242	0.0200	0.0228	0.0217	0.0290	0.0261	0.0264	0.0114	0.0215
	Females	0.0199	0.0095	0.0211	0.0225	0.0170	0.0210	0.0259	0.0234	0.0214	0.0154	0.0196
Hispanic	Both Sexes	0.0264	0.0123	0.0251	0.0265	0.0253	0.0296	0.0417	0.0388	0.0392	0.0235	0.0278
	Males	0.0275	0.0118	0.0262	0.0273	0.0246	0.0329	0.0465	0.0426	0.0371	0.0263	0.0292
	Females	0.0253	0.0128	0.0240	0.0256	0.0262	0.0262	0.0370	0.0354	0.0408	0.0219	0.0264

^a US 2016 cancer prevalence counts are based on 2016 cancer prevalence proportions from the SEER 13 Areas (excluding the Alaska Native Registry) and 1/1/2016 US population estimates based on the average of 2015 and 2016 population estimates from the US Bureau of the Census.

^b Prevalence was calculated using the first invasive tumor for each cancer site diagnosed during the previous 24 years. Maximum limited-duration prevalence is 24 years for 1992-2016 data.

^c Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^d Cases diagnosed more than 24 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000).

^e Complete prevalence is obtained by summing 0 to <24 and >=24.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 4.24
Cancer of the Breast (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2016
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <24	0 to <24 ^b	>=24 ^d	Complete ^e
<u>Race</u>	<u>Sex</u>								
All Races	Both Sexes	1,017,961	781,533	614,055	469,555	248,033	3,131,136	366,226	3,497,363
	Males	8,040	4,766	2,801	1,984	814	18,405	1,091	19,496
	Females	1,009,922	776,767	611,253	467,571	247,219	3,112,731	365,135	3,477,866
White	Both Sexes	843,120	663,539	534,432	413,399	219,892	2,674,382	328,144	3,008,268
	Males	6,591	4,062	2,367	1,711	673	15,403	+	+
	Females	836,530	659,478	532,065	411,688	219,219	2,658,979	328,144	3,008,268
Black	Both Sexes	112,746	75,971	51,618	36,535	18,893	295,762	30,695	324,210
	Males	1,148	516	343	179	60	2,246	+	+
	Females	111,597	75,455	51,275	36,356	18,832	293,515	30,695	324,210
Asian/ Pacific Islander	Both Sexes	46,680	31,992	22,348	15,092	6,872	122,984	+	+
	Males	223	112	53	38	24	451	+	+
	Females	46,457	31,879	22,295	15,054	6,847	122,533	+	+
Hispanic	Both Sexes	82,496	54,432	38,030	25,114	12,324	212,396	+	+
	Males	440	177	110	70	22	820	+	+
	Females	82,057	54,255	37,920	25,044	12,301	211,577	+	+

Estimated prevalence percent^a on January 1, 2016, of the SEER population diagnosed in the previous 24 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)										Age-Adjusted ^c
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races	Both Sexes	0.9717	0.0000	0.0001	0.0073	0.0938	0.5093	1.3510	2.5533	4.0039	4.7517	0.8206
	Males	0.0116	-	-	-	0.0006	0.0029	0.0102	0.0279	0.0677	0.1002	0.0110
	Females	1.9029	0.0001	0.0001	0.0148	0.1875	1.0073	2.6317	4.8408	7.2831	7.6133	1.5075
White	Both Sexes	1.0606	0.0001	0.0001	0.0068	0.0913	0.5103	1.3667	2.6114	4.1338	4.9105	0.8383
	Males	0.0123	-	-	-	0.0005	0.0024	0.0088	0.0275	0.0679	0.1018	0.0109
	Females	2.0901	0.0001	0.0001	0.0140	0.1851	1.0236	2.6917	5.0073	7.5942	7.9210	1.5552
Black	Both Sexes	0.6558	-	0.0001	0.0089	0.1040	0.4942	1.2574	2.2766	3.4061	3.9065	0.7294
	Males	0.0104	-	-	-	0.0008	0.0064	0.0190	0.0368	0.0798	0.0963	0.0136
	Females	1.2509	-	0.0003	0.0178	0.1987	0.9266	2.3364	4.0656	5.7216	5.7874	1.2600
Asian/ Pacific Islander	Both Sexes	0.6040	-	-	0.0073	0.0908	0.4933	1.2399	2.0447	2.6285	2.7875	0.6253
	Males	0.0046	-	-	-	0.0012	0.0004	0.0064	0.0151	0.0327	0.0490	0.0057
	Females	1.1553	-	-	0.0144	0.1716	0.9302	2.3071	3.6837	4.7140	4.5507	1.1254
Hispanic	Both Sexes	0.3719	-	0.0001	0.0058	0.0715	0.3599	0.9502	1.7962	2.6322	2.8541	0.5559
	Males	0.0028	-	-	-	0.0005	0.0020	0.0042	0.0136	0.0317	0.0534	0.0055
	Females	0.7489	-	0.0001	0.0121	0.1483	0.7280	1.8823	3.3728	4.6253	4.5832	1.0193

^a US 2016 cancer prevalence counts are based on 2016 cancer prevalence proportions from the SEER 13 Areas (excluding the Alaska Native Registry) and 1/1/2016 US population estimates based on the average of 2015 and 2016 population estimates from the US Bureau of the Census.

^b Prevalence was calculated using the first invasive tumor for each cancer site diagnosed during the previous 24 years. Maximum limited-duration prevalence is 24 years for 1992-2016 data.

^c Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^d Cases diagnosed more than 24 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000).

^e Complete prevalence is obtained by summing 0 to <24 and >=24.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 5.15
Cancer of the Cervix Uteri (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2016
By Race/Ethnicity and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <24	0 to <24 ^b	>=24 ^d	Complete ^e
<u>Race</u>								
All Races	43,027	37,307	36,007	37,748	27,225	181,314	108,382	289,696
White	34,247	30,268	29,636	31,086	22,910	148,148	94,604	245,766
Black	5,091	4,207	4,054	4,187	2,664	20,203	14,589	34,792
Asian/Pacific Islander	2,373	1,863	1,692	1,572	1,055	8,555	+	+
Hispanic	7,664	7,108	7,164	6,466	4,369	32,770	+	+

Estimated prevalence percent^a on January 1, 2016, of the SEER population diagnosed in the previous 24 years
By Age at Prevalence and Race/Ethnicity

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^c
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>											
All Races	0.1108	-	0.0002	0.0082	0.0682	0.1963	0.2416	0.2191	0.1835	0.1242	0.0995
White	0.1165	-	0.0002	0.0087	0.0746	0.2127	0.2504	0.2153	0.1741	0.1131	0.1029
Black	0.0861	-	0.0005	0.0064	0.0491	0.1385	0.1990	0.2215	0.2170	0.1734	0.0865
Asian/Pacific Islander	0.0807	-	-	0.0046	0.0282	0.1021	0.1768	0.2253	0.2316	0.2110	0.0774
Hispanic	0.1160	-	0.0002	0.0086	0.0722	0.2158	0.3486	0.3843	0.3754	0.2867	0.1440

^a US 2016 cancer prevalence counts are based on 2016 cancer prevalence proportions from the SEER 13 Areas (excluding the Alaska Native Registry) and 1/1/2016 US population estimates based on the average of 2015 and 2016 population estimates from the US Bureau of the Census.

^b Prevalence was calculated using the first invasive tumor for each cancer site diagnosed during the previous 24 years. Maximum limited-duration prevalence is 24 years for 1992-2016 data.

^c Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^d Cases diagnosed more than 24 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000).

^e Complete prevalence is obtained by summing 0 to <24 and >=24.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 6.28
Cancer of the Colon and Rectum (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2016
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <24	0 to <24 ^b	>=24 ^d	Complete ^e
<u>Race</u>	<u>Sex</u>								
All Races	Both Sexes	444,429	317,632	229,064	150,394	70,455	1,211,974	112,948	1,324,922
	Males	230,124	160,170	114,529	75,117	33,972	613,912	47,494	661,406
	Females	214,305	157,462	114,534	75,278	36,483	598,062	65,454	663,516
White	Both Sexes	360,394	260,557	192,176	128,096	59,919	1,001,141	97,331	1,113,103
	Males	187,381	132,866	97,733	64,653	29,146	511,778	41,357	561,266
	Females	173,013	127,691	94,443	63,443	30,773	489,363	55,974	551,837
Black	Both Sexes	53,043	36,471	23,891	14,824	7,043	135,272	10,020	145,292
	Males	26,142	16,561	10,260	6,622	3,106	62,692	4,004	66,696
	Females	26,901	19,910	13,631	8,202	3,937	72,580	6,016	78,596
Asian/ Pacific Islander	Both Sexes	21,293	14,221	9,467	5,573	2,655	53,209	+	+
	Males	11,306	7,247	4,636	2,774	1,269	27,231	+	+
	Females	9,988	6,974	4,831	2,799	1,387	25,978	+	+
Hispanic	Both Sexes	38,986	23,945	14,554	8,336	3,491	89,312	+	+
	Males	20,435	12,487	7,373	4,349	1,668	46,312	+	+
	Females	18,551	11,457	7,181	3,987	1,823	43,000	+	+

Estimated prevalence percent^a on January 1, 2016, of the SEER population diagnosed in the previous 24 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted ^c	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races	Both Sexes	0.3761	0.0001	0.0016	0.0074	0.0323	0.1242	0.3988	0.8329	1.6388	2.8341	0.3204
	Males	0.3870	0.0001	0.0015	0.0066	0.0323	0.1284	0.4329	0.9550	1.9166	3.2563	0.3656
	Females	0.3656	0.0002	0.0018	0.0083	0.0323	0.1200	0.3663	0.7223	1.4074	2.5744	0.2849
White	Both Sexes	0.3970	0.0002	0.0018	0.0080	0.0334	0.1237	0.3838	0.7948	1.6121	2.8527	0.3148
	Males	0.4096	0.0001	0.0017	0.0067	0.0333	0.1282	0.4173	0.9164	1.8930	3.2560	0.3595
	Females	0.3847	0.0002	0.0019	0.0093	0.0335	0.1191	0.3512	0.6820	1.3730	2.6003	0.2789
Black	Both Sexes	0.3000	-	0.0005	0.0043	0.0288	0.1255	0.4444	1.0353	1.8322	2.7160	0.3491
	Males	0.2898	-	0.0003	0.0042	0.0310	0.1235	0.4636	1.1528	2.0460	3.2824	0.3925
	Females	0.3093	-	0.0008	0.0045	0.0268	0.1272	0.4276	0.9414	1.6833	2.4364	0.3213
Asian/ Pacific Islander	Both Sexes	0.2613	-	0.0008	0.0051	0.0218	0.1050	0.3875	0.8164	1.4531	2.3489	0.2870
	Males	0.2792	-	0.0011	0.0051	0.0205	0.1118	0.4444	0.9562	1.7523	2.7523	0.3363
	Females	0.2449	-	0.0006	0.0051	0.0230	0.0991	0.3383	0.7036	1.2127	2.0892	0.2488
Hispanic	Both Sexes	0.1564	0.0002	0.0010	0.0050	0.0242	0.0879	0.3194	0.7058	1.3264	2.0502	0.2530
	Males	0.1605	0.0001	0.0007	0.0043	0.0246	0.0907	0.3398	0.7983	1.6371	2.5467	0.2989
	Females	0.1522	0.0003	0.0013	0.0058	0.0239	0.0851	0.2994	0.6241	1.0883	1.7437	0.2190

^a US 2016 cancer prevalence counts are based on 2016 cancer prevalence proportions from the SEER 13 Areas (excluding the Alaska Native Registry) and 1/1/2016 US population estimates based on the average of 2015 and 2016 population estimates from the US Bureau of the Census.

^b Prevalence was calculated using the first invasive tumor for each cancer site diagnosed during the previous 24 years. Maximum limited-duration prevalence is 24 years for 1992-2016 data.

^c Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^d Cases diagnosed more than 24 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000).

^e Complete prevalence is obtained by summing 0 to <24 and >=24.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 7.15
Cancer of the Corpus and Uterus, NOS (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2016
By Race/Ethnicity and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <24	0 to <24 ^b	>=24 ^d	Complete ^e
<u>Race</u>								
All Races	225,923	161,291	116,320	86,194	46,958	636,687	135,558	772,245
White	190,505	140,678	103,752	78,371	43,211	556,518	115,722	677,802
Black	21,411	11,777	7,015	4,029	1,919	46,152	5,451	51,603
Asian/Pacific Islander	9,979	6,356	4,242	2,678	1,383	24,638	+	+
Hispanic	20,762	11,784	8,068	4,746	2,486	47,847	+	+

Estimated prevalence percent^a on January 1, 2016, of the SEER population diagnosed in the previous 24 years
By Age at Prevalence and Race/Ethnicity

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^c
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>											
All Races	0.3892	0.0000	0.0001	0.0039	0.0359	0.1396	0.4412	1.1392	1.6248	1.4673	0.3031
White	0.4375	-	0.0001	0.0034	0.0362	0.1434	0.4620	1.2017	1.7198	1.5662	0.3190
Black	0.1967	0.0002	-	0.0052	0.0270	0.0948	0.2599	0.7780	1.1136	0.8228	0.1986
Asian/Pacific Islander	0.2323	-	-	0.0055	0.0360	0.1543	0.4831	0.8547	0.8978	0.6922	0.2210
Hispanic	0.1694	-	-	0.0056	0.0520	0.1565	0.3945	0.8469	1.0561	0.7990	0.2280

^a US 2016 cancer prevalence counts are based on 2016 cancer prevalence proportions from the SEER 13 Areas (excluding the Alaska Native Registry) and 1/1/2016 US population estimates based on the average of 2015 and 2016 population estimates from the US Bureau of the Census.

^b Prevalence was calculated using the first invasive tumor for each cancer site diagnosed during the previous 24 years. Maximum limited-duration prevalence is 24 years for 1992-2016 data.

^c Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^d Cases diagnosed more than 24 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000).

^e Complete prevalence is obtained by summing 0 to <24 and >=24.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 8.21
Cancer of the Esophagus (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2016
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <24	0 to <24 ^b	>=24 ^d	Complete ^e
<u>Race</u>	<u>Sex</u>								
All Races	Both Sexes	27,114	9,592	4,960	2,379	827	44,872	1,605	46,477
	Males	21,041	7,281	3,783	1,779	619	34,504	1,055	35,559
	Females	6,073	2,311	1,176	600	208	10,368	550	10,918
White	Both Sexes	24,421	8,819	4,574	2,089	672	40,576	1,297	42,147
	Males	19,236	6,816	3,554	1,629	507	31,742	891	32,837
	Females	5,185	2,003	1,019	461	165	8,834	406	9,310
Black	Both Sexes	1,881	508	251	215	107	2,963	208	3,171
	Males	1,226	272	129	109	79	1,814	84	1,898
	Females	655	237	122	107	28	1,148	124	1,273
Asian/ Pacific Islander	Both Sexes	552	169	85	57	25	888	+	+
	Males	392	112	57	35	20	616	+	+
	Females	160	57	28	21	5	271	+	+
Hispanic	Both Sexes	1,472	387	201	78	64	2,203	+	+
	Males	1,158	322	175	67	45	1,767	+	+
	Females	314	66	26	11	19	436	+	+

Estimated prevalence percent^a on January 1, 2016, of the SEER population diagnosed in the previous 24 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted ^c	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races	Both Sexes	0.0139	-	-	0.0001	0.0006	0.0028	0.0129	0.0407	0.0726	0.0725	0.0116
	Males	0.0217	-	-	0.0001	0.0009	0.0044	0.0204	0.0678	0.1256	0.1324	0.0199
	Females	0.0063	-	-	0.0001	0.0003	0.0012	0.0057	0.0161	0.0284	0.0356	0.0049
White	Both Sexes	0.0161	-	-	0.0001	0.0006	0.0031	0.0138	0.0442	0.0783	0.0764	0.0125
	Males	0.0254	-	-	0.0001	0.0010	0.0049	0.0221	0.0737	0.1368	0.1402	0.0215
	Females	0.0069	-	-	0.0000	0.0002	0.0012	0.0057	0.0168	0.0286	0.0365	0.0050
Black	Both Sexes	0.0066	-	-	-	0.0006	0.0020	0.0100	0.0262	0.0439	0.0428	0.0074
	Males	0.0084	-	-	-	0.0006	0.0025	0.0144	0.0389	0.0585	0.0714	0.0108
	Females	0.0049	-	-	-	0.0005	0.0014	0.0061	0.0160	0.0337	0.0287	0.0050
Asian/ Pacific Islander	Both Sexes	0.0044	-	-	0.0003	-	0.0013	0.0061	0.0145	0.0276	0.0355	0.0048
	Males	0.0063	-	-	0.0004	-	0.0018	0.0087	0.0252	0.0413	0.0591	0.0076
	Females	0.0026	-	-	-	-	0.0008	0.0038	0.0059	0.0166	0.0203	0.0027
Hispanic	Both Sexes	0.0039	-	-	0.0001	0.0004	0.0022	0.0070	0.0193	0.0390	0.0398	0.0063
	Males	0.0061	-	-	0.0001	0.0007	0.0035	0.0124	0.0306	0.0757	0.0766	0.0114
	Females	0.0015	-	-	-	-	0.0009	0.0017	0.0092	0.0109	0.0170	0.0022

^a US 2016 cancer prevalence counts are based on 2016 cancer prevalence proportions from the SEER 13 Areas (excluding the Alaska Native Registry) and 1/1/2016 US population estimates based on the average of 2015 and 2016 population estimates from the US Bureau of the Census.

^b Prevalence was calculated using the first invasive tumor for each cancer site diagnosed during the previous 24 years. Maximum limited-duration prevalence is 24 years for 1992-2016 data.

^c Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^d Cases diagnosed more than 24 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000).

^e Complete prevalence is obtained by summing 0 to <24 and >=24.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 9.21
Hodgkin LymphomaEstimated United States Cancer Prevalence Counts^a on January 1, 2016
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <24	0 to <24 ^b	>=24 ^d	Complete ^e
<u>Race</u>	<u>Sex</u>								
All Races	Both Sexes	37,437	35,145	29,981	25,990	18,237	146,789	64,184	210,974
	Males	20,890	18,902	15,287	13,181	8,972	77,233	31,343	108,576
	Females	16,547	16,243	14,694	12,808	9,264	69,556	32,841	102,397
White	Both Sexes	30,507	29,258	25,439	22,613	16,108	123,925	61,817	187,446
	Males	17,050	15,906	13,104	11,426	7,981	65,468	31,491	97,930
	Females	13,457	13,352	12,334	11,187	8,126	58,457	30,325	89,516
Black	Both Sexes	4,976	4,443	3,427	2,643	1,745	17,234	5,362	22,597
	Males	2,773	2,284	1,554	1,334	831	8,777	3,481	12,258
	Females	2,203	2,159	1,873	1,310	913	8,458	1,881	10,339
Asian/ Pacific Islander	Both Sexes	1,082	927	720	414	229	3,372	+	+
	Males	592	445	373	227	77	1,714	+	+
	Females	491	482	347	187	151	1,658	+	+
Hispanic	Both Sexes	4,772	4,081	2,993	2,316	1,344	15,506	+	+
	Males	2,636	2,220	1,628	1,260	734	8,480	+	+
	Females	2,136	1,861	1,364	1,056	610	7,026	+	+

Estimated prevalence percent^a on January 1, 2016, of the SEER population diagnosed in the previous 24 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)										Age-Adjusted ^c
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races	Both Sexes	0.0456	0.0006	0.0106	0.0412	0.0729	0.0801	0.0637	0.0502	0.0477	0.0372	0.0454
	Males	0.0487	0.0009	0.0120	0.0424	0.0700	0.0826	0.0722	0.0605	0.0580	0.0453	0.0483
	Females	0.0425	0.0004	0.0090	0.0400	0.0758	0.0776	0.0556	0.0410	0.0390	0.0323	0.0428
White	Both Sexes	0.0491	0.0005	0.0109	0.0450	0.0796	0.0861	0.0672	0.0525	0.0503	0.0403	0.0487
	Males	0.0524	0.0007	0.0120	0.0464	0.0758	0.0884	0.0754	0.0626	0.0611	0.0487	0.0514
	Females	0.0460	0.0003	0.0098	0.0436	0.0835	0.0838	0.0592	0.0431	0.0411	0.0350	0.0462
Black	Both Sexes	0.0382	0.0015	0.0102	0.0317	0.0622	0.0740	0.0609	0.0478	0.0392	0.0179	0.0400
	Males	0.0406	0.0021	0.0127	0.0313	0.0608	0.0744	0.0725	0.0602	0.0475	0.0256	0.0432
	Females	0.0360	0.0008	0.0075	0.0322	0.0635	0.0736	0.0508	0.0379	0.0335	0.0141	0.0375
Asian/ Pacific Islander	Both Sexes	0.0166	0.0003	0.0070	0.0203	0.0282	0.0233	0.0178	0.0143	0.0150	0.0096	0.0158
	Males	0.0176	-	0.0098	0.0245	0.0264	0.0244	0.0182	0.0141	0.0197	0.0121	0.0169
	Females	0.0156	-	0.0042	0.0162	0.0299	0.0224	0.0175	0.0145	0.0112	0.0080	0.0147
Hispanic	Both Sexes	0.0271	0.0007	0.0096	0.0300	0.0440	0.0423	0.0396	0.0429	0.0493	0.0313	0.0302
	Males	0.0294	0.0009	0.0112	0.0324	0.0437	0.0449	0.0446	0.0528	0.0569	0.0472	0.0333
	Females	0.0249	0.0005	0.0080	0.0275	0.0442	0.0395	0.0347	0.0341	0.0434	0.0215	0.0274

^a US 2016 cancer prevalence counts are based on 2016 cancer prevalence proportions from the SEER 13 Areas (excluding the Alaska Native Registry) and 1/1/2016 US population estimates based on the average of 2015 and 2016 population estimates from the US Bureau of the Census.

^b Prevalence was calculated using the first invasive tumor for each cancer site diagnosed during the previous 24 years. Maximum limited-duration prevalence is 24 years for 1992-2016 data.

^c Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^d Cases diagnosed more than 24 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000).

^e Complete prevalence is obtained by summing 0 to <24 and >=24.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 10.15
Kaposi SarcomaEstimated United States Cancer Prevalence Counts^a on January 1, 2016
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <24	0 to <24 ^b	>=24 ^d	Complete ^e
<u>Race</u>	<u>Sex</u>								
All Races	Both Sexes	8,281	7,394	6,306	6,738	5,106	33,825	+	+
	Males	7,549	6,889	6,078	6,502	5,003	32,021	+	+
	Females	732	504	227	237	103	1,804	+	+
White	Both Sexes	5,019	5,231	4,815	5,124	4,321	24,509	+	+
	Males	4,529	4,824	4,651	4,956	4,227	23,187	+	+
	Females	489	406	163	168	95	1,321	+	+
Black	Both Sexes	2,472	1,627	1,163	1,108	480	6,851	+	+
	Males	2,325	1,570	1,120	1,065	472	6,552	+	+
	Females	147	58	43	43	9	300	+	+
Asian/ Pacific Islander	Both Sexes	171	127	89	70	42	499	+	+
	Males	163	122	86	70	42	482	+	+
	Females	8	5	3	0	0	16	+	+
Hispanic	Both Sexes	1,281	1,116	1,027	1,038	667	5,129	+	+
	Males	1,145	1,001	994	1,000	656	4,795	+	+
	Females	136	115	33	38	11	334	+	+

Estimated prevalence percent^a on January 1, 2016, of the SEER population diagnosed in the previous 24 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)										Age-Adjusted ^c
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races	Both Sexes	0.0105	0.0000	0.0000	0.0015	0.0070	0.0163	0.0273	0.0178	0.0132	0.0207	0.0096
	Males	0.0202	0.0001	0.0000	0.0029	0.0137	0.0322	0.0544	0.0361	0.0257	0.0382	0.0189
	Females	0.0011	-	0.0000	-	0.0003	0.0008	0.0013	0.0011	0.0029	0.0098	0.0009
White	Both Sexes	0.0097	0.0000	0.0000	0.0005	0.0043	0.0133	0.0259	0.0173	0.0132	0.0207	0.0084
	Males	0.0186	0.0001	-	0.0010	0.0083	0.0259	0.0513	0.0349	0.0255	0.0384	0.0164
	Females	0.0010	-	0.0001	-	0.0001	0.0006	0.0010	0.0010	0.0028	0.0097	0.0008
Black	Both Sexes	0.0152	0.0001	0.0001	0.0063	0.0221	0.0333	0.0366	0.0206	0.0103	0.0117	0.0159
	Males	0.0303	0.0003	0.0003	0.0126	0.0444	0.0685	0.0751	0.0444	0.0228	0.0221	0.0327
	Females	0.0013	-	-	-	0.0016	0.0021	0.0030	0.0016	0.0017	0.0066	0.0013
Asian/ Pacific Islander	Both Sexes	0.0024	-	-	0.0003	0.0018	0.0045	0.0054	0.0051	0.0039	0.0062	0.0024
	Males	0.0049	-	-	0.0006	0.0039	0.0096	0.0112	0.0108	0.0088	0.0136	0.0052
	Females	0.0002	-	-	-	-	-	0.0005	0.0006	-	-	0.0001
Hispanic	Both Sexes	0.0090	0.0001	-	0.0007	0.0076	0.0173	0.0316	0.0205	0.0193	0.0408	0.0114
	Males	0.0166	0.0001	-	0.0013	0.0146	0.0333	0.0617	0.0415	0.0365	0.0610	0.0216
	Females	0.0012	-	-	-	0.0001	0.0009	0.0020	0.0018	0.0061	0.0283	0.0018

^a US 2016 cancer prevalence counts are based on 2016 cancer prevalence proportions from the SEER 13 Areas (excluding the Alaska Native Registry) and 1/1/2016 US population estimates based on the average of 2015 and 2016 population estimates from the US Bureau of the Census.

^b Prevalence was calculated using the first invasive tumor for each cancer site diagnosed during the previous 24 years. Maximum limited-duration prevalence is 24 years for 1992-2016 data.

^c Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^d Cases diagnosed more than 24 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000).

^e Complete prevalence is obtained by summing 0 to <24 and >=24.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 11.21
Cancer of the Kidney And Renal Pelvis (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2016
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <24	0 to <24 ^b	>=24 ^d	Complete ^e
<u>Race</u>	<u>Sex</u>								
All Races	Both Sexes	204,952	136,846	77,236	40,634	19,773	479,440	53,764	533,204
	Males	129,796	83,708	46,315	23,572	11,329	294,719	31,435	326,154
	Females	75,156	53,138	30,921	17,061	8,444	184,720	22,329	207,050
White	Both Sexes	169,767	114,748	66,313	34,810	17,383	403,021	51,538	459,398
	Males	108,041	70,416	39,790	20,653	10,128	249,028	26,810	278,804
	Females	61,726	44,332	26,524	14,157	7,254	153,993	24,729	180,594
Black	Both Sexes	25,424	16,219	8,023	4,521	1,821	56,009	7,697	63,706
	Males	15,638	9,671	4,814	2,207	873	33,202	3,486	36,689
	Females	9,787	6,549	3,209	2,314	949	22,807	4,210	27,017
Asian/ Pacific Islander	Both Sexes	6,056	3,469	1,845	862	353	12,585	+	+
	Males	3,840	2,139	1,076	493	206	7,755	+	+
	Females	2,216	1,330	769	368	147	4,831	+	+
Hispanic	Both Sexes	21,695	12,760	6,623	3,172	1,511	45,761	+	+
	Males	12,589	7,171	3,716	1,741	793	26,010	+	+
	Females	9,106	5,589	2,907	1,431	718	19,751	+	+

Estimated prevalence percent^a on January 1, 2016, of the SEER population diagnosed in the previous 24 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)										Age-Adjusted ^c
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races	Both Sexes	0.1488	0.0080	0.0113	0.0103	0.0200	0.0717	0.1876	0.3863	0.6473	0.6571	0.1272
	Males	0.1858	0.0075	0.0100	0.0090	0.0228	0.0876	0.2406	0.5189	0.9073	0.9629	0.1706
	Females	0.1129	0.0086	0.0126	0.0116	0.0171	0.0560	0.1369	0.2662	0.4308	0.4690	0.0914
White	Both Sexes	0.1598	0.0077	0.0115	0.0106	0.0204	0.0734	0.1894	0.3861	0.6555	0.6676	0.1285
	Males	0.1993	0.0068	0.0108	0.0095	0.0228	0.0882	0.2404	0.5140	0.9153	0.9788	0.1713
	Females	0.1210	0.0086	0.0122	0.0119	0.0179	0.0584	0.1396	0.2675	0.4344	0.4728	0.0925
Black	Both Sexes	0.1242	0.0102	0.0134	0.0113	0.0209	0.0723	0.2029	0.4498	0.6969	0.6468	0.1374
	Males	0.1535	0.0111	0.0095	0.0090	0.0257	0.0929	0.2717	0.6324	1.0112	0.9844	0.1912
	Females	0.0972	0.0094	0.0174	0.0135	0.0165	0.0540	0.1430	0.3040	0.4781	0.4801	0.0990
Asian/ Pacific Islander	Both Sexes	0.0618	0.0049	0.0042	0.0028	0.0090	0.0374	0.0970	0.2006	0.3198	0.3893	0.0661
	Males	0.0795	0.0041	0.0029	0.0028	0.0110	0.0495	0.1334	0.2905	0.4475	0.5515	0.0916
	Females	0.0455	0.0057	0.0054	0.0028	0.0071	0.0266	0.0655	0.1280	0.2172	0.2848	0.0461
Hispanic	Both Sexes	0.0801	0.0059	0.0092	0.0072	0.0199	0.0686	0.1693	0.3697	0.6059	0.6287	0.1198
	Males	0.0901	0.0049	0.0085	0.0047	0.0208	0.0743	0.2016	0.4717	0.8191	0.8511	0.1519
	Females	0.0699	0.0070	0.0100	0.0099	0.0189	0.0628	0.1374	0.2794	0.4425	0.4913	0.0946

^a US 2016 cancer prevalence counts are based on 2016 cancer prevalence proportions from the SEER 13 Areas (excluding the Alaska Native Registry) and 1/1/2016 US population estimates based on the average of 2015 and 2016 population estimates from the US Bureau of the Census.

^b Prevalence was calculated using the first invasive tumor for each cancer site diagnosed during the previous 24 years. Maximum limited-duration prevalence is 24 years for 1992-2016 data.

^c Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^d Cases diagnosed more than 24 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000).

^e Complete prevalence is obtained by summing 0 to <24 and >=24.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 12.21
Cancer of the Larynx (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2016
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <24	0 to <24 ^b	>=24 ^d	Complete ^e
<u>Race</u>	<u>Sex</u>								
All Races	Both Sexes	34,757	22,057	13,805	9,470	5,227	85,315	11,036	96,351
	Males	28,382	18,187	11,166	7,518	4,327	69,580	9,349	78,929
	Females	6,375	3,870	2,639	1,952	900	15,735	1,687	17,422
White	Both Sexes	28,856	18,609	11,841	8,141	4,489	71,936	8,929	81,619
	Males	23,737	15,365	9,602	6,504	3,698	58,906	7,570	67,116
	Females	5,119	3,244	2,239	1,636	791	13,030	1,359	14,502
Black	Both Sexes	4,799	2,771	1,575	1,054	621	10,820	949	11,769
	Males	3,715	2,238	1,244	762	546	8,505	806	9,312
	Females	1,084	534	331	291	76	2,315	142	2,457
Asian/ Pacific Islander	Both Sexes	598	421	259	164	77	1,519	+	+
	Males	493	360	215	149	63	1,281	+	+
	Females	105	61	44	15	14	238	+	+
Hispanic	Both Sexes	2,272	1,417	779	584	260	5,311	+	+
	Males	1,973	1,250	661	436	231	4,551	+	+
	Females	298	167	118	148	29	760	+	+

Estimated prevalence percent^a on January 1, 2016, of the SEER population diagnosed in the previous 24 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)										Age-Adjusted ^c
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races	Both Sexes	0.0265	0.0000	0.0001	0.0003	0.0009	0.0054	0.0271	0.0760	0.1368	0.1339	0.0221
	Males	0.0439	0.0000	0.0000	0.0004	0.0012	0.0077	0.0413	0.1299	0.2529	0.2994	0.0404
	Females	0.0096	-	0.0001	0.0001	0.0007	0.0032	0.0136	0.0272	0.0400	0.0321	0.0075
White	Both Sexes	0.0285	0.0000	0.0001	0.0003	0.0010	0.0058	0.0278	0.0756	0.1364	0.1343	0.0221
	Males	0.0471	0.0000	0.0001	0.0004	0.0012	0.0080	0.0420	0.1282	0.2502	0.2983	0.0401
	Females	0.0102	-	0.0001	0.0001	0.0008	0.0036	0.0140	0.0267	0.0396	0.0316	0.0075
Black	Both Sexes	0.0240	-	-	0.0004	0.0010	0.0053	0.0313	0.1005	0.1771	0.1497	0.0273
	Males	0.0393	-	-	0.0007	0.0015	0.0088	0.0478	0.1771	0.3500	0.3748	0.0530
	Females	0.0099	-	-	-	0.0005	0.0022	0.0168	0.0393	0.0567	0.0386	0.0099
Asian/ Pacific Islander	Both Sexes	0.0075	-	-	-	0.0005	0.0010	0.0071	0.0257	0.0519	0.0726	0.0084
	Males	0.0131	-	-	-	-	0.0016	0.0134	0.0474	0.1038	0.1530	0.0166
	Females	0.0022	-	-	-	0.0008	0.0004	0.0017	0.0081	0.0102	0.0209	0.0023
Hispanic	Both Sexes	0.0093	-	0.0001	0.0003	0.0005	0.0040	0.0156	0.0463	0.0930	0.1191	0.0156
	Males	0.0158	-	0.0001	0.0004	0.0007	0.0058	0.0246	0.0859	0.1955	0.2728	0.0312
	Females	0.0027	-	0.0001	0.0002	0.0004	0.0022	0.0068	0.0113	0.0144	0.0241	0.0037

^a US 2016 cancer prevalence counts are based on 2016 cancer prevalence proportions from the SEER 13 Areas (excluding the Alaska Native Registry) and 1/1/2016 US population estimates based on the average of 2015 and 2016 population estimates from the US Bureau of the Census.

^b Prevalence was calculated using the first invasive tumor for each cancer site diagnosed during the previous 24 years. Maximum limited-duration prevalence is 24 years for 1992-2016 data.

^c Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^d Cases diagnosed more than 24 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000).

^e Complete prevalence is obtained by summing 0 to <24 and >=24.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 13.29
LeukemiaEstimated United States Cancer Prevalence Counts^a on January 1, 2016
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <24	0 to <24 ^b	>=24 ^d	Complete ^e
<u>Race</u>	<u>Sex</u>								
All Races	Both Sexes	153,253	100,309	63,581	38,012	21,353	376,508	38,265	414,773
	Males	90,192	57,683	36,029	21,220	12,020	217,145	18,411	235,556
	Females	63,061	42,625	27,552	16,792	9,333	159,363	19,854	179,217
White	Both Sexes	132,173	87,961	56,051	33,596	19,035	328,814	32,999	370,135
	Males	78,368	50,894	32,033	18,770	10,845	190,910	15,102	210,930
	Females	53,805	37,067	24,018	14,826	8,190	137,905	17,897	159,205
Black	Both Sexes	12,734	6,564	4,233	2,487	1,259	27,277	4,070	31,347
	Males	6,847	3,556	2,204	1,334	637	14,578	1,776	16,354
	Females	5,887	3,008	2,029	1,153	622	12,699	2,294	14,992
Asian/ Pacific Islander	Both Sexes	3,926	2,561	1,778	1,164	690	10,120	+	+
	Males	2,235	1,462	936	646	351	5,630	+	+
	Females	1,691	1,100	842	519	339	4,490	+	+
Hispanic	Both Sexes	14,464	9,581	6,796	4,462	2,630	37,933	+	+
	Males	8,166	5,374	3,794	2,408	1,538	21,280	+	+
	Females	6,298	4,207	3,002	2,055	1,092	16,654	+	+

Estimated prevalence percent^a on January 1, 2016, of the SEER population diagnosed in the previous 24 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted ^c	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races	Both Sexes	0.1168	0.0328	0.0640	0.0628	0.0377	0.0561	0.1053	0.2263	0.4023	0.5111	0.1054
	Males	0.1369	0.0339	0.0706	0.0685	0.0419	0.0657	0.1287	0.2888	0.5282	0.7048	0.1306
	Females	0.0974	0.0317	0.0572	0.0570	0.0333	0.0466	0.0829	0.1696	0.2974	0.3919	0.0844
White	Both Sexes	0.1304	0.0369	0.0715	0.0712	0.0410	0.0584	0.1106	0.2389	0.4249	0.5386	0.1127
	Males	0.1528	0.0379	0.0789	0.0771	0.0457	0.0682	0.1349	0.3042	0.5558	0.7387	0.1389
	Females	0.1084	0.0359	0.0637	0.0650	0.0362	0.0484	0.0869	0.1784	0.3135	0.4133	0.0904
Black	Both Sexes	0.0605	0.0145	0.0322	0.0320	0.0267	0.0463	0.0752	0.1455	0.2470	0.3076	0.0660
	Males	0.0674	0.0142	0.0357	0.0338	0.0281	0.0530	0.0919	0.1737	0.3249	0.4580	0.0816
	Females	0.0541	0.0148	0.0286	0.0301	0.0255	0.0402	0.0605	0.1230	0.1927	0.2334	0.0552
Asian/ Pacific Islander	Both Sexes	0.0497	0.0264	0.0531	0.0426	0.0230	0.0390	0.0553	0.0882	0.1245	0.1389	0.0513
	Males	0.0577	0.0298	0.0555	0.0507	0.0249	0.0461	0.0665	0.1122	0.1552	0.1904	0.0613
	Females	0.0423	0.0228	0.0507	0.0345	0.0212	0.0327	0.0456	0.0688	0.0999	0.1057	0.0430
Hispanic	Both Sexes	0.0664	0.0375	0.0737	0.0728	0.0395	0.0485	0.0723	0.1191	0.1987	0.2074	0.0746
	Males	0.0737	0.0397	0.0827	0.0807	0.0471	0.0576	0.0831	0.1350	0.2221	0.2551	0.0854
	Females	0.0589	0.0352	0.0642	0.0641	0.0312	0.0391	0.0617	0.1050	0.1809	0.1780	0.0647

^a US 2016 cancer prevalence counts are based on 2016 cancer prevalence proportions from the SEER 13 Areas (excluding the Alaska Native Registry) and 1/1/2016 US population estimates based on the average of 2015 and 2016 population estimates from the US Bureau of the Census.

^b Prevalence was calculated using the first invasive tumor for each cancer site diagnosed during the previous 24 years. Maximum limited-duration prevalence is 24 years for 1992-2016 data.

^c Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^d Cases diagnosed more than 24 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000).

^e Complete prevalence is obtained by summing 0 to <24 and >=24.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 13.30
Acute Lymphocytic Leukemia

Estimated United States Cancer Prevalence Counts^a on January 1, 2016
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <24	0 to <24 ^b	>=24 ^d	Complete ^e
<u>Race</u>	<u>Sex</u>								
All Races	Both Sexes	20,277	16,033	13,721	12,074	8,203	70,308	25,456	95,764
	Males	11,203	9,049	7,751	6,542	4,634	39,179	12,997	52,176
	Females	9,074	6,984	5,971	5,531	3,568	31,129	12,460	43,589
White	Both Sexes	17,123	13,855	11,870	10,663	7,154	60,665	21,823	83,282
	Males	9,549	7,741	6,768	5,696	4,084	33,839	11,286	45,559
	Females	7,573	6,114	5,102	4,967	3,070	26,826	10,537	37,723
Black	Both Sexes	1,590	1,133	923	727	553	4,926	1,609	6,535
	Males	809	734	473	411	290	2,716	592	3,308
	Females	781	399	450	317	263	2,210	1,017	3,227
Asian/ Pacific Islander	Both Sexes	998	720	669	486	361	3,234	+	+
	Males	527	406	369	313	178	1,793	+	+
	Females	472	314	300	173	183	1,441	+	+
Hispanic	Both Sexes	5,905	4,543	3,352	2,717	1,749	18,266	+	+
	Males	3,302	2,686	1,899	1,465	1,058	10,409	+	+
	Females	2,603	1,857	1,454	1,252	692	7,857	+	+

Estimated prevalence percent^a on January 1, 2016, of the SEER population diagnosed in the previous 24 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)										Age-Adjusted ^c
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races	Both Sexes	0.0218	0.0276	0.0541	0.0467	0.0139	0.0078	0.0059	0.0065	0.0053	0.0033	0.0228
	Males	0.0247	0.0286	0.0591	0.0520	0.0163	0.0095	0.0066	0.0069	0.0056	0.0035	0.0251
	Females	0.0190	0.0266	0.0488	0.0410	0.0115	0.0062	0.0053	0.0062	0.0050	0.0032	0.0204
White	Both Sexes	0.0241	0.0315	0.0612	0.0542	0.0161	0.0086	0.0063	0.0070	0.0056	0.0033	0.0259
	Males	0.0271	0.0323	0.0670	0.0598	0.0186	0.0107	0.0069	0.0074	0.0060	0.0032	0.0284
	Females	0.0211	0.0306	0.0551	0.0482	0.0135	0.0064	0.0057	0.0066	0.0053	0.0033	0.0232
Black	Both Sexes	0.0109	0.0107	0.0250	0.0197	0.0067	0.0042	0.0039	0.0029	0.0012	0.0027	0.0102
	Males	0.0126	0.0112	0.0281	0.0224	0.0079	0.0035	0.0052	0.0026	0.0012	0.0027	0.0112
	Females	0.0094	0.0102	0.0219	0.0171	0.0056	0.0047	0.0028	0.0032	0.0013	0.0027	0.0092
Asian/ Pacific Islander	Both Sexes	0.0159	0.0208	0.0418	0.0289	0.0071	0.0064	0.0045	0.0050	0.0054	0.0036	0.0163
	Males	0.0184	0.0226	0.0427	0.0360	0.0086	0.0069	0.0055	0.0054	0.0033	0.0055	0.0180
	Females	0.0136	0.0190	0.0408	0.0219	0.0058	0.0059	0.0036	0.0047	0.0071	0.0023	0.0146
Hispanic	Both Sexes	0.0320	0.0328	0.0637	0.0574	0.0165	0.0107	0.0093	0.0100	0.0084	0.0012	0.0279
	Males	0.0361	0.0342	0.0710	0.0648	0.0205	0.0131	0.0110	0.0063	0.0072	-	0.0309
	Females	0.0278	0.0312	0.0562	0.0494	0.0122	0.0082	0.0077	0.0133	0.0093	0.0020	0.0246

^a US 2016 cancer prevalence counts are based on 2016 cancer prevalence proportions from the SEER 13 Areas (excluding the Alaska Native Registry) and 1/1/2016 US population estimates based on the average of 2015 and 2016 population estimates from the US Bureau of the Census.

^b Prevalence was calculated using the first invasive tumor for each cancer site diagnosed during the previous 24 years. Maximum limited-duration prevalence is 24 years for 1992-2016 data.

^c Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^d Cases diagnosed more than 24 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000).

^e Complete prevalence is obtained by summing 0 to <24 and >=24.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 14.21
Cancer of the Liver and Intrahepatic Bile Duct (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2016
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <24	0 to <24 ^b	>=24 ^d	Complete ^e
<u>Race</u>	<u>Sex</u>								
All Races	Both Sexes	56,127	15,028	5,865	2,197	850	80,067	3,015	83,081
	Males	39,846	10,628	3,814	1,417	520	56,224	875	57,099
	Females	16,282	4,400	2,052	780	330	23,843	2,139	25,982
White	Both Sexes	42,087	11,264	4,300	1,603	655	59,910	+	+
	Males	29,951	8,010	2,834	1,000	423	42,217	+	+
	Females	12,137	3,255	1,466	603	232	17,693	+	+
Black	Both Sexes	7,886	1,509	504	122	56	10,077	+	+
	Males	5,574	1,035	241	84	9	6,942	+	+
	Females	2,313	474	263	38	47	3,135	+	+
Asian/ Pacific Islander	Both Sexes	4,781	1,883	903	395	88	8,051	+	+
	Males	3,311	1,335	618	284	64	5,612	+	+
	Females	1,470	548	285	111	24	2,438	+	+
Hispanic	Both Sexes	8,095	2,114	767	280	146	11,401	+	+
	Males	5,570	1,495	502	170	101	7,839	+	+
	Females	2,525	618	265	110	45	3,562	+	+

Estimated prevalence percent^a on January 1, 2016, of the SEER population diagnosed in the previous 24 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)										Age-Adjusted ^c
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races	Both Sexes	0.0248	0.0029	0.0034	0.0018	0.0018	0.0065	0.0320	0.0978	0.0856	0.0598	0.0200
	Males	0.0354	0.0030	0.0036	0.0021	0.0018	0.0081	0.0488	0.1558	0.1226	0.0934	0.0296
	Females	0.0146	0.0027	0.0032	0.0015	0.0018	0.0048	0.0159	0.0452	0.0549	0.0391	0.0117
White	Both Sexes	0.0238	0.0032	0.0038	0.0020	0.0017	0.0057	0.0288	0.0875	0.0760	0.0533	0.0181
	Males	0.0338	0.0032	0.0040	0.0023	0.0017	0.0072	0.0437	0.1395	0.1070	0.0825	0.0265
	Females	0.0139	0.0032	0.0036	0.0017	0.0016	0.0042	0.0142	0.0394	0.0497	0.0350	0.0107
Black	Both Sexes	0.0223	0.0009	0.0012	0.0009	0.0022	0.0051	0.0337	0.1399	0.0884	0.0470	0.0221
	Males	0.0321	0.0014	0.0013	0.0014	0.0017	0.0057	0.0474	0.2297	0.1460	0.0817	0.0349
	Females	0.0134	0.0003	0.0011	0.0005	0.0026	0.0046	0.0218	0.0681	0.0482	0.0298	0.0124
Asian/ Pacific Islander	Both Sexes	0.0395	0.0035	0.0039	0.0012	0.0021	0.0145	0.0618	0.1493	0.2298	0.2044	0.0420
	Males	0.0575	0.0041	0.0044	0.0010	0.0027	0.0207	0.1065	0.2419	0.3403	0.3151	0.0650
	Females	0.0230	0.0029	0.0033	0.0014	0.0015	0.0090	0.0231	0.0746	0.1411	0.1332	0.0236
Hispanic	Both Sexes	0.0200	0.0037	0.0037	0.0017	0.0013	0.0066	0.0455	0.1274	0.1506	0.0898	0.0288
	Males	0.0272	0.0040	0.0043	0.0016	0.0014	0.0094	0.0703	0.2040	0.2022	0.1299	0.0419
	Females	0.0126	0.0034	0.0030	0.0018	0.0011	0.0037	0.0211	0.0596	0.1110	0.0651	0.0175

^a US 2016 cancer prevalence counts are based on 2016 cancer prevalence proportions from the SEER 13 Areas (excluding the Alaska Native Registry) and 1/1/2016 US population estimates based on the average of 2015 and 2016 population estimates from the US Bureau of the Census.

^b Prevalence was calculated using the first invasive tumor for each cancer site diagnosed during the previous 24 years. Maximum limited-duration prevalence is 24 years for 1992-2016 data.

^c Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^d Cases diagnosed more than 24 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000).

^e Complete prevalence is obtained by summing 0 to <24 and >=24.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 15.27
Cancer of the Lung And Bronchus (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2016
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <24	0 to <24 ^b	>=24 ^d	Complete ^e
<u>Race</u>	<u>Sex</u>								
All Races	Both Sexes	296,258	106,837	52,045	27,542	12,276	494,958	43,284	538,243
	Males	135,269	44,604	20,996	11,092	5,095	217,057	22,799	239,856
	Females	160,989	62,233	31,049	16,450	7,181	277,901	20,485	298,387
White	Both Sexes	247,227	91,706	45,073	24,033	10,557	418,596	26,947	447,921
	Males	112,082	37,908	17,888	9,484	4,290	181,652	12,781	195,508
	Females	135,145	53,798	27,185	14,548	6,267	236,944	14,166	252,413
Black	Both Sexes	34,379	10,866	5,121	2,396	1,283	54,045	2,986	57,031
	Males	16,053	4,731	2,283	1,075	592	24,734	1,520	26,254
	Females	18,326	6,134	2,838	1,322	691	29,311	1,466	30,777
Asian/ Pacific Islander	Both Sexes	11,557	3,291	1,527	827	294	17,495	+	+
	Males	5,652	1,530	702	404	144	8,433	+	+
	Females	5,905	1,761	824	423	150	9,063	+	+
Hispanic	Both Sexes	12,554	4,359	2,023	1,048	477	20,462	+	+
	Males	5,566	1,752	878	418	203	8,817	+	+
	Females	6,988	2,607	1,145	630	274	11,645	+	+

Estimated prevalence percent^a on January 1, 2016, of the SEER population diagnosed in the previous 24 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)										Age-Adjusted ^c
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races	Both Sexes	0.1536	0.0002	0.0005	0.0017	0.0063	0.0250	0.1217	0.3745	0.8893	0.9606	0.1308
	Males	0.1368	0.0002	0.0006	0.0016	0.0057	0.0217	0.1031	0.3611	0.8754	1.0432	0.1290
	Females	0.1699	0.0002	0.0003	0.0019	0.0069	0.0283	0.1395	0.3867	0.9009	0.9099	0.1332
White	Both Sexes	0.1660	0.0002	0.0005	0.0018	0.0066	0.0251	0.1165	0.3666	0.9048	0.9817	0.1313
	Males	0.1454	0.0002	0.0006	0.0015	0.0058	0.0217	0.0977	0.3429	0.8725	1.0468	0.1270
	Females	0.1863	0.0002	0.0005	0.0022	0.0074	0.0286	0.1348	0.3885	0.9322	0.9410	0.1358
Black	Both Sexes	0.1198	0.0004	0.0004	0.0017	0.0058	0.0254	0.1655	0.4706	0.8818	0.8197	0.1378
	Males	0.1143	0.0004	0.0008	0.0024	0.0066	0.0232	0.1414	0.5274	0.9944	1.0415	0.1534
	Females	0.1249	0.0003	-	0.0010	0.0050	0.0274	0.1864	0.4251	0.8034	0.7103	0.1286
Asian/ Pacific Islander	Both Sexes	0.0859	-	-	0.0011	0.0048	0.0214	0.0938	0.2817	0.6035	0.7546	0.0963
	Males	0.0864	-	-	0.0012	0.0037	0.0181	0.0915	0.3086	0.6896	0.9019	0.1076
	Females	0.0854	-	-	0.0011	0.0058	0.0243	0.0959	0.2599	0.5343	0.6598	0.0882
Hispanic	Both Sexes	0.0358	0.0001	0.0002	0.0015	0.0040	0.0126	0.0507	0.1569	0.4091	0.5149	0.0619
	Males	0.0306	0.0001	0.0003	0.0011	0.0034	0.0122	0.0426	0.1436	0.3983	0.5928	0.0618
	Females	0.0412	-	0.0001	0.0019	0.0047	0.0130	0.0586	0.1687	0.4173	0.4668	0.0626

^a US 2016 cancer prevalence counts are based on 2016 cancer prevalence proportions from the SEER 13 Areas (excluding the Alaska Native Registry) and 1/1/2016 US population estimates based on the average of 2015 and 2016 population estimates from the US Bureau of the Census.

^b Prevalence was calculated using the first invasive tumor for each cancer site diagnosed during the previous 24 years. Maximum limited-duration prevalence is 24 years for 1992-2016 data.

^c Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^d Cases diagnosed more than 24 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000).

^e Complete prevalence is obtained by summing 0 to <24 and >=24.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 16.21
Melanoma of the Skin (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2016
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <24	0 to <24 ^b	>=24 ^d	Complete ^e
<u>Race</u>	<u>Sex</u>								
All Races	Both Sexes	340,956	252,815	188,823	141,530	79,525	1,003,650	191,958	1,195,608
	Males	191,955	135,055	95,771	69,992	37,823	530,596	80,562	611,158
	Females	149,001	117,760	93,052	71,538	41,703	473,054	111,397	584,451
White	Both Sexes	322,955	243,898	182,478	134,944	75,013	959,288	187,178	1,183,599
	Males	182,373	130,757	92,628	66,932	35,701	508,391	79,164	606,511
	Females	140,583	113,140	89,850	68,011	39,313	450,897	108,015	577,088
Black	Both Sexes	1,210	817	583	521	306	3,437	518	3,955
	Males	543	357	246	204	181	1,531	238	1,769
	Females	667	459	337	317	126	1,906	281	2,186
Asian/ Pacific Islander	Both Sexes	1,037	645	549	390	151	2,772	+	+
	Males	503	286	214	153	67	1,223	+	+
	Females	534	359	335	238	84	1,549	+	+
Hispanic	Both Sexes	7,370	4,706	3,769	2,800	1,400	20,045	+	+
	Males	3,099	1,935	1,178	980	464	7,655	+	+
	Females	4,271	2,772	2,591	1,820	936	12,390	+	+

Estimated prevalence percent^a on January 1, 2016, of the SEER population diagnosed in the previous 24 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)										Age-Adjusted ^c
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races	Both Sexes	0.3115	0.0004	0.0028	0.0272	0.1112	0.2434	0.4490	0.7350	1.1135	1.3328	0.2713
	Males	0.3345	0.0003	0.0025	0.0185	0.0749	0.1879	0.4233	0.8607	1.5266	2.0883	0.3122
	Females	0.2892	0.0006	0.0032	0.0363	0.1477	0.2980	0.4736	0.6212	0.7693	0.8681	0.2449
White	Both Sexes	0.3804	0.0005	0.0034	0.0337	0.1372	0.2949	0.5285	0.8476	1.2565	1.4827	0.3142
	Males	0.4069	0.0004	0.0030	0.0229	0.0916	0.2232	0.4920	0.9822	1.7036	2.3064	0.3547
	Females	0.3544	0.0006	0.0039	0.0451	0.1843	0.3673	0.5640	0.7228	0.8759	0.9669	0.2892
Black	Both Sexes	0.0076	-	0.0005	0.0006	0.0044	0.0068	0.0113	0.0205	0.0395	0.0556	0.0088
	Males	0.0071	-	0.0003	0.0005	0.0026	0.0047	0.0098	0.0236	0.0477	0.0851	0.0097
	Females	0.0081	-	0.0008	0.0007	0.0060	0.0086	0.0126	0.0180	0.0338	0.0410	0.0085
Asian/ Pacific Islander	Both Sexes	0.0136	0.0006	0.0011	0.0025	0.0068	0.0143	0.0244	0.0345	0.0489	0.0657	0.0141
	Males	0.0125	-	0.0008	0.0024	0.0060	0.0114	0.0209	0.0364	0.0582	0.0775	0.0142
	Females	0.0146	0.0011	0.0013	0.0027	0.0075	0.0170	0.0274	0.0329	0.0415	0.0582	0.0142
Hispanic	Both Sexes	0.0351	0.0002	0.0012	0.0048	0.0188	0.0438	0.0841	0.1385	0.1969	0.2490	0.0500
	Males	0.0265	0.0002	0.0008	0.0033	0.0100	0.0251	0.0587	0.1211	0.2071	0.3352	0.0450
	Females	0.0439	0.0002	0.0017	0.0064	0.0282	0.0631	0.1092	0.1539	0.1891	0.1958	0.0562

^a US 2016 cancer prevalence counts are based on 2016 cancer prevalence proportions from the SEER 13 Areas (excluding the Alaska Native Registry) and 1/1/2016 US population estimates based on the average of 2015 and 2016 population estimates from the US Bureau of the Census.

^b Prevalence was calculated using the first invasive tumor for each cancer site diagnosed during the previous 24 years. Maximum limited-duration prevalence is 24 years for 1992-2016 data.

^c Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^d Cases diagnosed more than 24 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000).

^e Complete prevalence is obtained by summing 0 to <24 and >=24.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 17.13
MesotheliomaEstimated United States Cancer Prevalence Counts^a on January 1, 2016
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <24	0 to <24 ^b	>=24 ^d	Complete ^e
<u>Race</u>	<u>Sex</u>								
All Races	Both Sexes	4,257	693	358	285	174	5,767	+	+
	Males	2,983	368	172	109	55	3,687	+	+
	Females	1,274	324	186	176	120	2,080	+	+
White	Both Sexes	3,946	658	302	237	153	5,296	+	+
	Males	2,776	368	152	101	51	3,449	+	+
	Females	1,170	290	150	136	102	1,847	+	+
Black	Both Sexes	184	34	26	35	9	289	+	+
	Males	112	0	9	8	0	129	+	+
	Females	72	34	17	27	9	160	+	+
Asian/ Pacific Islander	Both Sexes	57	0	17	3	3	81	+	+
	Males	40	0	7	0	3	50	+	+
	Females	17	0	10	3	0	30	+	+
Hispanic	Both Sexes	354	66	14	48	5	487	+	+
	Males	245	25	3	38	0	310	+	+
	Females	109	41	11	11	5	177	+	+

Estimated prevalence percent^a on January 1, 2016, of the SEER population diagnosed in the previous 24 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)										Age-Adjusted ^c
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races	Both Sexes	0.0018	-	-	0.0001	0.0004	0.0009	0.0016	0.0038	0.0095	0.0104	0.0016
	Males	0.0023	-	-	0.0000	0.0003	0.0007	0.0015	0.0050	0.0153	0.0196	0.0023
	Females	0.0013	-	-	0.0001	0.0005	0.0010	0.0016	0.0027	0.0048	0.0048	0.0011
White	Both Sexes	0.0021	-	-	0.0001	0.0004	0.0010	0.0018	0.0042	0.0103	0.0112	0.0017
	Males	0.0028	-	-	0.0000	0.0003	0.0008	0.0017	0.0056	0.0168	0.0210	0.0025
	Females	0.0015	-	-	0.0001	0.0005	0.0012	0.0019	0.0028	0.0048	0.0051	0.0012
Black	Both Sexes	0.0006	-	-	-	0.0001	0.0006	0.0003	0.0016	0.0054	0.0069	0.0008
	Males	0.0006	-	-	-	0.0003	0.0006	0.0003	0.0015	0.0035	0.0127	0.0009
	Females	0.0007	-	-	-	-	0.0006	0.0003	0.0016	0.0066	0.0040	0.0008
Asian/ Pacific Islander	Both Sexes	0.0004	-	-	-	0.0003	0.0003	0.0004	0.0013	0.0024	0.0014	0.0004
	Males	0.0005	-	-	-	-	0.0005	-	0.0021	0.0048	-	0.0006
	Females	0.0003	-	-	-	0.0006	-	0.0005	0.0006	-	-	0.0003
Hispanic	Both Sexes	0.0009	-	-	0.0001	0.0003	0.0009	0.0015	0.0034	0.0079	0.0075	0.0013
	Males	0.0011	-	-	-	0.0001	0.0006	0.0013	0.0059	0.0149	0.0159	0.0021
	Females	0.0006	-	-	0.0002	0.0004	0.0013	0.0016	0.0012	0.0025	0.0023	0.0008

^a US 2016 cancer prevalence counts are based on 2016 cancer prevalence proportions from the SEER 13 Areas (excluding the Alaska Native Registry) and 1/1/2016 US population estimates based on the average of 2015 and 2016 population estimates from the US Bureau of the Census.

^b Prevalence was calculated using the first invasive tumor for each cancer site diagnosed during the previous 24 years. Maximum limited-duration prevalence is 24 years for 1992-2016 data.

^c Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^d Cases diagnosed more than 24 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000).

^e Complete prevalence is obtained by summing 0 to <24 and >=24.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 18.21
Myeloma (Invasive)Estimated United States Cancer Prevalence Counts^a on January 1, 2016
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <24	0 to <24 ^b	>=24 ^d	Complete ^e
Race	Sex								
All Races	Both Sexes	78,454	32,030	12,390	4,580	1,515	128,969	2,423	131,392
	Males	44,371	17,567	6,651	2,646	818	72,052	947	72,999
	Females	34,083	14,463	5,740	1,934	697	56,917	1,475	58,392
White	Both Sexes	58,668	24,166	9,726	3,553	1,216	97,330	+	+
	Males	34,273	13,673	5,361	2,174	660	56,140	+	+
	Females	24,395	10,494	4,366	1,380	556	41,190	+	+
Black	Both Sexes	16,092	6,595	2,225	772	216	25,900	+	+
	Males	8,066	3,203	1,002	354	127	12,751	+	+
	Females	8,027	3,392	1,223	418	90	13,149	+	+
Asian/ Pacific Islander	Both Sexes	2,202	821	243	129	55	3,450	+	+
	Males	1,205	447	141	52	17	1,861	+	+
	Females	998	375	102	77	38	1,589	+	+
Hispanic	Both Sexes	6,254	2,272	955	421	106	10,008	+	+
	Males	3,491	1,278	517	242	76	5,603	+	+
	Females	2,763	995	439	178	30	4,404	+	+

Estimated prevalence percent^a on January 1, 2016, of the SEER population diagnosed in the previous 24 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)										Age-Adjusted ^c
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
Race	Sex											
All Races	Both Sexes	0.0400	-	0.0000	0.0003	0.0026	0.0146	0.0464	0.1106	0.1970	0.1862	0.0337
	Males	0.0454	-	0.0000	0.0004	0.0032	0.0167	0.0540	0.1311	0.2459	0.2554	0.0416
	Females	0.0348	-	0.0000	0.0003	0.0020	0.0126	0.0391	0.0921	0.1563	0.1437	0.0274
White	Both Sexes	0.0386	-	0.0000	0.0004	0.0022	0.0126	0.0408	0.0989	0.1785	0.1734	0.0304
	Males	0.0449	-	0.0001	0.0004	0.0028	0.0144	0.0488	0.1203	0.2258	0.2447	0.0383
	Females	0.0324	-	-	0.0003	0.0017	0.0108	0.0330	0.0790	0.1383	0.1288	0.0239
Black	Both Sexes	0.0574	-	0.0001	0.0001	0.0049	0.0307	0.0885	0.2156	0.3846	0.3403	0.0652
	Males	0.0589	-	-	-	0.0059	0.0360	0.0959	0.2386	0.4730	0.4272	0.0766
	Females	0.0560	-	0.0003	0.0002	0.0040	0.0260	0.0821	0.1971	0.3231	0.2973	0.0572
Asian/ Pacific Islander	Both Sexes	0.0169	-	-	0.0003	0.0017	0.0067	0.0221	0.0594	0.1125	0.1085	0.0184
	Males	0.0191	-	-	0.0004	0.0027	0.0082	0.0259	0.0706	0.1417	0.1314	0.0226
	Females	0.0150	-	-	-	0.0008	0.0054	0.0188	0.0503	0.0891	0.0937	0.0151
Hispanic	Both Sexes	0.0175	0.0000	0.0001	0.0005	0.0026	0.0125	0.0377	0.0870	0.1684	0.1325	0.0275
	Males	0.0194	-	0.0001	0.0006	0.0030	0.0143	0.0466	0.1013	0.2056	0.1864	0.0338
	Females	0.0156	0.0001	-	0.0003	0.0022	0.0106	0.0289	0.0744	0.1399	0.0992	0.0225

^a US 2016 cancer prevalence counts are based on 2016 cancer prevalence proportions from the SEER 13 Areas (excluding the Alaska Native Registry) and 1/1/2016 US population estimates based on the average of 2015 and 2016 population estimates from the US Bureau of the Census.

^b Prevalence was calculated using the first invasive tumor for each cancer site diagnosed during the previous 24 years. Maximum limited-duration prevalence is 24 years for 1992-2016 data.

^c Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^d Cases diagnosed more than 24 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000).

^e Complete prevalence is obtained by summing 0 to <24 and >=24.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 19.25
Non-Hodgkin LymphomaEstimated United States Cancer Prevalence Counts^a on January 1, 2016
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <24	0 to <24 ^b	>=24 ^d	Complete ^e
Race	Sex								
All Races	Both Sexes	248,946	177,490	117,326	69,109	31,890	644,761	49,943	694,704
	Males	135,137	94,173	60,159	36,020	16,581	342,069	25,489	367,558
	Females	113,809	83,317	57,167	33,089	15,309	302,692	24,454	327,146
White	Both Sexes	213,764	154,454	101,984	60,537	28,026	558,765	49,625	620,368
	Males	116,527	82,172	52,535	31,627	14,543	297,404	26,526	330,121
	Females	97,237	72,282	49,448	28,910	13,483	261,361	23,099	290,247
Black	Both Sexes	20,500	13,640	9,750	5,271	2,215	51,376	4,115	55,491
	Males	10,712	7,188	4,768	2,684	1,174	26,526	1,911	28,437
	Females	9,789	6,452	4,981	2,587	1,041	24,850	2,204	27,054
Asian/ Pacific Islander	Both Sexes	8,405	5,210	3,415	1,867	844	19,741	+	+
	Males	4,548	2,661	1,757	1,013	407	10,386	+	+
	Females	3,856	2,549	1,657	855	437	9,355	+	+
Hispanic	Both Sexes	21,339	14,049	8,272	4,881	2,136	50,676	+	+
	Males	10,918	7,243	4,331	2,617	1,094	26,202	+	+
	Females	10,421	6,805	3,941	2,264	1,043	24,474	+	+

Estimated prevalence percent^a on January 1, 2016, of the SEER population diagnosed in the previous 24 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)										Age-Adjusted ^c
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
Race	Sex											
All Races	Both Sexes	0.2001	0.0035	0.0139	0.0283	0.0538	0.1121	0.2352	0.4640	0.8181	1.0224	0.1738
	Males	0.2156	0.0044	0.0189	0.0354	0.0625	0.1293	0.2710	0.5243	0.9364	1.2449	0.2023
	Females	0.1850	0.0026	0.0087	0.0208	0.0450	0.0952	0.2010	0.4094	0.7196	0.8856	0.1498
White	Both Sexes	0.2216	0.0035	0.0143	0.0289	0.0534	0.1148	0.2427	0.4858	0.8597	1.0738	0.1807
	Males	0.2380	0.0045	0.0198	0.0359	0.0622	0.1317	0.2790	0.5442	0.9778	1.3060	0.2095
	Females	0.2054	0.0025	0.0085	0.0214	0.0442	0.0976	0.2072	0.4317	0.7593	0.9285	0.1560
Black	Both Sexes	0.1139	0.0029	0.0113	0.0265	0.0599	0.1027	0.2010	0.3340	0.5117	0.5333	0.1251
	Males	0.1226	0.0029	0.0140	0.0351	0.0696	0.1223	0.2333	0.3969	0.5849	0.6055	0.1457
	Females	0.1059	0.0030	0.0086	0.0178	0.0510	0.0853	0.1728	0.2836	0.4608	0.4976	0.1088
Asian/ Pacific Islander	Both Sexes	0.0970	0.0039	0.0133	0.0207	0.0353	0.0680	0.1409	0.2692	0.4470	0.6094	0.1032
	Males	0.1065	0.0044	0.0175	0.0240	0.0398	0.0744	0.1640	0.3072	0.5516	0.7472	0.1222
	Females	0.0882	0.0033	0.0092	0.0175	0.0312	0.0623	0.1209	0.2384	0.3629	0.5207	0.0880
Hispanic	Both Sexes	0.0887	0.0030	0.0117	0.0206	0.0398	0.0850	0.1769	0.3632	0.5888	0.7209	0.1294
	Males	0.0908	0.0035	0.0147	0.0256	0.0455	0.0932	0.1924	0.3940	0.6326	0.8003	0.1420
	Females	0.0866	0.0024	0.0086	0.0151	0.0337	0.0766	0.1616	0.3360	0.5552	0.6719	0.1185

^a US 2016 cancer prevalence counts are based on 2016 cancer prevalence proportions from the SEER 13 Areas (excluding the Alaska Native Registry) and 1/1/2016 US population estimates based on the average of 2015 and 2016 population estimates from the US Bureau of the Census.

^b Prevalence was calculated using the first invasive tumor for each cancer site diagnosed during the previous 24 years. Maximum limited-duration prevalence is 24 years for 1992-2016 data.

^c Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^d Cases diagnosed more than 24 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000).

^e Complete prevalence is obtained by summing 0 to <24 and >=24.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 20.27
Cancer of the Oral Cavity and Pharynx (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2016
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <24	0 to <24 ^b	>=24 ^d	Complete ^e
Race	Sex								
All Races	Both Sexes	137,601	83,078	52,272	33,186	16,846	322,983	47,326	370,309
	Males	97,382	56,660	34,723	21,268	10,139	220,172	29,888	250,059
	Females	40,219	26,418	17,549	11,918	6,707	102,811	17,438	120,250
White	Both Sexes	119,389	71,998	45,504	28,485	14,475	279,851	40,358	325,100
	Males	85,471	49,690	30,750	18,599	8,853	193,362	25,529	222,024
	Females	33,919	22,309	14,754	9,885	5,622	86,489	14,829	103,077
Black	Both Sexes	10,229	5,844	3,613	2,443	1,333	23,462	2,368	25,830
	Males	6,966	3,712	2,101	1,354	678	14,811	1,246	16,057
	Females	3,262	2,132	1,512	1,090	655	8,651	1,121	9,773
Asian/ Pacific Islander	Both Sexes	5,206	3,370	2,197	1,497	670	12,941	+	+
	Males	3,161	2,040	1,290	850	380	7,721	+	+
	Females	2,045	1,330	907	647	290	5,220	+	+
Hispanic	Both Sexes	7,864	4,600	2,787	1,690	868	17,809	+	+
	Males	5,180	2,704	1,697	983	418	10,982	+	+
	Females	2,684	1,896	1,090	707	450	6,828	+	+

Estimated prevalence percent^a on January 1, 2016, of the SEER population diagnosed in the previous 24 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		All Ages	0-9	10-19	20-29	Age Specific (Crude)					Age-Adjusted ^c	
Race	Sex											All Ages
All Races	Both Sexes	0.1002	0.0004	0.0023	0.0066	0.0168	0.0488	0.1514	0.2996	0.3762	0.3571	0.0835
	Males	0.1388	0.0004	0.0023	0.0062	0.0162	0.0604	0.2212	0.4620	0.5721	0.5369	0.1218
	Females	0.0629	0.0004	0.0024	0.0071	0.0174	0.0374	0.0848	0.1524	0.2130	0.2464	0.0509
White	Both Sexes	0.1110	0.0004	0.0021	0.0065	0.0162	0.0480	0.1574	0.3148	0.3973	0.3757	0.0868
	Males	0.1548	0.0004	0.0020	0.0059	0.0153	0.0603	0.2321	0.4855	0.6023	0.5628	0.1269
	Females	0.0680	0.0004	0.0023	0.0072	0.0172	0.0356	0.0846	0.1566	0.2228	0.2585	0.0518
Black	Both Sexes	0.0520	0.0004	0.0029	0.0068	0.0144	0.0395	0.1077	0.2014	0.2167	0.1750	0.0546
	Males	0.0685	0.0005	0.0031	0.0073	0.0142	0.0444	0.1503	0.3108	0.3419	0.2814	0.0791
	Females	0.0369	0.0003	0.0027	0.0063	0.0146	0.0352	0.0705	0.1141	0.1295	0.1224	0.0366
Asian/ Pacific Islander	Both Sexes	0.0636	-	0.0022	0.0061	0.0237	0.0607	0.1289	0.1964	0.2441	0.2337	0.0650
	Males	0.0792	-	0.0027	0.0058	0.0268	0.0733	0.1718	0.2771	0.3360	0.3056	0.0859
	Females	0.0492	-	0.0017	0.0064	0.0208	0.0496	0.0918	0.1312	0.1703	0.1873	0.0479
Hispanic	Both Sexes	0.0312	0.0005	0.0019	0.0048	0.0093	0.0284	0.0713	0.1499	0.2128	0.2184	0.0456
	Males	0.0381	0.0006	0.0019	0.0047	0.0086	0.0323	0.0941	0.2145	0.2996	0.3112	0.0616
	Females	0.0242	0.0003	0.0019	0.0050	0.0100	0.0243	0.0489	0.0927	0.1463	0.1610	0.0326

^a US 2016 cancer prevalence counts are based on 2016 cancer prevalence proportions from the SEER 13 Areas (excluding the Alaska Native Registry) and 1/1/2016 US population estimates based on the average of 2015 and 2016 population estimates from the US Bureau of the Census.

^b Prevalence was calculated using the first invasive tumor for each cancer site diagnosed during the previous 24 years. Maximum limited-duration prevalence is 24 years for 1992-2016 data.

^c Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^d Cases diagnosed more than 24 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000).

^e Complete prevalence is obtained by summing 0 to <24 and >=24.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 21.15
Cancer of the Ovary (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2016
By Race/Ethnicity and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <24	0 to <24 ^b	>=24 ^d	Complete ^e
<u>Race</u>								
All Races	71,655	40,457	29,594	22,902	15,418	180,026	49,849	229,875
White	61,218	35,163	25,759	20,146	13,438	155,723	42,717	199,796
Black	5,771	2,492	1,929	1,452	1,154	12,799	4,395	17,193
Asian/Pacific Islander	3,549	2,137	1,546	1,001	605	8,838	+	+
Hispanic	7,562	4,112	2,702	1,700	1,135	17,210	+	+

Estimated prevalence percent^a on January 1, 2016, of the SEER population diagnosed in the previous 24 years
By Age at Prevalence and Race/Ethnicity

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^c
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>											
All Races	0.1101	0.0005	0.0055	0.0202	0.0387	0.0814	0.1760	0.2810	0.3388	0.2710	0.0906
White	0.1224	0.0006	0.0054	0.0212	0.0405	0.0843	0.1857	0.3014	0.3631	0.2861	0.0958
Black	0.0545	0.0003	0.0053	0.0147	0.0310	0.0582	0.1029	0.1520	0.1822	0.1496	0.0544
Asian/Pacific Islander	0.0833	0.0008	0.0050	0.0193	0.0362	0.0868	0.1850	0.2315	0.2271	0.1867	0.0790
Hispanic	0.0609	0.0006	0.0066	0.0254	0.0401	0.0724	0.1442	0.2241	0.2500	0.2174	0.0761

^a Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

^a US 2016 cancer prevalence counts are based on 2016 cancer prevalence proportions from the SEER 13 Areas (excluding the Alaska Native Registry) and 1/1/2016 US population estimates based on the average of 2015 and 2016 population estimates from the US Bureau of the Census.

^b Prevalence was calculated using the first invasive tumor for each cancer site diagnosed during the previous 24 years.

^c Maximum limited-duration prevalence is 24 years for 1992-2016 data.

^d Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^d Cases diagnosed more than 24 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000).

^e Complete prevalence is obtained by summing 0 to <24 and >=24.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 22.21
Cancer of the Pancreas (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2016
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <24	0 to <24 ^b	>=24 ^d	Complete ^e
Race	Sex								
All Races	Both Sexes	53,929	10,127	4,024	1,824	832	70,735	2,819	73,554
	Males	27,124	5,187	1,759	810	208	35,088	1,330	36,418
	Females	26,805	4,940	2,265	1,014	623	35,647	1,489	37,136
White	Both Sexes	44,556	8,296	3,377	1,411	655	58,294	2,308	61,221
	Males	22,760	4,311	1,562	690	172	29,495	1,163	31,041
	Females	21,796	3,985	1,814	721	482	28,799	1,145	30,180
Black	Both Sexes	6,478	1,238	411	260	89	8,476	881	9,358
	Males	2,875	583	108	76	0	3,642	406	4,048
	Females	3,602	655	303	185	89	4,834	475	5,309
Asian/ Pacific Islander	Both Sexes	2,167	415	185	87	34	2,889	+	+
	Males	1,038	200	67	24	10	1,339	+	+
	Females	1,129	215	118	63	25	1,550	+	+
Hispanic	Both Sexes	4,279	724	362	177	96	5,638	+	+
	Males	1,873	332	121	67	6	2,400	+	+
	Females	2,406	392	241	110	89	3,238	+	+

Estimated prevalence percent^a on January 1, 2016, of the SEER population diagnosed in the previous 24 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		All Ages	0-9	10-19	20-29	Age Specific (Crude)					Age-Adjusted ^c	
Race	Sex											All Ages
All Races	Both Sexes	0.0220	0.0000	0.0003	0.0010	0.0029	0.0095	0.0272	0.0613	0.0992	0.0938	0.0186
	Males	0.0221	0.0000	0.0003	0.0005	0.0025	0.0085	0.0269	0.0682	0.1132	0.1093	0.0201
	Females	0.0218	0.0001	0.0003	0.0015	0.0034	0.0105	0.0274	0.0551	0.0876	0.0842	0.0174
White	Both Sexes	0.0231	0.0001	0.0003	0.0011	0.0029	0.0091	0.0257	0.0604	0.1002	0.0940	0.0184
	Males	0.0236	0.0001	0.0002	0.0006	0.0025	0.0081	0.0261	0.0674	0.1148	0.1103	0.0200
	Females	0.0226	0.0001	0.0003	0.0016	0.0034	0.0101	0.0253	0.0539	0.0878	0.0838	0.0171
Black	Both Sexes	0.0188	-	0.0003	0.0007	0.0024	0.0123	0.0374	0.0721	0.0961	0.0891	0.0204
	Males	0.0168	-	0.0003	0.0005	0.0017	0.0105	0.0331	0.0791	0.0999	0.0904	0.0203
	Females	0.0206	-	0.0003	0.0009	0.0030	0.0139	0.0412	0.0666	0.0934	0.0885	0.0206
Asian/ Pacific Islander	Both Sexes	0.0142	-	-	0.0008	0.0037	0.0077	0.0209	0.0502	0.0759	0.0791	0.0151
	Males	0.0137	-	-	-	0.0036	0.0083	0.0194	0.0512	0.0906	0.0812	0.0159
	Females	0.0146	-	-	0.0013	0.0038	0.0072	0.0221	0.0494	0.0642	0.0777	0.0144
Hispanic	Both Sexes	0.0099	0.0001	0.0003	0.0011	0.0027	0.0076	0.0203	0.0467	0.0812	0.0836	0.0151
	Males	0.0083	-	0.0004	0.0003	0.0014	0.0057	0.0177	0.0449	0.0958	0.0717	0.0147
	Females	0.0115	0.0001	0.0003	0.0020	0.0041	0.0095	0.0229	0.0482	0.0700	0.0909	0.0157

^a US 2016 cancer prevalence counts are based on 2016 cancer prevalence proportions from the SEER 13 Areas (excluding the Alaska Native Registry) and 1/1/2016 US population estimates based on the average of 2015 and 2016 population estimates from the US Bureau of the Census.

^b Prevalence was calculated using the first invasive tumor for each cancer site diagnosed during the previous 24 years. Maximum limited-duration prevalence is 24 years for 1992-2016 data.

^c Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^d Cases diagnosed more than 24 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000).

^e Complete prevalence is obtained by summing 0 to <24 and >=24.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 23.15
Cancer of the Prostate (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2016
By Race/Ethnicity and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <24	0 to <24 ^b	>=24 ^d	Complete ^e
<u>Race</u>								
All Races	925,846	945,574	660,380	387,565	150,033	3,069,398	41,006	3,110,403
White	724,023	764,898	550,849	327,374	129,883	2,497,026	44,331	2,628,705
Black	138,607	128,648	83,217	46,936	16,450	413,858	4,779	418,637
Asian/Pacific Islander	21,563	19,811	12,673	5,940	1,897	61,883	+	+
Hispanic	60,114	54,746	36,830	17,966	6,233	175,889	+	+

Estimated prevalence percent^a on January 1, 2016, of the SEER population diagnosed in the previous 24 years
By Age at Prevalence and Race/Ethnicity

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^c
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>											
All Races	1.9347	0.0001	0.0002	0.0002	0.0010	0.0692	1.0107	5.0637	12.8866	17.5576	1.8440
White	1.9984	0.0002	0.0003	0.0003	0.0008	0.0538	0.8859	4.7049	12.3728	17.0503	1.7533
Black	1.9131	-	-	0.0002	0.0021	0.1784	1.9464	8.3071	18.4946	22.5283	2.7086
Asian/Pacific Islander	0.6344	-	-	-	-	0.0130	0.2989	1.9809	5.7864	9.5868	0.8474
Hispanic	0.6095	0.0002	0.0002	0.0001	0.0006	0.0334	0.5553	3.2066	9.3286	13.5466	1.3218

^a US 2016 cancer prevalence counts are based on 2016 cancer prevalence proportions from the SEER 13 Areas (excluding the Alaska Native Registry) and 1/1/2016 US population estimates based on the average of 2015 and 2016 population estimates from the US Bureau of the Census.

^b Prevalence was calculated using the first invasive tumor for each cancer site diagnosed during the previous 24 years. Maximum limited-duration prevalence is 24 years for 1992-2016 data.

^c Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^d Cases diagnosed more than 24 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000).

^e Complete prevalence is obtained by summing 0 to <24 and >=24.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 24.21
Cancer of the Stomach (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2016
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <24	0 to <24 ^b	>=24 ^d	Complete ^e
<u>Race</u>	<u>Sex</u>								
All Races	Both Sexes	53,400	25,301	13,855	7,885	3,686	104,126	8,928	113,054
	Males	31,498	14,259	7,709	4,287	1,970	59,722	5,115	64,837
	Females	21,901	11,042	6,147	3,597	1,716	44,404	3,813	48,217
White	Both Sexes	40,321	19,117	10,308	5,880	2,783	78,409	6,979	86,508
	Males	24,380	11,125	5,946	3,236	1,422	46,109	3,972	50,671
	Females	15,942	7,992	4,361	2,645	1,361	32,301	3,007	35,836
Black	Both Sexes	7,845	3,338	1,909	1,092	410	14,594	1,156	15,750
	Males	4,110	1,557	905	528	253	7,352	614	7,966
	Females	3,735	1,781	1,005	564	157	7,241	542	7,784
Asian/ Pacific Islander	Both Sexes	4,048	2,310	1,377	783	400	8,918	+	+
	Males	2,382	1,251	739	441	239	5,052	+	+
	Females	1,666	1,059	638	341	161	3,865	+	+
Hispanic	Both Sexes	7,356	3,456	1,883	996	475	14,166	+	+
	Males	3,866	1,810	987	506	208	7,377	+	+
	Females	3,490	1,646	896	490	267	6,789	+	+

Estimated prevalence percent^a on January 1, 2016, of the SEER population diagnosed in the previous 24 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted ^c	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races	Both Sexes	0.0323	-	0.0001	0.0008	0.0039	0.0130	0.0354	0.0759	0.1462	0.2055	0.0277
	Males	0.0376	-	0.0001	0.0007	0.0037	0.0133	0.0408	0.0959	0.2002	0.2791	0.0355
	Females	0.0271	-	0.0002	0.0010	0.0041	0.0128	0.0302	0.0578	0.1012	0.1603	0.0215
White	Both Sexes	0.0311	-	0.0002	0.0008	0.0038	0.0122	0.0314	0.0679	0.1317	0.1830	0.0249
	Males	0.0369	-	0.0001	0.0006	0.0038	0.0126	0.0364	0.0862	0.1849	0.2506	0.0323
	Females	0.0254	-	0.0002	0.0009	0.0038	0.0117	0.0264	0.0510	0.0864	0.1406	0.0189
Black	Both Sexes	0.0324	-	-	0.0010	0.0036	0.0135	0.0504	0.1072	0.1999	0.2867	0.0375
	Males	0.0340	-	-	0.0007	0.0036	0.0126	0.0557	0.1330	0.2512	0.3732	0.0461
	Females	0.0309	-	-	0.0012	0.0035	0.0144	0.0458	0.0865	0.1642	0.2440	0.0319
Asian/ Pacific Islander	Both Sexes	0.0438	-	0.0003	0.0012	0.0040	0.0185	0.0503	0.1205	0.2676	0.4688	0.0496
	Males	0.0518	-	-	0.0012	0.0033	0.0200	0.0632	0.1610	0.3478	0.6499	0.0650
	Females	0.0364	-	0.0006	0.0011	0.0046	0.0171	0.0391	0.0878	0.2032	0.3523	0.0381
Hispanic	Both Sexes	0.0248	-	0.0001	0.0008	0.0057	0.0173	0.0463	0.1072	0.2083	0.3279	0.0401
	Males	0.0256	-	0.0001	0.0006	0.0057	0.0171	0.0482	0.1220	0.2769	0.3913	0.0477
	Females	0.0240	-	0.0001	0.0010	0.0058	0.0175	0.0444	0.0942	0.1556	0.2887	0.0345

^a US 2016 cancer prevalence counts are based on 2016 cancer prevalence proportions from the SEER 13 Areas (excluding the Alaska Native Registry) and 1/1/2016 US population estimates based on the average of 2015 and 2016 population estimates from the US Bureau of the Census.

^b Prevalence was calculated using the first invasive tumor for each cancer site diagnosed during the previous 24 years. Maximum limited-duration prevalence is 24 years for 1992-2016 data.

^c Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^d Cases diagnosed more than 24 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000).

^e Complete prevalence is obtained by summing 0 to <24 and >=24.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 25.15
Cancer of the Testis (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2016
By Race/Ethnicity and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <24	0 to <24 ^b	>=24 ^d	Complete ^e
<u>Race</u>								
All Races	44,309	41,137	38,167	35,138	25,040	183,792	79,346	263,137
White	40,170	37,762	35,963	33,087	23,810	170,792	75,409	249,218
Black	1,451	1,289	951	900	575	5,167	1,874	7,041
Asian/Pacific Islander	1,222	877	668	567	393	3,727	+	+
Hispanic	7,880	6,596	4,732	3,512	2,007	24,726	+	+

Estimated prevalence percent^a on January 1, 2016, of the SEER population diagnosed in the previous 24 years
By Age at Prevalence and Race/Ethnicity

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^c	
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages	
<u>Race</u>												
All Races	0.1159	0.0025	0.0068	0.0745	0.1997	0.2470	0.2202	0.1153	0.0479	0.0227	0.1165	
White	0.1367	0.0024	0.0080	0.0906	0.2380	0.2918	0.2535	0.1292	0.0529	0.0249	0.1367	
Black	0.0239	0.0029	0.0021	0.0108	0.0389	0.0516	0.0553	0.0343	0.0082	0.0027	0.0253	
Asian/Pacific Islander	0.0382	0.0031	0.0048	0.0337	0.0636	0.0692	0.0634	0.0392	0.0115	0.0061	0.0366	
Hispanic	0.0857	0.0026	0.0108	0.0898	0.1855	0.1726	0.1286	0.0561	0.0312	0.0080	0.0890	

^a US 2016 cancer prevalence counts are based on 2016 cancer prevalence proportions from the SEER 13 Areas (excluding the Alaska Native Registry) and 1/1/2016 US population estimates based on the average of 2015 and 2016 population estimates from the US Bureau of the Census.

^b Prevalence was calculated using the first invasive tumor for each cancer site diagnosed during the previous 24 years. Maximum limited-duration prevalence is 24 years for 1992-2016 data.

^c Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^d Cases diagnosed more than 24 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000).

^e Complete prevalence is obtained by summing 0 to <24 and >=24.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 26.21
Cancer of the Thyroid (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2016
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <24	0 to <24 ^b	>=24 ^d	Complete ^e
<u>Race</u>	<u>Sex</u>								
All Races	Both Sexes	229,536	178,307	117,358	75,019	44,352	644,572	177,670	822,242
	Males	55,079	39,028	24,894	15,512	9,579	144,092	41,510	185,603
	Females	174,457	139,279	92,464	59,507	34,772	500,479	136,160	636,639
White	Both Sexes	194,120	153,635	102,911	66,508	39,197	556,372	159,207	724,029
	Males	48,082	34,690	22,384	14,132	8,661	127,948	35,724	165,484
	Females	146,039	118,945	80,527	52,377	30,537	428,424	123,483	558,545
Black	Both Sexes	16,940	12,042	7,223	3,846	2,299	42,351	9,745	52,096
	Males	3,033	1,940	1,223	595	424	7,216	1,457	8,673
	Females	13,907	10,102	6,000	3,251	1,876	35,135	8,288	43,423
Asian/ Pacific Islander	Both Sexes	12,778	8,972	5,547	3,447	2,035	32,779	+	+
	Males	2,759	1,623	921	590	322	6,214	+	+
	Females	10,019	7,349	4,626	2,857	1,713	26,565	+	+
Hispanic	Both Sexes	27,552	18,698	11,559	7,140	4,052	69,000	+	+
	Males	5,118	2,980	1,911	1,134	616	11,759	+	+
	Females	22,434	15,718	9,648	6,006	3,436	57,241	+	+

Estimated prevalence percent^a on January 1, 2016, of the SEER population diagnosed in the previous 24 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)										Age-Adjusted ^c
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races	Both Sexes	0.2000	0.0002	0.0067	0.0490	0.1582	0.2959	0.3791	0.4108	0.4157	0.2932	0.1832
	Males	0.0908	0.0002	0.0027	0.0168	0.0504	0.1124	0.1612	0.2225	0.2658	0.2103	0.0842
	Females	0.3060	0.0002	0.0109	0.0827	0.2666	0.4765	0.5873	0.5814	0.5406	0.3442	0.2773
White	Both Sexes	0.2206	0.0002	0.0075	0.0560	0.1764	0.3217	0.4085	0.4282	0.4282	0.3006	0.1965
	Males	0.1024	0.0003	0.0030	0.0195	0.0559	0.1233	0.1759	0.2369	0.2790	0.2179	0.0908
	Females	0.3368	0.0002	0.0123	0.0944	0.3009	0.5223	0.6355	0.6056	0.5552	0.3523	0.2996
Black	Both Sexes	0.0939	-	0.0020	0.0150	0.0630	0.1529	0.1988	0.2628	0.2903	0.1967	0.1003
	Males	0.0334	-	0.0010	0.0033	0.0170	0.0464	0.0725	0.1103	0.1455	0.1325	0.0394
	Females	0.1497	-	0.0030	0.0268	0.1052	0.2474	0.3088	0.3845	0.3911	0.2284	0.1511
Asian/ Pacific Islander	Both Sexes	0.1610	-	0.0060	0.0417	0.1281	0.2505	0.3186	0.3812	0.3769	0.2802	0.1595
	Males	0.0637	-	0.0027	0.0132	0.0425	0.0970	0.1229	0.1725	0.2037	0.1631	0.0673
	Females	0.2505	-	0.0095	0.0703	0.2054	0.3865	0.4880	0.5498	0.5161	0.3556	0.2384
Hispanic	Both Sexes	0.1208	0.0001	0.0072	0.0410	0.1255	0.2228	0.3018	0.3534	0.3793	0.2723	0.1508
	Males	0.0407	0.0001	0.0033	0.0117	0.0307	0.0691	0.1011	0.1515	0.1986	0.1503	0.0566
	Females	0.2026	-	0.0112	0.0730	0.2280	0.3810	0.4995	0.5320	0.5178	0.3477	0.2413

^a US 2016 cancer prevalence counts are based on 2016 cancer prevalence proportions from the SEER 13 Areas (excluding the Alaska Native Registry) and 1/1/2016 US population estimates based on the average of 2015 and 2016 population estimates from the US Bureau of the Census.

^b Prevalence was calculated using the first invasive tumor for each cancer site diagnosed during the previous 24 years. Maximum limited-duration prevalence is 24 years for 1992-2016 data.

^c Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^d Cases diagnosed more than 24 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000).

^e Complete prevalence is obtained by summing 0 to <24 and >=24.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 27.21
Cancer of the Urinary Bladder (Invasive and *In Situ*)

Estimated United States Cancer Prevalence Counts^a on January 1, 2016
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <24	0 to <24 ^b	>=24 ^d	Complete ^e
<u>Race</u>	<u>Sex</u>								
All Races	Both Sexes	254,236	164,468	109,426	67,889	35,394	631,413	68,037	699,450
	Males	195,876	124,874	81,422	49,926	25,688	477,786	47,928	525,713
	Females	58,359	39,595	28,005	17,963	9,705	153,627	20,109	173,737
White	Both Sexes	229,902	149,755	100,692	63,306	32,990	576,645	54,539	640,868
	Males	177,987	114,172	75,052	46,755	23,912	437,878	38,280	483,506
	Females	51,915	35,583	25,640	16,551	9,079	138,768	16,259	157,361
Black	Both Sexes	14,181	7,937	5,008	2,693	1,478	31,297	1,899	33,196
	Males	10,118	5,683	3,495	1,763	1,077	22,137	1,506	23,642
	Females	4,063	2,254	1,513	930	401	9,161	394	9,554
Asian/ Pacific Islander	Both Sexes	5,234	3,346	2,007	1,007	495	12,088	+	+
	Males	3,973	2,547	1,523	700	378	9,121	+	+
	Females	1,261	799	484	307	116	2,967	+	+
Hispanic	Both Sexes	10,965	6,555	4,063	2,397	1,252	25,233	+	+
	Males	8,292	4,882	3,019	1,760	887	18,841	+	+
	Females	2,673	1,673	1,044	637	365	6,391	+	+

Estimated prevalence percent^a on January 1, 2016, of the SEER population diagnosed in the previous 24 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted ^c	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races	Both Sexes	0.1960	0.0003	0.0004	0.0018	0.0074	0.0285	0.1163	0.4081	1.0482	1.7281	0.1679
	Males	0.3012	0.0003	0.0005	0.0023	0.0101	0.0428	0.1788	0.6621	1.7797	3.3369	0.2938
	Females	0.0939	0.0002	0.0003	0.0011	0.0046	0.0144	0.0565	0.1781	0.4388	0.7385	0.0726
White	Both Sexes	0.2287	0.0002	0.0004	0.0018	0.0084	0.0318	0.1261	0.4405	1.1342	1.8405	0.1807
	Males	0.3504	0.0001	0.0004	0.0027	0.0115	0.0467	0.1926	0.7058	1.9111	3.5373	0.3135
	Females	0.1091	0.0002	0.0004	0.0009	0.0053	0.0168	0.0612	0.1945	0.4730	0.7782	0.0781
Black	Both Sexes	0.0694	0.0008	0.0001	0.0012	0.0032	0.0152	0.0699	0.2225	0.5057	0.8221	0.0848
	Males	0.1023	0.0011	-	0.0005	0.0038	0.0265	0.1070	0.3854	0.8725	1.5812	0.1508
	Females	0.0390	0.0005	0.0003	0.0019	0.0027	0.0052	0.0375	0.0924	0.2503	0.4474	0.0421
Asian/ Pacific Islander	Both Sexes	0.0594	0.0003	-	0.0007	0.0025	0.0100	0.0469	0.1621	0.3869	0.8154	0.0692
	Males	0.0935	0.0005	-	0.0010	0.0035	0.0154	0.0775	0.2801	0.6696	1.5166	0.1230
	Females	0.0280	-	-	0.0004	0.0017	0.0053	0.0204	0.0668	0.1597	0.3639	0.0297
Hispanic	Both Sexes	0.0442	0.0002	0.0003	0.0011	0.0044	0.0151	0.0533	0.1742	0.4869	0.8113	0.0782
	Males	0.0653	0.0003	0.0004	0.0017	0.0056	0.0211	0.0806	0.2828	0.8485	1.5600	0.1379
	Females	0.0226	0.0001	0.0002	0.0006	0.0031	0.0090	0.0265	0.0781	0.2097	0.3490	0.0347

^a US 2016 cancer prevalence counts are based on 2016 cancer prevalence proportions from the SEER 13 Areas (excluding the Alaska Native Registry) and 1/1/2016 US population estimates based on the average of 2015 and 2016 population estimates from the US Bureau of the Census.

^b Prevalence was calculated using the first invasive tumor for each cancer site diagnosed during the previous 24 years. Maximum limited-duration prevalence is 24 years for 1992-2016 data.

^c Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^d Cases diagnosed more than 24 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000).

^e Complete prevalence is obtained by summing 0 to <24 and >=24.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.