


Table 2.18  
All Cancer Sites (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates by State, All Races, 2012-2016

Males and Females

State	Rate	SE	Rank(CI) <sup>d</sup>	PD
TOTAL U.S.	160.96	0.10		
High Five States				
Kentucky	197.93	0.89	01(01,01)	22.97 <sup>c</sup>
Mississippi	193.62	1.09	02(02,03)	20.29 <sup>c</sup>
West Virginia	189.41	1.25	03(02,04)	17.67 <sup>c</sup>
Arkansas	185.17	1.03	04(03,07)	15.04 <sup>c</sup>
Tennessee	183.95	0.70	05(04,07)	14.28 <sup>c</sup>
Low Five States				
Arizona	144.08	0.61	47(44,48)	-10.49 <sup>c</sup>
New Mexico	143.99	1.11	48(44,48)	-10.54 <sup>c</sup>
Colorado	138.87	0.73	49(48,49)	-13.72 <sup>c</sup>
Hawaii	133.13	1.25	50(50,50)	-17.29 <sup>c</sup>
Utah	127.01	1.05	51(51,51)	-21.09 <sup>c</sup>

State	Rate	SE	Rank(CI) <sup>d</sup>	PD	State	Rate	SE	Rank(CI) <sup>d</sup>	PD
Alabama	179.00	0.80	08(08,10)	11.21 <sup>c</sup>	Montana	154.97	1.57	36(32,43)	-3.72
Alaska	164.85	2.56	22(13,34)	2.42	Nebraska	159.29	1.23	33(26,35)	-1.04
Arizona	144.08	0.61	47(44,48)	-10.49 <sup>c</sup>	Nevada	162.29	1.06	28(22,33)	0.83
Arkansas	185.17	1.03	04(03,07)	15.04 <sup>c</sup>	New Hampshire	162.04	1.43	29(20,33)	0.67
California	145.27	0.27	45(44,48)	-9.75 <sup>c</sup>	New Jersey	153.50	0.55	39(36,42)	-4.63
Colorado	138.87	0.73	49(48,49)	-13.72 <sup>c</sup>	New Mexico	143.99	1.11	48(44,48)	-10.54 <sup>c</sup>
Connecticut	145.81	0.82	44(43,48)	-9.41 <sup>c</sup>	New York	150.96	0.36	42(41,43)	-6.21 <sup>c</sup>
Delaware	169.57	1.73	15(12,24)	5.35 <sup>c</sup>	North Carolina	166.90	0.55	19(17,23)	3.69
D.C.	169.04	2.35	17(11,27)	5.02 <sup>c</sup>	North Dakota	149.80	1.92	43(36,46)	-6.93 <sup>c</sup>
Florida	152.56	0.33	41(38,42)	-5.22 <sup>c</sup>	Ohio	177.12	0.51	10(08,11)	10.04 <sup>c</sup>
Georgia	165.57	0.59	21(18,25)	2.86	Oklahoma	183.69	0.93	06(04,07)	14.12 <sup>c</sup>
Hawaii	133.13	1.25	50(50,50)	-17.29 <sup>c</sup>	Oregon	162.79	0.84	26(22,32)	1.14
Idaho	154.81	1.34	37(33,42)	-3.82	Pennsylvania	169.04	0.46	16(14,19)	5.02 <sup>c</sup>
Illinois	168.48	0.49	18(15,20)	4.67	Rhode Island	164.75	1.61	24(16,32)	2.35
Indiana	178.39	0.70	09(08,11)	10.83 <sup>c</sup>	South Carolina	171.81	0.79	13(12,16)	6.74 <sup>c</sup>
Iowa	165.63	0.94	20(17,27)	2.90	South Dakota	160.46	1.81	31(21,36)	-0.31
Kansas	164.24	1.01	25(19,30)	2.04	Tennessee	183.95	0.70	05(04,07)	14.28 <sup>c</sup>
Kentucky	197.93	0.89	01(01,01)	22.97 <sup>c</sup>	Texas	154.01	0.36	38(36,40)	-4.32
Louisiana	183.02	0.86	07(04,07)	13.71 <sup>c</sup>	Utah	127.01	1.05	51(51,51)	-21.09 <sup>c</sup>
Maine	174.06	1.39	12(09,15)	8.14 <sup>c</sup>	Vermont	164.82	2.06	23(14,33)	2.40
Maryland	160.28	0.71	32(27,33)	-0.42	Virginia	160.84	0.60	30(26,33)	-0.07
Massachusetts	155.82	0.63	35(33,38)	-3.19	Washington	157.09	0.65	34(32,37)	-2.40
Michigan	170.42	0.54	14(13,17)	5.88 <sup>c</sup>	West Virginia	189.41	1.25	03(02,04)	17.67 <sup>c</sup>
Minnesota	153.44	0.71	40(36,42)	-4.67	Wisconsin	162.77	0.69	27(22,31)	1.12
Mississippi	193.62	1.09	02(02,03)	20.29 <sup>c</sup>	Wyoming	144.66	2.16	46(43,49)	-10.12 <sup>c</sup>
Missouri	175.59	0.70	11(10,12)	9.09 <sup>c</sup>					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).

<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 5% or more.

<sup>d</sup> Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.