


Table 15.25
Cancer of the Lung and Bronchus (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2012-2016

Males

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	51.64	0.08		
High Five States				
Kentucky	84.55	0.88	01(01,01)	63.73 ^c
Mississippi	78.59	1.06	02(02,03)	52.19 ^c
Arkansas	75.70	0.98	03(02,05)	46.58 ^c
Tennessee	73.31	0.67	04(03,06)	41.95 ^c
West Virginia	72.53	1.15	05(03,06)	40.45 ^c
Low Five States				
Wyoming	37.21	1.62	47(42,50)	-27.95 ^c
California	36.47	0.21	48(47,49)	-29.39 ^c
New Mexico	35.22	0.81	49(47,50)	-31.80 ^c
Colorado	32.52	0.54	50(49,50)	-37.03 ^c
Utah	23.45	0.67	51(51,51)	-54.60 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	70.90	0.76	06(04,06)	37.28 ^c	Montana	41.87	1.20	42(38,47)	-18.92 ^c
Alaska	50.31	2.13	26(17,38)	-2.58	Nebraska	50.49	1.04	25(22,34)	-2.24
Arizona	40.06	0.47	45(42,47)	-22.42 ^c	Nevada	47.57	0.84	33(27,37)	-7.89 ^c
Arkansas	75.70	0.98	03(02,05)	46.58 ^c	New Hampshire	50.22	1.18	27(22,35)	-2.76
California	36.47	0.21	48(47,49)	-29.39 ^c	New Jersey	43.37	0.45	40(38,43)	-16.02 ^c
Colorado	32.52	0.54	50(49,50)	-37.03 ^c	New Mexico	35.22	0.81	49(47,50)	-31.80 ^c
Connecticut	41.52	0.67	43(40,46)	-19.61 ^c	New York	45.11	0.31	38(35,40)	-12.65 ^c
Delaware	57.00	1.51	16(14,23)	10.38 ^c	North Carolina	62.53	0.51	12(11,14)	21.09 ^c
D.C.	43.11	1.84	41(31,47)	-16.52 ^c	North Dakota	47.37	1.61	35(25,41)	-8.26 ^c
Florida	49.79	0.28	30(25,31)	-3.59	Ohio	62.72	0.46	11(11,14)	21.46 ^c
Georgia	59.80	0.55	15(14,16)	15.79 ^c	Oklahoma	67.20	0.84	08(07,10)	30.12 ^c
Hawaii	39.13	1.01	46(42,48)	-24.23 ^c	Oregon	46.49	0.67	36(30,39)	-9.97 ^c
Idaho	40.36	1.01	44(41,47)	-21.84 ^c	Pennsylvania	55.23	0.39	20(17,22)	6.96 ^c
Illinois	54.81	0.43	21(18,23)	6.13 ^c	Rhode Island	56.38	1.44	18(15,24)	9.17 ^c
Indiana	66.66	0.65	09(07,10)	29.09 ^c	South Carolina	61.92	0.71	13(11,15)	19.90 ^c
Iowa	55.93	0.82	19(16,22)	8.30 ^c	South Dakota	52.40	1.53	24(17,32)	1.48
Kansas	53.41	0.86	22(18,26)	3.42	Tennessee	73.31	0.67	04(03,06)	41.95 ^c
Kentucky	84.55	0.88	01(01,01)	63.73 ^c	Texas	47.62	0.30	32(30,36)	-7.79 ^c
Louisiana	67.44	0.80	07(07,10)	30.60 ^c	Utah	23.45	0.67	51(51,51)	-54.60 ^c
Maine	61.72	1.23	14(10,16)	19.51 ^c	Vermont	50.01	1.68	28(20,37)	-3.15
Maryland	48.50	0.60	31(26,36)	-6.09 ^c	Virginia	52.96	0.53	23(21,26)	2.55
Massachusetts	47.45	0.53	34(29,37)	-8.12 ^c	Washington	45.16	0.52	37(35,40)	-12.56 ^c
Michigan	56.47	0.47	17(16,20)	9.35 ^c	West Virginia	72.53	1.15	05(03,06)	40.45 ^c
Minnesota	44.12	0.57	39(36,42)	-14.56 ^c	Wisconsin	49.91	0.58	29(24,32)	-3.35
Mississippi	78.59	1.06	02(02,03)	52.19 ^c	Wyoming	37.21	1.62	47(42,50)	-27.95 ^c
Missouri	65.04	0.64	10(08,11)	25.95 ^c					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 5% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.