

Table 18.10
MyelomaSEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Males

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
Survival Time																									
1-year	67.8	69.6	73.2	72.6	74.3	71.0	74.1	76.3	75.8	73.7	74.4	74.3	77.0	76.9	76.4	73.9	77.7	76.9	83.9	83.2	80.3	80.4	82.1	84.8	82.4
2-year	52.3	54.7	59.3	58.5	59.8	58.1	61.1	63.7	61.5	61.8	61.8	61.6	65.4	68.3	64.5	67.1	68.3	68.0	74.2	73.9	72.7	71.3	73.0	76.8	
3-year	39.0	42.8	46.2	46.7	49.3	44.8	50.4	53.6	50.9	52.6	51.2	52.7	56.4	60.6	57.0	58.6	60.3	59.5	66.9	66.6	64.6	63.8	66.8		
4-year	30.8	33.6	36.6	37.1	41.3	39.6	40.9	43.8	42.9	43.4	46.3	46.9	47.9	51.9	50.9	52.2	52.3	52.9	61.5	59.3	57.4	58.7			
5-year	24.3	25.8	28.3	30.4	36.3	33.5	31.6	38.2	34.8	36.0	40.2	41.8	42.4	46.5	47.4	47.9	48.1	47.7	53.5	53.7	51.9				
6-year	19.2	20.8	23.5	25.3	31.8	29.5	26.5	34.1	30.0	30.2	35.3	36.6	37.9	40.8	41.9	43.5	43.1	41.3	48.4	50.3					
7-year	15.4	16.9	19.2	21.4	26.1	25.5	22.5	28.2	26.1	26.3	31.6	33.2	34.6	36.6	36.5	38.1	39.0	39.3	44.0						
8-year	13.0	13.8	16.2	18.7	22.2	23.7	20.2	25.8	21.7	23.6	27.7	29.9	31.0	33.4	34.4	34.4	36.5	35.8							
9-year	11.0	11.7	13.0	16.8	20.2	20.9	18.3	23.6	19.5	20.0	24.9	26.9	28.7	31.5	31.8	32.1	33.3								
10-year	9.2	9.9	11.2	14.9	19.2	18.0	16.5	22.0	17.6	18.8	22.3	24.3	24.1	29.0	28.5	29.4									
11-year	7.5	8.9	9.9	13.8	17.6	16.7	15.0	21.1	14.5	17.4	21.1	21.7	21.8	27.0	25.0										
12-year	6.9	7.7	9.4	12.2	15.7	16.3	14.5	19.1	13.8	15.8	19.9	21.0	20.7	25.8											
13-year	5.8	6.7	8.6	11.4	15.1	14.2	13.2	17.6	13.1	14.9	17.9	20.1	19.1												
14-year	5.2	6.1	8.0	10.6	13.5	13.8	12.8	17.2	11.5	14.2	17.5	18.8													
15-year	4.6	5.7	7.1	10.2	13.3	12.4	12.3	16.1	11.1	13.8	15.9														
16-year	4.4	5.2	6.8	9.8	11.6	9.5	11.8	14.7	10.2	13.8															
17-year	3.9	4.8	6.4	9.1	10.8	9.1	10.9	14.3	10.1																
18-year	3.5	4.4	6.2	8.4	10.5	9.1	9.1	13.4																	
19-year	3.3	4.2	6.1	7.5	9.5	8.7	9.0																		
20-year	2.7	3.9	6.0	7.3	8.2	7.4																			

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). [Expected survival rates](#) are derived from the U.S. Annual Life Tables.