Table 19.3 Non-Hodgkin Lymphoma

Trends in SEER Incidence^{ab} and U.S. Mortality^c Using the Joinpoint Regression Program, 1975-2016 With up to Five Joinpoints, 2000-2016 With up to Three Joinpoints, Females by Race/Ethnicity

| | JP Trend 1 | | JP Trend 2 | | JP Trend 3 | | JP Trend 4 | | JP Trend 5 | | JP Trend 6 | | AAPC ^d | |
|---|------------|-------|------------|-------|------------|-------|------------|-------|------------|-----|------------|-----|-------------------|---------|
| | Years | APC | Years | APC | Years | APC | Years | APC | Years | APC | Years | APC | 2007-16 | 2012-16 |
| SEER 9 Delay-Adjusted Incidence ^a , 1975-2016 | | | | | | | | | | | | | | |
| All Races | 1975-88 | 3.0* | 1988-04 | 1.4* | 2004-16 | -0.7* | | | | | | | -0.7* | -0.7* |
| White | 1975-88 | 3.1* | 1988-04 | 1.4* | 2004-16 | -0.6* | | | | | | | -0.6* | -0.6* |
| Black | 1975-04 | 2.9* | 2004-16 | -0.6 | | | | | | | | | -0.6 | -0.6 |
| SEER 21 Delay-Adjusted Incidence ^b , 2000-2016 | | | | | | | | | | | | | | |
| All Races | 2000-04 | 1.3* | 2004-16 | -0.4* | | | | | | | | | -0.4* | -0.4* |
| White | 2000-06 | 0.8* | 2006-16 | -0.5* | | | | | | | | | -0.5* | -0.5* |
| White NH ^{ef} | 2000-06 | 0.8* | 2006-16 | -0.5* | | | | | | | | | -0.5* | -0.5* |
| Black | 2000-16 | 0.5* | | | | | | | | | | | 0.5* | 0.5* |
| Black NH ^{ef} | 2000-16 | 0.5* | | | | | | | | | | | 0.5* | 0.5* |
| API ^e | 2000-16 | 0.3 | | | | | | | | | | | 0.3 | 0.3 |
| AI/AN ^{eg} | 2000-16 | 0.7 | | | | | | | | | | | 0.7 | 0.7 |
| $	ext{Hispanic}^{	ext{f}}$ | 2000-16 | 0.4* | | | | | | | | | | | 0.4* | 0.4* |
| U.S. Cancer Mortal | | | | | | | | | | | | | | |
| All Races | 1975-94 | 2.2* | 1994-97 | 1.0 | 1997-08 | -3.3* | 2008-16 | -2.5* | | | | | -2.6* | -2.5* |
| White | 1975-97 | 2.1* | 1997-04 | -3.8* | 2004-16 | -2.7* | | | | | | | -2.7* | -2.7* |
| Black | 1975-96 | 2.7* | 1996-16 | -2.2* | | | | | | | | | -2.2* | -2.2* |
| U.S. Cancer Mortality ^c , 2000-2016 | | | | | | | | | | | | | | |
| All Races | 2000-04 | -3.9* | 2004-16 | -2.7* | | | | | | | | | -2.7* | -2.7* |
| White | 2000-03 | -4.4* | 2003-16 | -2.7* | | | | | | | | | -2.7* | -2.7* |
| White NH ^{ef} | 2000-03 | -4.5* | 2003-16 | -2.7* | | | | | | | | | -2.7* | -2.7* |
| Black | 2000-16 | -2.1* | | | | | | | | | | | -2.1* | -2.1* |
| Black NH ^{ef} | 2000-16 | -2.0* | | | | | | | | | | | -2.0* | -2.0* |
| API ^e | 2000-16 | -1.9* | | | | | | | | | | | -1.9* | -1.9* |
| AI/AN ^{eg} | 2000-16 | -3.7* | | | | | | | | | | | -3.7* | -3.7* |
| Hispanic ^f | 2000-16 | -2.1* | | | | | | | | | | | -2.1* | -2.1* |

Joinpoint Regression Program Version 4.7, February 2019, National Cancer Institute. (https://surveillance.cancer.gov/joinpoint/). The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Trends are from the SEER 21 areas (SEER 9 areas, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding

SF/SJM/LA, Kentucky, Louisiana, New Jersey, Georgia excluding ATL/RG, Idaho, New York and Massachusetts).

Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

d The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic

Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

Data for American Indian/Alaska Native are based on the Purchased/Referred Care Delivery Area (PRCDA) counties.

The APC/AAPC is significantly different from zero (p<.05).

Joinpoint regression line analysis could not be performed on data series.